

Distribution Modeling for Colorado SWAP Plants of Greatest Conservation Need Interim Report- April 2022



WARNER COLLEGE
OF NATURAL RESOURCES
COLORADO STATE UNIVERSITY

CNHP's mission is to advance the conservation of Colorado's native species and ecosystems through science, planning, and education for the benefit of current and future generations.

Colorado Natural Heritage Program

Warner College of Natural Resources
Colorado State University
1475 Campus Delivery
Fort Collins, CO 80523

Recommended Citation:

Decker, K., J.P. Smith, J. Handwerk, G. Doyle, and S. Panjabi. 2022. Distribution Modeling for Colorado SWAP Plants of Greatest Conservation Need. Colorado Natural Heritage Program, Colorado State University, Fort Collins, Colorado.

Cover photo: *Penstemon acaulis* var. *yampaensis* in Moffat County, Colorado. © 2020, Jessica Smith.

Contents

Contents	3
Acknowledgement	3
Introduction	4
Background	4
Objectives.....	4
Methods	8
Occurrence update and review.....	8
Modeling process.....	8
Model review and threshold setting.....	10
Post-review processing	11
Results	11
Individual species results	11
Discussion	40
Coverage and use of available species models.....	40
Additional modeling needs	41
Future data development	43
Conclusion.....	46
References	48
Appendix A: Data sources	50
Appendix B: Basic modeling methods	77
Appendix C: Expert review of SDM for rare plant species	80
Introduction	80
Methods.....	80
Results.....	82
Discussion.....	83
References	87
Appendix D: Projected suitable habitat under potential future climate conditions	89

Acknowledgement

The authors would like to thank the Colorado Natural Areas Program (CNAP)/Colorado Parks & Wildlife (CPW) for financial support of this project. Special thanks to CNAP Program Coordinator, Raquel Wertsbaugh, for her support of the project and CNHP staff, Amy Greenwell, for her GIS expertise and coordination with CODEX.

Introduction

Background

The revised 2015 Colorado State Wildlife Action Plan (SWAP) includes 117 Plant Species of Greatest Conservation Need (PGCN) in the Rare Plant Addendum (Colorado Parks and Wildlife 2015). These species are ranked globally critically imperiled (G1) or imperiled (G2), at risk throughout their range and under threat of extinction. Known locations of these species are recorded in a statewide geospatial database maintained by the Colorado Natural Heritage Program (CNHP). However, surveys for these species have not been comprehensive, and therefore information on their distribution is incomplete. For this project, models of species distribution were created using information on known locations and species habitat requirements. These distribution models will be included in the conservation data sharing platform, the Colorado Conservation Data Explorer (CODEX), in order to help conserve and protect these PGCN through environmental review and conservation planning. The following objectives were met in this project:

Objectives

- 1) Create distribution models for 80 of the 117 plant species listed in the Rare Plant Addendum of the SWAP (Table 1). Species included in this project were prioritized by species conservation need and development threats. Whenever possible, distribution models were produced as probability surfaces. These raster digital datasets can be converted to other formats as needed, although with some loss of information.
- 2) Produce binary versions of the distribution models for inclusion in the Colorado Conservation Data Explorer (CODEX), released in 2021 and hosted by CNHP, in order for PGCN to be considered in statewide conservation planning and environmental review.

Table 1. PGCN species modeled. Common names are those used in Colorado.

Scientific Name	Common Name	Species Priority	Global & State Status Ranks	Federal Agency Status	% of Range in Colorado	Priority
<i>Aliciella sedifolia</i>	Stonecrop gilia	Tier 1	G1 / S1	USFS	Endemic	Highest - Tier 1
<i>Ipomopsis ramosa</i>	Coral ipomopsis	Tier 1	G1 / S1		Endemic	Highest - Tier 1
<i>Lepidium huberi</i>	Huber's pepperwort	Tier 1	G1G2 / S1S2		High	Highest - Tier 1
<i>Lygodesmia doloresensis</i>	Dolores River skeletonplant	Tier 1	G1G2 / S1S2	BLM	High	Highest - Tier 1
<i>Mimulus gemmiparus</i>	Budding monkey flower	Tier 1	G1 / S1	USFS	Endemic	Highest - Tier 1
<i>Oenothera coloradensis</i> ssp. <i>coloradensis</i>	Colorado butterfly plant	Tier 1	G3T2 / S1	LT	Medium	Highest - Tier 1
<i>Packera mancosana</i>	Mancos shale packera	Tier 1	G1 / S1		Endemic	Highest - Tier 1
<i>Pediocactus knowltonii</i>	Knowlton cactus	Tier 1	G1 / SNA	LE	Historical	Highest - Tier 1
<i>Penstemon gibbensii</i>	Gibben's beardtongue	Tier 1	G1G2 / S1	BLM	High	Highest - Tier 1
<i>Penstemon scariosus</i> var. <i>albifluvis</i>	White River penstemon	Tier 1	G4T1 / S1	BLM	Low	Highest - Tier 1
<i>Phacelia gina-glenneae</i>	Troublesome phacelia	Tier 1	G1/S1		Endemic	Highest - Tier 1
<i>Physaria rollinsii</i>	Rollins twinpod	Tier 1	G1 / S1		Endemic	Highest - Tier 1
<i>Physaria scrotiformis</i>	West Silver bladderpod	Tier 1	G1 / S1		Endemic	Highest - Tier 1
<i>Asclepias uncialis</i> ssp. <i>uncialis</i>	Dwarf milkweed	Tier 2	G3G4T2T3 / S2	BLM/USFS	Very High	High
<i>Astragalus sparsiflorus</i>	Front Range milkvetch	Tier 2	G2 / S2		Endemic	High
<i>Mentzelia rhizomata</i>	Roan Cliffs blazing star	Tier 2	G2 / S2	BLM	Endemic	High
<i>Nuttallia chrysantha</i>	Golden blazing star	Tier 2	G2 / S2	BLM	Endemic	High
<i>Oonopsis puebloensis</i>	Pueblo goldenweed	Tier 2	G2 / S2		Endemic	High
<i>Oxybaphus rotundifolius</i>	Round-leaf four o'clock	Tier 2	G2 / S2		Endemic	High
<i>Thalictrum heliophilum</i>	Sun-loving meadow rue	Tier 2	G2 / S2	BLM/USFS	Endemic	High
<i>Aletes humilis</i>	Larimer aletes	Tier 2	G2G3 / S2S3		Endemic	Medium
<i>Astragalus rafaelensis</i>	San Rafael milkvetch	Tier 2	G2G3 / S1	BLM	High	Medium
<i>Camissonia eastwoodiae</i>	Eastwood evening primrose	Tier 2	G2 / S1	BLM	Medium	Medium
<i>Castilleja puberula</i>	Downy Indian-paintbrush	Tier 2	G2G3 / S2S3		Endemic	Medium
<i>Cleome multicaulis</i>	Slender spiderflower	Tier 2	G2G3 / S2S3	BLM	High	Medium

Scientific Name	Common Name	Species Priority	Global & State Status Ranks	Federal Agency Status	% of Range in Colorado	Priority
<i>Draba smithii</i>	Smith whitlow-grass	Tier 2	G2 / S2	USFS	Endemic	Medium
<i>Frasera coloradensis</i>	Colorado green gentian	Tier 2	G2G3 / S2S3		Endemic	Medium
<i>Herrickia horrida</i>	Canadian River spiny aster	Tier 2	G2? / S1		Medium	Medium
<i>Ipomopsis globularis</i>	Globe gilia	Tier 2	G2 / S2	USFS	Endemic	Medium
<i>Lupinus crassus</i>	Payson lupine	Tier 2	G2 / S2	BLM	Endemic	Medium
<i>Nuttallia densa</i>	Arkansas Canyon stickleaf	Tier 2	G2 / S2	BLM	Endemic	Medium
<i>Oenothera acutissima</i>	Narrow-leaf evening primrose	Tier 2	G2 / S2	BLM	Medium	Medium
<i>Oonopsis foliosa</i> var. <i>monocephala</i>	Rayless goldenweed	Tier 2	G3G4T2 / S2		Endemic	Medium
<i>Oreocarya revealii</i>	Gypsum Valley cat's-eye	Tier 2	G2 / S2	BLM	Endemic	Medium
<i>Oxytropis besseyi</i> var. <i>obnapiformis</i>	Bessey locoweed	Tier 2	G5T2 / S2		Very High	Medium
<i>Penstemon acaulis</i> var. <i>yampaensis</i>	Yampa beardtongue	Tier 2	G3T2 / S2		High	Medium
<i>Penstemon degeneri</i>	Degener beardtongue	Tier 2	G2 / S2	BLM/USFS	Endemic	Medium
<i>Penstemon fremontii</i> var. <i>glabrescens</i>	Fremont's beardtongue	Tier 2	G3G4T2 / S2		Endemic	Medium
<i>Penstemon scariosus</i> var. <i>cyanomontanus</i>	Plateau penstemon	Tier 2	G4T2 / S2		High	Medium
<i>Physaria bellii</i>	Bell's twinpod	Tier 2	G2G3 / S2S3		Endemic	Medium
<i>Physaria parviflora</i>	Piceance bladderpod	Tier 2	G2 / S2	BLM	Endemic	Medium
<i>Physaria vicina</i>	Good-neighbor bladderpod	Tier 2	G2 / S2	BLM	Endemic	Medium
<i>Ptilagrostis porteri</i>	Porter feathergrass	Tier 2	G2 / S2	USFS	Endemic	Medium
<i>Puccinellia parishii</i>	Parish's alkali grass	Tier 2	G2G3 / S1		Low	Medium
<i>Townsendia glabella</i>	Gray's townsend-daisy	Tier 2	G2 / S2		Endemic	Medium
<i>Eriogonum coloradense</i>	Colorado wild buckwheat	Tier 2	G2 / S2	BLM	Endemic	Low
<i>Lepidium crenatum</i>	Alkaline pepperwort	Tier 2	G2 / S2		Medium	Low
<i>Lomatium concinnum</i>	Colorado desert-parsley	Tier 2	G2G3 / S2S3	BLM	Endemic	Low
<i>Penstemon mensarum</i>	Grand Mesa penstemon	Tier 2	G2 / S2		Endemic	Low
<i>Physaria alpina</i>	Avery Peak twinpod	Tier 2	G2 / S2		Endemic	Low
<i>Physaria pruinosa</i>	Pagosa bladderpod	Tier 2	G2 / S2	BLM/USFS	Endemic	Low
<i>Thelypodiopsis juniperorum</i>	Juniper tumble mustard	Tier 2	G2 / S2		Endemic	Low

Scientific Name	Common Name	Species Priority	Global & State Status Ranks	Federal Agency Status	% of Range in Colorado	Priority
<i>Aletes macdougallii</i> ssp. <i>breviradiatus</i>	Mesa Verde aletes	Tier 2	G3T2T3 / S1		Medium	Lowest
<i>Anticlea vaginatus</i>	Alcove death camas	Tier 2	G2 / S2		Low	Lowest
<i>Astragalus cronquistii</i>	Cronquist milkvetch	Tier 2	G2 / S2		High	Lowest
<i>Astragalus equisolensis</i>	Horseshoe milkvetch	Tier 2	G5T1 / S1	BLM	Low	Lowest
<i>Astragalus iodopetalus</i>	Violet milkvetch	Tier 2	G2 / S1	USFS	Medium	Lowest
<i>Astragalus missouriensis</i> var. <i>humistratus</i>	Missouri milkvetch	Tier 2	G5T1 / S1	USFS	Endemic	Lowest
<i>Astragalus naturitensis</i>	Naturita milkvetch	Tier 2	G2G3 / S2S3	BLM	High	Lowest
<i>Astragalus piscator</i>	Fisher Towers milkvetch	Tier 2	G2G3 / S1	BLM	Low	Lowest
<i>Boechnera crandallii</i>	Crandall's rock-cress	Tier 2	G2 / S2	BLM	High	Lowest
<i>Calochortus ciscoensis</i>	Cisco sego lily	Tier 2	G2 / S1		Low	Lowest
<i>Cirsium perplexans</i>	Adobe thistle	Tier 2	G2G3 / S2S3		Endemic	Lowest
<i>Delphinium ramosum</i> var. <i>alpestre</i>	Colorado larkspur	Tier 2	G4T2 / S2		High	Lowest
<i>Delphinium robustum</i>	Wahatoya Creek larkspur	Tier 2	G2? / S2?		Medium	Lowest
<i>Draba graminea</i>	San Juan whitlow-grass	Tier 2	G2 / S2		Endemic	Lowest
<i>Erigeron kachinensis</i>	Kachina daisy	Tier 2	G2 / S1	BLM	Low	Lowest
<i>Eriogonum clavellatum</i>	Comb Wash buckwheat	Tier 2	G2 / S1	BLM	Medium	Lowest
<i>Limnorchis zothecina</i>	Alcove bog orchid	Tier 2	G2 / S1		Low	Lowest
<i>Mentzelia paradoxensis</i>	Paradox stickleaf	Tier 2	G2? / S2?		Endemic	Lowest
<i>Mertensia humilis</i>	Rocky Mountain bluebells	Tier 2	G2 / S1		Medium	Lowest
<i>Oreocarya osterhoutii</i>	Osterhout cat's-eye	Tier 2	G2G3 / S2	BLM	Low	Lowest
<i>Salix arizonica</i>	Arizona willow	Tier 2	G2G3 / S1	USFS	Low	Lowest
<i>Thelypodium paniculatum</i>	Northwestern thelypody	Tier 2	G2 / SH		Low	Lowest
<i>Townsendia fendleri</i>	Fendler's townsend-daisy	Tier 2	G2 / S2		High	Lowest
<i>Trifolium dasyphyllum</i> ssp. <i>anemophilum</i>	Whip-root clover	Tier 2	G5T2? / S1		Low	Lowest
<i>Eriogonum brandegeei</i>	Brandegeee wild buckwheat	Tier 1	G1G2 / S1S2	BLM/USFS	Endemic	New model
<i>Eriogonum pelinophilum</i>	Clay-loving wild buckwheat	Tier 1	G2 / S2	LE	Endemic	New model
<i>Potentilla rupincola</i>	Rocky Mountain cinquefoil	Tier 2	G2 / S2	USFS	Endemic	New model
<i>Telesonix jamesii</i>	James telesonix	Tier 2	G2 / S2		Very High	New model

Methods

Occurrence update and review

Element occurrences for the 80 species were updated prior to the modeling effort. The primary data source searched for new information was SEINet, the online herbarium database (SEINet 2021). SEINet records were compared to existing CNHP BIOTICS database records (CNHP 2021-2022) and SEINet specimens representing new locations were mapped as new or updated element occurrences in the CNHP database. All SEINet records dated 2000 and newer were mapped as well as selected older SEINet records that represented range extensions. In addition, all backlog data from CNHP files for the species of interest were incorporated into the CNHP database. A total of 265 new or updated Element Occurrence (EO) records were produced during the project.

A shapefile of occurrence polygons for each species was exported from BIOTICS. Using ArcGIS 10.4 (ESRI 1999-2015), this multipart shapefile was converted to a single-part shapefile, separating polygons belonging to the same EO into individual features. These polygon features were converted to centroid points, with the constraint that the point fall within the polygon. Coordinates (XY in UTM NAD83 zone 13) were added to the points and used to produce the location input csv file for each species. Older historic records with poor location precision were sometimes omitted from the modeling dataset but retained for model review. In a few cases, additional points were added to very large polygons, using a 500m square net to ensure that added points were not exactly duplicating environmental locations. In contrast, for a few species that had many mapped polygons within a smaller well-surveyed area, a randomly selected subset of points in these areas was used in the model to decrease the influence of one smaller area on the predicted range as a whole.

Modeling process

Species distribution models (also called environmental/ecological niche models, or predictive habitat models), are based on the premise of finding places on the landscape where environmental conditions (climate, soils, exposure, etc.) are similar to conditions at documented locations of the species of interest. This can be as simple as extracting environmental covariate values for known points and using them to select portions of spatial datasets that match those values or value ranges (deductive modeling) or can involve using complex algorithms that compute an approximate probability that a species could occur at a particular point (inductive modeling). Distribution models can, but do not typically, take into account the biogeographic history of a species, or its ability to disperse to new areas or tolerate novel conditions.

For this project at least one inductive model using the maximum entropy (Maxent) modeling procedure (Phillips et al. 2004, 2006) was produced for each species with more than two occurrence records. This procedure is particularly useful for modeling species where absence data is lacking. We used the Maxent version 3.4.3 java-based software (Phillips et al. 2020). Maxent has been widely used in species distribution modeling and performs well in comparison with other methods (Elith et al. 2011). This procedure uses the environmental covariate values from occurrence points plus 10,000 randomly selected background points to estimate a probability distribution that is consistent with data from known locations. This estimate is as close as possible (has maximum entropy) to the estimate from the background data (the null model), since, without any data, we would have no reason to think that the

species would be more likely to be in one location than any other. Species distribution is estimated by minimizing the distance between the occupied and background, subject to constraining the means of estimated occupied factors to be close to observed means. Constraints ensure that the mean for a variable in the estimated distribution is close to the mean across the locations with occurrences. The raw solution is transformed to complementary log-log (cloglog) output with a potential range of 0 to 1, becoming more-or-less a probability estimate of occurrence.

In addition to the spatial location of known occurrences, inputs for the model are generally data matrices as raster digital data representing the value of an environmental factor for every cell across the entire study area. The Maxent software requires that environmental factors be in ASCII grid format, and all grids must share a common spatial reference, extent, cell size, and alignment, and be in the same folder. Environmental inputs were produced in ArcGIS using a 30m resolution digital elevation model (DEM) raster with a rectangular extent covering the state of Colorado plus a buffer of approximately 8km on each edge of the boundary as the reference extent to which all other rasters were aligned. All data used the NAD 1983 UTM Zone 13N spatial projection. Input rasters were produced as geotiffs, then converted to ASCII and stored together. In a few instances values for a particular environmental factor were not available across the entire study area. If, however, these areas of “no data” fell outside the reasonable expected range of a species, the raster was used anyway. The resulting slight reduction in background point data available for that factor was a reasonable tradeoff for the potential contribution of additional environmental information.

For each species, habitat description information was extracted from individual element occurrence records in BIOTICS (typically in the General Description field, but sometimes useful information was in additional fields) and compiled in a spreadsheet. This information was used to identify important environmental factors such as characteristic geologic substrate, vegetation type, landform, aspect, slope, elevation and others, if known. Some environmental inputs were chosen to reflect particular documented habitat details for a species (e.g., a single geologic formation, or a habitat type), but general climatic (temperature and precipitation), soil, and topographic inputs were also used for all species (Appendix A).

Climate data for precipitation were grouped seasonally. Winter includes the months December, January, and February; spring includes March, April, and May; summer includes June, July, and August and fall includes September, October, and November. Temperature-related climate data were either based on monthly averages (monthly minimum temperatures), seasonal extremes (winter minimum or summer maximum), or growing season boundaries (first and last frost dates, or total number of frost-days). Seasonal extreme temperatures indicate the lowest winter or highest summer temperature for a location over a 30-year normal period, not an annual average low or high.

If a species was reliably reported as being associated with a particular geologic substrate or substrates, a Euclidian distance to mapped geologic unit areas input layer was generally preferred over a categorical geology input layer. This technique compensates for the fact that geology mapping is highly inexact at the scale which matters to individual plants. Local erosive processes may also spread the appropriate substrate beyond its formation of origin. Finally, identification of geologic substrates by field botanists can be incorrect. Moreover, the continuous surface of the distance layer produces model surfaces characterized by gradual suitability changes that are more likely to reflect ecological conditions on the ground. If, however, substrate appeared to be important but not described in detail, a categorical

geology layer was used. Likewise, soil characteristics represented as continuous values (e.g., percent silt, clay, or sand), were used preferentially rather than individual soils units. In some instances, distance to a particular type of vegetation, or a categorical vegetation layer was used. A brief summary of the modeling process is found in Appendix B.

The use of statewide input layers facilitated an economy of scale because layers did not have to be adjusted to the range of each species (a potentially time-consuming process). However, one potential consequence of this practice is to introduce areas of predicted suitable habitat that are simply too far from the known distribution into the final model. The use of numerous “distance to factor” layers mitigated against this issue to some extent. Ideally, future modeling efforts will be able to develop a cost-effective method of either selecting small portions of statewide environmental datasets, or else constraining background samples to be nearer known occurrences.

Model review and threshold setting

Maximum entropy model results, in the form of the model raster image and a layer file classified into three tiers of probability, were reviewed in ArcMap for acceptable geographic extent, inclusion of element occurrence records, and overall correctness. The analysis provided in the Maxent results was also reviewed, with special attention paid to which environmental factors were the most important in creating the model.

Maxent returns a continuous probability surface of approximately 0-100% (0 to 1) likelihood that a species would be present at a location (assuming the model adequately represents required environmental conditions). Models were primarily intended to suggest a need for field survey in the sense of indicating if an area of interest to a CODEX user was likely to contain suitable habitat for a particular PGCN species. Our threshold standard is based on this use; other potential model uses (e.g., identifying critical habitat, targeted inventory, etc.) might require a different threshold standard. Our threshold decision tree is based on commonly used thresholding methods reported in Pearson (2010, Table 4).

Modeled area to be included in CODEX was determined by setting a cut-off value for the probability of species presence to return a yes or no (binary) value for potential species presence in the environmental review. The typical probability used to classify the CODEX models was 0.50, i.e., at least a 50% chance that the species would be present at the location according to the model. Thus, the top threshold level was the fixed value of probability = 0.50. If this threshold was not useful, the second threshold level considered was the value of equal training sensitivity and specificity. If neither of these thresholds was satisfactory, a method of looking for the lowest predicted value corresponding to documented occurrences was used. This last threshold was applied using a flexible “best professional judgement” method that reflected the variable quality of occurrence record mapping and the limitations of the modeling process. In general, we sought to identify a nuanced threshold that would cover portions of all high-quality occurrences while not simultaneously greatly increasing the area suggested for survey. Reasons for threshold adjustment included:

- Including all occurrences or occurrence polygons for Tier 1 species, or all high-quality occurrences for Tier 2 species, in the binary model
- Generally increasing the number and/or area of occurrences included in the binary model

- Widening the predicted area to avoid too close a fit to known occurrences, or to include at least a portion of historic or low confidence occurrences.
- Expanding the binary model to increase area represented in key habitats or edge of range, or in general for poorly known species with few occurrences

Expert review of 46 species models was provided by regional botanists, and comments and notes on revisions are included below in the Individual species results. Details and results of the review process in Appendix C.

All models were produced with a statewide extent; if the modeled range extended far outside the known species range, a decision was made on where to clip the model. Typically, models were clipped to exclude areas further than one county away from the known range or further than 30-40 miles from an occurrence record. This wide buffer accounted for uncertainties in cases where a species had few element occurrence records, limited survey attention, or a wide range of ecological conditions.

Post-review processing

Reviewed and approved models were converted to binary rasters using the Reclassify tool in ArcGIS. Cells below the cutoff value were reclassified as NoData; cells \geq threshold value were classified with a value of 1. The binary raster was then clipped as needed, as specified by the reviewer. Clipping was done using an appropriate polygon shape as the clipping geometry. The final binary models with metadata will be converted to vector format for use in CODEX. Binary and full raster models and classified vector (shapefile) model versions will be retained in CNHP botany files for use in future survey work.

Metadata was created and included with the models in GIS. Metadata includes the list of input environmental factors, and indicates which factors made a non-zero contribution to the result. Use constraints and caveats are also included.

Results

Altogether, about 200 Maxent model runs were made in order to produce the final 80 CODEX models. Twenty-six species required only a single run to produce a satisfactory model, 25 needed a second run to incorporate corrected or additional factors, and 26 species required additional runs. Three species could not be modeled using maximum entropy methods (too few documented locations) and a deductive model was constructed instead, using substrate, vegetation types, elevations, and/or climate factors matching conditions at the documented locations.

Ninety-three separate environmental input layers were produced for use in this modeling effort (Appendix A), although not all of these proved useful in final model results. The full list of important environmental variables and their relative percentage of contribution is included in the metadata for each species model.

Individual species results

Model input details and key results, including important environmental variables are summarized by species below. Common names are Colorado state common names. For more detail on model inputs and results, see the metadata for the GIS model.

Aletes humilis (Larimer aletes), Tier 2

This Colorado endemic is known from 27 locations in Larimer and Boulder counties in the northern Front Range. Habitat is primarily tied to outcrops of granitic rock in the 1400-MY age group. Spring warm-up timing, slope, and seasonal temperature extremes are also contributing factors. The modeled range generally extends west from the mountain front in Larimer County north of the Poudre River up to elevations between 7500 and 8500 feet (2290-2590 m), and from the vicinity of Glen Haven near the Big Thompson Canyon in southern Larimer County down to Left Hand Creek in central Boulder County. A threshold value of 0.5 was used for the probability in the CODEX version of the model and the full extent of the model was retained.

This model was reviewed by one expert reviewer, who ranked it as reasonable. This reviewer noted that the model included areas where they had observed the species in the field, but it was on the coarse side and should exclude Lake Estes and Halligan Reservoir. The model was revised to exclude the reservoirs.

Aletes macdougallii ssp. *breviradiatus* (Mesa Verde aletes), Tier 2

In southwestern Colorado this species is documented from five locations in Mesa Verde National Park, where it grows in crevices of the sandstone canyon walls. Habitat was limited to areas where extreme low winter temperatures do not fall below -16.6°F (-27°C). Steep slopes were an important factor. Substrates east of the Navajo Wash valley are primarily Cretaceous sandstones characteristic of the area, including Cliff House and Point Lookout sandstones, while habitat on the Sleeping Ute Mountain laccolith is on younger Laramide intrusives. Modeled habitat is largely west-facing canyon walls and similar slopes from School Section Canyon to the west rim of the larger Mesa Verde area, with additional areas of potential habitat to the south of the mesa in canyons tributary to the Mancos or San Juan rivers, and in scattered mid-elevation areas of Sleeping Ute Mountain on Ute Mountain Ute tribal lands. A threshold value of 0.5 was used for the probability in the CODEX version of the model and the full extent of the model was retained.

This model was created in Round II of the project and expert review has not yet been solicited.

Aliciella sedifolia (Stonecrop gilia), Tier 1

A Colorado endemic, this species is known from four locations in San Juan and Hinsdale counties in the San Juan Mountains. Habitats are barren alpine gravelly soils below ridgelines. Elevation (generally above 12000 ft; 3660 m) was the primary factor contributing to the model. Distance to surface geology of Tertiary volcanic tuff was also important and could explain nearly 85% of the distribution in a single factor model. Extreme maximum summer temperatures are generally not above 77°F (25°C). A threshold value of 0.5 was used for the probability in the CODEX version of the model. The modeled range extended 75 km (47 miles) east from nearest EO record, which seemed a reasonable extent for this under-surveyed species; therefore, the full extent of the model was retained.

This model was reviewed by one expert reviewer, who ranked it as poor. Field observations by this reviewer point to the species being restricted to areas of fine-grained texture of more widespread volcanic soils above 13,000' feet. This model should be considered for eventual revision.

Anticlea vaginatus (Alcove death camas), Tier 2

This is a species of canyon wall alcove habitats, found where perennial seeps supply pocket wetland habitats in otherwise dry sandstone dominated cliffs. Seven Colorado occurrences have been documented from the Yampa River canyon in Dinosaur National Monument in western Moffat County. Modeled habitat was constrained to canyon wall areas by using distance to steep slopes (≥ 30 degrees) factor. This habitat is among the driest in Colorado, with annual precipitation less than 12 inches (30 cm), which was an important contributing factor in the model, along with the greater frost-free period characteristic of lower elevations. A threshold value of 0.5 was used for the probability in the CODEX version of the model. Modeled habitat was clipped to constrain predicted suitable areas to the Yampa River canyon from the vicinity of Schoonover Buttes west to the Utah border, and the Green River south of the Gates of Lodore.

This model was created in Round II of the project and expert review has not yet been solicited.

Asclepias uncialis ssp. *uncialis* (Dwarf milkweed), Tier 2

In Colorado, this species is found on a variety of soil types and microsites, generally associated with grasslands. The large but sparsely populated range and lack of obvious narrow environmental influence on this diminutive, early flowering species make it extremely challenging to model. Furthermore, much of the original species' habitat has probably been converted to agricultural use, causing occurrences in the northern portion of the Colorado range to appear as outliers in the species' environmental niche. Numerous Maxent model runs with different inputs were made, but as they appeared to converge on a common solution, the version that included more of the northern habitat was selected for comparison with a deductive model. Ultimately, the Maxent model was chosen, as it captured a greater number of highly ranked EOs. Important environmental factors included distance to shortgrass prairie and soil depth. A lower probability cut-off of 0.107 was used to include modeled habitat in the northeast corner of the state, and the full extent of the model was retained.

This model was reviewed by two expert reviewers. Two reviewers responded with comments but did not return a survey form. They agreed the model looked reasonable. One reviewer believed the distribution of the model corresponded well with their field observations. The other thought the distribution may extend too far to the west and wondered if a different environmental factor might show a different projection.

Astragalus cronquistii (Cronquist milkvetch), Tier 2

In Colorado this species is primarily known from Ute Mountain Ute tribal lands in southwestern Montezuma County; it also occurs in Utah. Nearly all the 11 Colorado occurrences have not been observed for a couple of decades or more. Predicted suitable habitat was limited to areas where extreme low winter temperatures do not fall below -15°F (-26°C). Distance to substrates derived from Mancos Shale was an important factor. Seasonal precipitation patterns also contributed to constrain predicted suitable habitat to arid area; plants are apparently able to tolerate spring or summer precipitation amounts less than 2 inches (5 cm). A threshold value of 0.5 was used for the probability in the CODEX version of the model, and modeled habitat was clipped to the southwestern corner of Montezuma County. Modeled habitat included primarily alluvial fans below Sleeping Ute Mountain or the western edge of Mesa Verde where saltbush shrublands intermix with more barren areas.

This model was created in Round II of the project and expert review has not yet been solicited.

Astragalus equisolensis (Horseshoe milkvetch), Tier 2

This narrow Utah-Colorado endemic is known from nine occurrences in the Dolores River drainage in western Mesa County, which are disjunct from the Utah population. Geology was the primary factor determining potential suitable habitat; occurrences are mapped on the Moenkopi and Cutler formations as a combined unit in that area. Early spring (March average) low temperatures that are just below freezing were also a notable contributing factor, along with extreme low winter temperatures that remain above -15°F C (-26°). The suitable habitat predicted by the model aligns fairly well with the canyon slopes near the inner Dolores River canyon from Roc Creek confluence in northwestern Montrose County northwest to the Utah state line. A few scattered bands of habitat further east in Mesa County are also present. A threshold value of 0.5 was used for the probability in the CODEX version of the model and the full extent of the model was retained.

This model was created in Round II of the project and expert review has not yet been solicited.

Astragalus iodopetalus (Violet milkvetch), Tier 2

This species has a large range in southwestern Colorado, from southern Gunnison County to the New Mexico border. A total of 17 occurrences are documented; nearly all are considered historical as they have not been observed for more than 20 years. Vegetation type was the most important contributing factor, although variable from sagebrush shrublands in the north to a selection of dry woodland types such as ponderosa pine or Gambel oak in the south. Soils are characterized by higher clay and lower sand percentage. Climatic environmental factors contributing to the model are characteristic of drier valleys and foothills on north and south flanks of the San Juan Mountains. Seasonal patterns indicate a somewhat drier spring, but otherwise precipitation is more-or-less equally distributed across the seasons. Extreme winter low temperatures are -31°F (-35°C) at the highest elevations, while extreme maximum summer temperatures can reach near 100°F (37.5°C) in the warmer locations. A threshold value of 0.5 was used for the probability in the CODEX version of the model. Modeled suitable habitat was clipped to include all sides of the San Juan Mountains and includes areas of interest in eastern San Miguel County as well as the western San Luis Valley north of the Rio Grande River.

This model was created in Round II of the project and expert review has not yet been solicited.

Astragalus missouriensis var. *humistratus* (Missouri milkvetch), Tier 2

A narrow endemic known from 13 occurrences in Colorado and several in New Mexico, the species is restricted to the upper basin of the San Juan River in southwestern Colorado and northwestern New Mexico. Distance to Mancos Shale substrates was the most important contributing factor in the model. Soils have a fairly high clay content. The overall climate envelope of the distribution reflects moderately arid and warm conditions with average seasonal precipitation of around 5-6 in (12-15 cm), together with an average of 220 frost days per year. The model threshold was adjusted to 0.36 to better capture documented locations, and the model extent was clipped to exclude habitat outside of Archuleta, La Plata and small amounts of southern Hinsdale and Mineral counties.

This model was created in Round II of the project and expert review has not yet been solicited.

Astragalus naturitensis (Naturita milkvetch), Tier 2

This species has been documented from 63 occurrences in Colorado and is also found in Utah and New Mexico. Locations stretch from southern Garfield County to the Four-Corners area. Recorded habitat characteristics for the species are fairly broad, but it tends to occur on substrates primarily derived from

sandstone, which was an important contributing factor in the model. May minimum temperatures comfortably above freezing were also important. A subset of occurrences in the De Beque Canyon area were used for modeling to decrease the influence of this densely populated region in model results. The model threshold was adjusted to 0.25 to better capture documented locations, and the model extent was clipped to small exclude habitat patches on the eastern slope.

This model was created in Round II of the project and expert review has not yet been solicited.

Astragalus piscator (Fisher Towers milkvetch), Tier 2

This species is documented from four occurrences in western Mesa County, none further than 5 miles from the Utah border. Distance to substrates derived from the Permian sandstone dominated Cutler Formation were the most important contributing factor. April minimum temperatures generally not below freezing were also important, as were extreme minimum winter temperatures not below about -15°F. The model threshold was adjusted to the equal training sensitivity and specificity (0.242) to include all documented locations. Modeled suitable habitat is restricted to the Dolores River drainage in the vicinity of Gateway.

This model was created in Round II of the project and expert review has not yet been solicited.

Astragalus rafaensis (San Rafael milkvetch), Tier 2

This species is known from eastern Utah and western Colorado, where it is documented from 28 occurrences in Mesa, Delta, Montrose, and northern San Miguel counties. Habitats are generally on soils derived from Morrison Formation units; distance to this type of surface geology was the major contributing factor in the model. Precipitation in the dry season of summer was important, with a minimum requirement of generally in the range of 6-8 cm; southern occurrences receive more precipitation in comparison with northern stands. Extreme winter minimum temperatures are generally not below -22°F (-30°C). The model predicted suitable habitat in Montezuma County, but this was excluded from the final version, as the predicted habitat was over 50 miles and 2 counties away from known EOs. A probability cut-off of 0.35 was used to include additional habitat matching highly ranked EOs which were excluded at the 0.5 threshold. Predicted habitat follows the known distribution fairly closely, with the addition of habitat just downstream from the Black Canyon of the Gunnison River, where Morrison Formation units are common. The model extent was clipped to exclude areas in Montezuma County.

This model was reviewed by two expert reviewers. Two reviewers agreed the model was reasonable. One reviewer stated that the model correctly matched potential habitat but also included areas in Mesa County not currently considered potential habitat. The second reviewer also believed the model overstated potential a bit, but it did pick up all areas where surveys for the species would be recommended in their management area.

Astragalus sparsiflorus (Front Range milkvetch), Tier 2

This Colorado endemic is documented from 21 occurrences at mid-montane elevations ranging in a north-south distribution from Boulder to Custer County. Granitic substrates are common in this region. Distance to surface geology of Precambrian age metamorphic and igneous rock was the most important environmental factor in the model. These rocky soils are typically shallow. Summer precipitation generally greater than 20 cm and May minimum temperatures averaging just above freezing were also contributing factors. A threshold value of 0.5 was used for the probability in the CODEX version of the

model. The modeled area was clipped to include only areas of the Front Range and southern mountain front, from northern Larimer County to northern Huerfano County. Predicted habitat is especially prevalent at elevations of 7000-9500 feet (2130-2895 m) in the vicinity of the Platte Canyon, Rampart Range, Pikes Peak, and the eastern flank of the Wet Mountains.

This model was reviewed by one expert reviewer, who ranked it as good. The reviewer noted that the model appeared to capture the range of the species and identified habitat in the expected topographic position. The reviewer chose "Suitable" for fit, but noted the model was a bit underfit due to presumed coarseness of data.

Boechea crandallii (Crandall's rock-cress), Tier 2

The Colorado distribution of this species is centered in the Gunnison Basin, but occurrences are known from Grand County to northern Hinsdale County, as well as from southern Wyoming. A total of 55 Colorado occurrences were used in modeling. Habitats are generally open, rocky slopes within montane shrubland or forest types, which was reflected in the importance of both geology and biophysical type as model inputs. The species is found in areas where there are an average of 250 frost days per year and seasonal precipitation is generally on the low end, especially in winter. The model threshold was adjusted to the equal training sensitivity and specificity (0.213) to include most documented locations. Modeled suitable habitat was clipped to exclude areas along the southern mountain front, or west of Delta County.

This model was created in Round II of the project and expert review has not yet been solicited.

Calochortus ciscoensis (Cisco sego lily), Tier 2

A regional endemic from western Colorado / eastern Utah, this species is documented in Colorado from only three occurrences in the Grand Valley west of Fruita where Mesa and Garfield counties about the Utah state line. Habitats are in saltbush shrublands on substrates derived from Mancos Shale; distance to these two features were primary contributing factors in the model. The area is dry, averaging less than 3 inches (75 cm) in both spring and summer seasons. March minimum temperatures warm to just under freezing, while extreme summer high temperatures are generally over 105°F (41°C). A handful of extra points were added to the model set to better represent some of the larger polygon occurrence features. A threshold value of 0.5 was used for the probability in the CODEX version of the model. Modeled suitable habitat is constrained to the western end of the Grand Valley.

This model was created in Round II of the project and expert review has not yet been solicited.

Camissonia eastwoodiae (Eastwood evening primrose), Tier 2

This Colorado Plateau endemic has been documented from 11 Colorado locations in Mesa and Delta counties. Occurrences are concentrated on nearly barren Mancos shale salt-shrub habitats in the Grand Valley north and west of Grand Junction, and on lower mesa slopes north of the Gunnison River valley near Hotchkiss. The most important environmental factor contributing to the model was distance to Mancos shale. An average day of last frost in early May and summer precipitation of at least 7 cm in what is regionally a dry season were also important. Extreme minimum winter temperatures are also fairly warm, generally not reaching below -20°F (-29°C). Predicted habitat follows the known distribution around the two separated population centers. A threshold value of 0.5 was used for the probability in the CODEX version of the model and the full extent of the model was retained.

This model was reviewed by one expert reviewer, who ranked it as reasonable with a suitable fit. The reviewer noted the model covers a broader area than known occurrence records, but soils, elevations and overall setting seems correct.

Castilleja puberula (Downy Indian-paintbrush), Tier 2

This Colorado endemic is found in rocky alpine habitats on high peaks of the Continental Divide, with 22 documented occurrences ranging from Larimer to Park County. As could be expected for a species of high elevation cool habitat, extreme maximum summer temperatures were an important factor, rarely exceeding 82°F (28°C). Precipitation amounts in all seasons were also contributing factors, especially for winter and spring; totals across all seasons average nearly 35 inches (90 cm) per year. Aspects tend towards east-facing, and elevations were generally above 10,000 ft (3000 m). Modeled habitat ranges from the vicinity of Hague's Peak (highest point of the Mummy Range) in Rocky Mountain National Park south to around Weston Peak in the Mosquito Range, with smaller areas to the west in the Sawatch Range and is generally concentrated within 25 km of the Continental Divide. A threshold value of 0.5 was used for the probability in the CODEX version of the model and the full extent of the model was retained.

This model was reviewed by two expert reviewers, who believed the model was either reasonable or excellent. One believed the model was underfit, the other suitable. One reviewer questioned the southern extent of the model but the second agreed with this. The second reviewer believed this species is likely more widespread than is currently documented and thought the model showed potentially suitable places which would be good candidates for finding the species.

Cirsium perplexans (Adobe thistle), Tier 2

A Colorado endemic of barren clay "adobe" soils derived from shales of the Mancos or Wasatch formations, this species is documented from 45 locations in the Gunnison and Colorado river drainages of Colorado's western slope. Distances to the two key surface geology types were the primary contributing factors in the model. Days without frost were also a contributing factor; the species appears to tolerate the cooler conditions found in drainages above the Grand Valley, but rarely occurs in more exposed areas of higher elevations to the east. The model threshold was adjusted to 0.25 to better capture documented locations. Predicted suitable habitat extends from western Rio Blanco County southeast to the lower Gunnison Basin and northeastern edge of the Uncompahgre Plateau north of Ridgeway. The full extent of the model was retained.

This model was created in Round II of the project and expert review has not yet been solicited.

Cleome multicaulis (Slender spiderflower), Tier 2

In Colorado, this species is limited to the high intermountain San Luis Valley in Saguache, Rio Grande, Alamosa, Conejos, and Costilla counties, where it occurs in saline or alkaline wetland soils. Fifty-two documented occurrences range from Russell Lakes in the northern valley south to the Rio Grande River valley near the San Luis Hills but are especially frequent in the *sabkha* wetlands south and west of the Great Sand Dunes. The most important environmental factors included roughly equal contributions of soil pH (basic soils preferred) and spring precipitation of at least 5 cm, and, to a lesser extent, distance to palustrine emergent wetland types. Maximum temperatures in summer are slightly higher on the eastern side of the valley where occurrences are most frequent. Modeled predicted habitat follows the known distribution fairly closely but includes quite a bit of additional area in the closed basin wetlands

and greasewood flats west of Saguache Creek. A threshold value of 0.5 was used for the probability in the CODEX version of the model and the full extent of the model was retained.

This model was reviewed by two expert reviewers, who both believed the model was good. One reviewer believed the model was suitable and agreed pH should be an important factor in habitat prediction. A second reviewer the scale was too coarse and should exclude irrigated farmland and urban areas around Alamosa. This model should be considered for eventual revision to exclude unsuitable habitat.

Delphinium alpestre (Colorado larkspur), Tier 2

In Colorado this species of high elevation habitats is documented from 20 occurrences ranging from near the Continental Divide in northwestern Park County south to the Sangre de Cristo Range near the New Mexico border, with a single disjunct historical occurrence from the northern edge of Hinsdale County. Habitats are upper sub-alpine forested types such as spruce-fir and aspen or adjacent alpine rocky areas. At these elevations, average spring minimum temperatures even in May are below freezing, and there can be more than 300 days with frost per year; both factors contributed to the model. Also contributing were surface geology types characteristic of Colorado's central mountain mass, primarily substrates derived from metamorphic, granitic, or tertiary volcanic formations. The model threshold was adjusted to 0.21 to better capture documented locations and clipped to exclude disjunct patches of predicted habitat away from the main range.

This model was created in Round II of the project and expert review has not yet been solicited.

Delphinium robustum (Wahatoya Creek larkspur), Tier 2

This enigmatic species (Sivinski in NMRPTC 1999) is the subject of some taxonomic disagreement, in that some specimens have been annotated to *D. ramosum* while others from the same collection, but at different herbaria, have not. The species is found in north-central New Mexico and south-central Colorado, where it is currently documented from 10 occurrences. Eleven different model runs failed to identify a good habitat-narrowing factor present in available data, indicating that additional research on this species is needed. The selected model primarily used temperature envelope, vegetation type, and aspect to identify a fairly broad extent of likely habitat. Extreme maximum summer temperatures for this prediction are on average are below 90°F (32°C), while extreme minimum winter temperatures average -25°F (-32°C) but may be as low as -33°F (-36°C) following the temperature envelope of Colorado's mid- to upper-elevation regions. Aspects tended to be somewhat west-facing. Vegetation type represented many of the common forest and shrubland types of southern Colorado, including aspen, mixed conifer, spruce-fir, oak-mountain shrub, and sagebrush. A threshold value of 0.5 was used for the probability in the CODEX version of the model, and the model extent was clipped to restrict habitat to central and southwestern Colorado.

This model was created in Round II of the project and expert review has not yet been solicited.

Draba graminea (San Juan whitlow-grass), Tier 2

A Colorado endemic of the San Juan Mountains in southwestern Colorado, this species is documented from 40 occurrences in alpine or upper sub-alpine habitats. Climatic conditions at high elevations were the primary contributing factors to the model. In particular, extreme maximum summer temperatures are generally lower than 80°F (27°C) and most areas experience more than 300 days a year where temperatures dip below freezing, and snow can fall in any month. Due to the southern Colorado

location, these locations also receive monsoon moisture in late summer and early fall that contributes to the model prediction. The model threshold was adjusted to 0.10 to better capture documented locations and clipped to exclude disjunct patches of predicted habitat away from the core range in the San Juans.

This model was created in Round II of the project and expert review has not yet been solicited.

Draba smithii (Smith whitlow-grass), Tier 2

This species is essentially a Colorado endemic, although it may also occur in adjacent New Mexico. The 31 documented Colorado occurrences are clustered in a few widely separated areas, including the San Juan Mountains near and south of Creede, the Sangre de Cristo Range north of Blanca Peak, and the vicinity of Fishers Peak south of Trinidad. Occurrences are generally on talus and scree slopes from upper foothills to lower alpine elevations. Important environmental factors for this species included distance to selected Tertiary volcanic formations in south-central Colorado, terrain roughness index (which, together with slope indicates rugged, steep terrain), and winter (driest season) and summer (wettest season) precipitation. Only modeled habitat in southern Colorado counties was included (areas of Teller and Montrose counties were omitted). Predicted suitable habitat is generally within the three regions described above, with the addition of substantial habitat in the southern Wet Mountains, around and north of Greenhorn Mountain. A threshold value of 0.5 was used for the probability in the CODEX version of the model.

This model had three reviewers, one with limited experience, one with experience in the San Luis Valley and another with experience east of the Continental Divide. Comments centered around the distribution of the model. Two reviewers commented that the distribution was suitable, although one wondered why the Spanish Peaks were not included. The geology of the Spanish Peaks is similar to the Raton Mesa, and therefore another factor must have excluded this habitat. A third reviewer questioned the inclusion of the Wet Mountains. This reviewer had many negative survey results in that mountain range. The Wet Mountains did have patchy areas of similar geology to the Sangre de Cristo Range, but based on negative surveys of the expert reviewer, this model was revised to exclude the Wet Mountains.

Erigeron kachinensis (Kachina daisy), Tier 2

A species of canyon wall “alcove” wetlands that form where water seeps between less permeable geologic layers, this narrow endemic of southwestern Colorado and adjacent Utah is documented in Colorado from just three occurrences in western Montrose and Mesa counties. As might be expected, steep slopes were the most important contributing factor in the model, which was also limited by using distance to very steep slopes as a factor. The only other contributing factor was the number of frost days per year. Although not in the warmest of west slope habitats, these locations generally experience fewer than 160 days with freezing temperatures in a year. A threshold value of 0.5 was used for the probability in the CODEX version of the model. Modeled suitable habitat extent was clipped to the Dolores and San Miguel River drainages.

This model was created in Round II of the project and expert review has not yet been solicited.

Eriogonum brandegeei (Brandeggee wild buckwheat), Tier 1

This species was not originally included in the project. However, recent communication with federal partners regarding important climate factors for this species made it expedient to produce a revised model to replace the original deductive version for CODEX.

The nine documented occurrences of this Colorado endemic are centered around the upper Arkansas River drainage in central Colorado. Occurrences are closely associated with bentonite clay soils derived from steep, eroding outcrops of the Tertiary Dry Union Formation (in Chaffee County) and lower Cretaceous/upper Jurassic sedimentary layers of Dakota, Purgatoire, Morrison, and Rolston Creek Formations in the vicinity of Cañon City (Fremont County). These are generally very sparsely vegetated light-colored soils with an overstory of open pinyon-juniper woodland. Distance to either of the two geologic groups was the highest contributor to the model. The most important climate factor was fall precipitation, which is generally quite low (5-6 cm or about 2 inches) within the range of the species. Areas in Colorado with comparable low fall precipitation include most of the upper Arkansas River drainage, the central San Luis Valley, central South Park in the vicinity of Antero Reservoir, and the Point of Rocks vicinity east of Greeley. The model only included the upper Arkansas River drainage and a small area around the vicinity of Antero Reservoir in South Park. Annual precipitation for the range of the species is not exceptionally low for Colorado, but late growing season climate water deficit appears to be limiting to most other understory species. Soils are somewhat alkaline. Occurrences also tend to be on eastern exposures, and are able to tolerate extreme summer temperatures well over 100°F. The moisture retention capacity of bentonite clay-bearing soils may support the persistence of *Eriogonum brandegeei* in an otherwise challenging habitat. A threshold value of 0.5 was used for the probability in the CODEX version of the model and the full extent of the model was retained.

This model was reviewed by two expert reviewers, both who ranked it as reasonable. The reviewers noted the model excluded a known, well-documented location around Cañon City and did a better job picking up habitat around Salida. Additionally, the reviewers believed the model included too many flat or grassland areas. The species can be found on in flat areas but is more likely on steep eroded habitat. To address the model review, the threshold level for the CODEX model was lowered to a probability of 0.25 to include the known location outside of Cañon City. This added more modeled habitat throughout the range, and the model appears underfit in the Chaffee County. This model should be considered for eventual review, possibly running separate models for the Chaffee County and Fremont County populations and making an ensemble with the two.

Eriogonum clavellatum (Comb Wash buckwheat), Tier 2

A regional endemic of the Four-Corners area, this species is documented from 13 occurrences in Montezuma County in Colorado, none of which have been observed more recently than 2003. The fine textured soils of its mat saltbush shrubland habitats are derived from Mancos Shale. Together with extreme minimum winter temperatures not below -15°F (-26°C), distance to substrates of Mancos Shale accounted for more than 85% of model contribution. Soils have high clay content. Spring is the driest season; occurrences are in an area where precipitation is this season is less than 2.7 inches (6.8 cm). The model threshold was adjusted to 0.48 to better capture documented locations and clipped to restrict predicted habitat to Montezuma County.

This model was created in Round II of the project and expert review has not yet been solicited.

Eriogonum coloradense (Colorado wild buckwheat), Tier 2

A central Colorado endemic or alpine or near-alpine elevations documented from 26 locations in Chaffee, Gunnison, Park, Pitkin, and Saguache counties, this species is closely tied to soil type in some, but apparently not all locations (although it is difficult to be sure since soil data is of variable quality across the species' range). Although not associated with a particular geologic substrate, geology did

contribute to the model, picking out locally mapped areas of a comprehensive selection of formations from ancient to more recent. As expected for a species of high elevations, temperature was most important. Extreme winter low temperatures of -40°F (-40°C) or lower are characteristic of the habitat. Mean April minimum temperatures are below freezing, and snow cover persists longer than at lower elevations. The model threshold was adjusted to 0.20 to better capture documented locations and clipped to exclude predicted habitat too far outside the known range. The model did a poor job of picking up known locations in atypical habitat of Redcloud channery loam, 3 to 30 percent slopes.

This model was reviewed by one expert reviewer who believed the model was reasonable, covering areas near occurrences which they had found. However, the model generally missed the primary soil type where occurrences are in their field office (Redcloud channery loam). The modeling process included inputs of known species locations in this habitat type, but similar habitat was not included in the model despite several test runs. More surveys and data collection in this habitat type could help provide species input data for future model revisions. Additionally, the reviewer noted the model includes unsuitable habitat of riparian areas and developed areas.

Eriogonum pelinophilum (Clay-loving wild buckwheat), Tier 1

This species was not originally included in the project. However, recent communication with federal partners made it expedient to produce a revised model to replace the original deductive version for CODEX.

This Colorado endemic is federally listed as Endangered, with a small range in Delta and Montrose counties. The species has been the object of extensive survey and monitoring efforts and is currently documented from 22 well mapped occurrences on adobe clay soils derived from Mancos Shale west of the Uncompahgre River. As expected, distance to this substrate was the primary contributing factor in the model. These areas are also closely associated with mat saltbush shrublands. Under the constraints of data used for this project, we were not able to further refine the habitat beyond what is already known. The model threshold was adjusted to 0.30 to pick up mapped features, with the eastern edge of the range only picked up at this lower threshold, and habitat outside the known range was excluded.

This model was reviewed informally by members of the *Eriogonum pelinophilum* SSA technical team and critiqued as being too coarse-scale. Using LiDAR and finer scale geology data was considered but rejected as unsuitable for our current methodology. Fine-scale geology was not available across the entire range of the species and relying on this would result in a gap in the model across the species range. To incorporate LiDAR data, all other data, including climate data, would need to be down sampled to match the fine grid of LiDAR data. Down sampling climate data to this degree over-exaggerates the accuracy of this data.

Frasera coloradensis (Colorado green gentian), Tier 2

A Colorado endemic of shale and sandstone breaks in grasslands in extreme southeastern Colorado, this species has been documented across less than 300 acres in 32 occurrences. Documented locations range from small outcrops on plains below the slopes of Black Mesa, along a northeast trending line of shallowly dissected hills following the general direction of Two Butte Creek. This stretch more-or-less outlines the southern limb of surface exposures of Cretaceous age Carlisle shale/Greenhorn Limestone and Graneros shale (Kcg). Close proximity to this group of sedimentary, outcrop-forming formations was the most important environmental variable in the model. Most occurrences were at increasing distance

from shale outcrops of the Niobrara Formation, which is commonly adjacent to the Kcg north of the canyon of the Purgatoire River, with occasional surface presence on the south side. Extreme maximum summer temperatures in this area are somewhat cooler than in the valley of the Arkansas River to the north. Modeled habitat follows the overall range of the species fairly closely. A threshold value of 0.5 was used for the probability in the CODEX version of the model and the full extent of the model was retained.

This model was reviewed by one expert reviewer, who provided comments over email that it appeared to be spot-on.

Herrickia horrida (Canadian River spiny aster), Tier 2

This species is found in extreme south-central Colorado and northern New Mexico, with only 10 element occurrence records in the state. Two Maxent models were reviewed for this species: one with the Raton Formation included and one without. The model with the Raton Formation was chosen to include a larger high probability modeled area. The most important environmental variables for this model were distance to the Raton Formation (81.4% contribution) with northness values near -1 (i.e., south-facing slopes) and summer precipitation combined explaining another 10%. A threshold value of 0.5 was used for the probability in the CODEX version of the model and the full extent of the model in Colorado was retained.

This model had two expert reviewers. One believed it was spot-on. The second commented that areas to the western edge were not likely suitable habitat, although there could be some potential microsites, based on their survey work in nearby areas in New Mexico. There is one known location, documented by an EO, in the area this reviewer suggested should be excluded; therefore, we retained all the modeled area.

Ipomopsis globularis (Globe gilia), Tier 2

This Colorado endemic is restricted to the Mosquito Range in central Colorado except for a disjunct population found on Mt. Elbert in the Sawatch Range in 2015. The species is found on alpine ridges with gravelly, calcareous soils. Two Maxent models were run for this species: one without and one including glacial drift in an attempt to pick-up high-quality occurrences on erosional substrates near high elevation limestone. The model with glacial drift was chosen, but areas around the Collegiate Peaks, which are primarily granite, were excluded. The most important environmental factors explaining the model were distance to units containing Leadville (and Manitou) Limestone, elevation and distance to glacial drift of the Pinedale and Bull Lake age. The probability for the cutoff of the binary model in CODEX was set to 0.177 to include medium tier values of the model, which picked up the Mt. Elbert area. The model was also clipped to exclude the Cottonwood Pass/Collegiate Peaks areas as the geology is significantly different with granite versus limestone.

This model was reviewed by two expert reviewers, both who ranked it as good. Both reviewers believed the model to be at least a bit broad but recognized the ability to create a finer scale model could be limited.

Ipomopsis ramosa (Coral ipomopsis), Tier 1

This is a narrow Colorado endemic, found in two side canyons of the Dolores River Canyon in Montezuma County. Three occurrences are documented on soils derived from the red sandstones, siltstones, and shales of the Permian age Cutler Formation. The important environmental variables

defining this model are distance to the Cutler Formation, northness (prefers south facing slopes), and average minimum May temperature. A threshold value of 0.5 was used for the probability in the CODEX version of the model. The entire extent of the model, which extended approximately 45 miles from EO records, was included in the version for CODEX. We felt this represented an acceptable potential range for this under-surveyed species.

This model was reviewed by two expert reviewers, one who believed the model was un-usable and the other reasonable. Both reviewers were experienced with this species and thought the model covered too broad of a habitat. One reviewer reported negative surveys within 8 miles of the type locale for this species. This reviewer believed a model of the Cutler formation, north facing aspect and elevation between 8,000-9,200 would be more accurate. The second reviewer believes the species may be of a young evolutionary age and has not expanded into all potential habitat, making modeling difficult. This reviewer has surveyed the Hermosa Creek drainage, particularly the large valley west of Purgatory Ski Resort, many times and does not believe the species is there. The reviewer was also skeptical of habitat to the west of the core known area, although they were less familiar with this area. This model should be considered for eventual revision.

Lepidium crenatum (Alkaline pepperwort), Tier 2

In western Colorado this species is largely known from pinyon-juniper woodlands or adjacent vegetation types, and ranges from Moffat County in the north to Montezuma and La Plata counties in the south. It is documented from 27 occurrences in Colorado, many of which are historical and would benefit from better mapping. The top contributing factor in the model was surface geology, which picked up a variety of Jurassic- to Cretaceous-age sandstone and shale bearing formations that are characteristic of Colorado's west slope. Biophysical types were also widespread varieties typical of western Colorado. Extreme winter minimum temperatures are typically not below -22°F (-30°C). The model threshold was adjusted to 0.47 to better capture documented locations and the full extent of the model was retained. This species could benefit from additional survey and study; the known habitat does not well explain its rarity. If additional factors can be identified, a new model should be produced.

This model was created in Round II of the project and expert review has not yet been solicited.

Lepidium huberi (Huber's pepperwort), Tier 1

Little is known about this Tier 1 SWAP PGCN. Its range extends from eastern Utah to western Colorado, and the species is documented from 19 widely scattered occurrences in sagebrush to pinyon-juniper in Rio Blanco, Garfield, and northern Mesa counties. All EO records for this species are historical or extant. The best model included both distance to Green River Formation (widespread in this region) and a categorical surface geology layer. Together these two factors accounted for 85% of the model prediction. The species also appears to prefer areas where extreme minimum winter temperatures do not generally fall below -31°F (-35°C). The classification cutoff used for CODEX was 0.45 to include more modeled habitat associated with known EO records and the full extent of the model was retained.

This model was reviewed by one expert reviewer, who believed the model to be reasonable. The reviewer had made one collection of this rare species. The collection fit well within modeled boundaries and the modeled area looked to be of similar habitat.

Limnorchis zothecina (Alcove bog orchid), Tier 2

This species is found in canyon seep habitats of the Colorado Plateau where it occurs in Utah, Arizona, and Colorado. The Colorado range of this canyon-wall alcove species coincides with that of *Anticlea vaginatus*, although the two species do not necessarily occur together. It is documented from five occurrences in the canyon and side drainages of the Yampa River in Dinosaur National Monument. Annual precipitation in this dry region is less than 12 in. (30 cm). This factor was important in constraining the modeled range but is clearly not a requirement for the species since it occupies small wetlands within the arid environment. Distance to springs was correspondingly important, as the proximity of places where groundwater emerges indicates the likely presence of seep habitat in the vicinity, even if not mapped. Finally, as could be expected, proximity to slopes steeper than 30 degrees was an important contributing factor. The model threshold was adjusted to the equal training sensitivity and specificity (0.362) to include documented locations, and habitat outside of Dinosaur NM was excluded.

This model was created in Round II of the project and expert review has not yet been solicited.

Lomatium concinnum (Colorado desert-parsley), Tier 2

This Colorado endemic species has a range that largely overlaps that of *Eriogonum pelinophilum* in the valley of the Uncompahgre River between Montrose and Delta, but extending further south towards Ridgeway, and reaching east up the Gunnison River drainage to the vicinity of Hotchkiss. It is documented from 38 occurrences, typically in mat saltbush shrubland on soils derived from Mancos Shale. As expected, distance to Mancos Shale was the primary contributing factor in the model. Terrain roughness index also picked up the gentle to moderately sloping habitats of this species. Winter precipitation of at least 2 in. (5 cm) was important as well. The model threshold was adjusted to 0.15 to pick up mapped features, and habitat outside the known range was excluded.

This model was created in Round II of the project and expert review has not yet been solicited.

Lupinus crassus (Payson lupine), Tier 2

This Colorado endemic is documented from 17 occurrences in western Montrose County, where it is associated with sparsely vegetated pinyon-juniper woodland understory. Substrates are alluvium derived from Mancos shale or Chinle formation (upper Triassic mud/silt/sandstone). Primary environmental factors in the model were distance to Quaternary alluvium and eolian deposits, and to a lesser extent, extreme maximum summer temperatures exceeding 102°F (39°C). Predicted habitat follows the known distribution in Paradox Valley and on mesa parklands northeast of the San Miguel River canyon. Additional potential habitat is predicted in the Sinbad Valley at the Mesa/Montrose County line. A threshold value of 0.5 was used for the probability in the CODEX version of the model and the full extent of the model in Colorado was retained.

This model was reviewed by one expert reviewer, who believed the model to be poor. The reviewer believed the model greatly overstated habitat suitability where modeled, and understated extent. This reviewer believes the species is found in or very near ephemeral drainages where either coal bearing, or relatively high clay content soils are present. Fine-scale mapping of these factors (ephemeral drainages and specific geology) are not available over the extent of our modeling area. The reviewer also noted the model missed known locations to the northeast and east of Nucla and believed modeled habitat should have included an expanded range of precipitation and elevation. Unfortunately, data on locations

around Nucla were not available at the time of modeling (Spring 2021). This model should be considered for eventual revision.

Lygodesmia doloresensis (Dolores River skeletonplant), Tier 1

This species is known from extreme eastern Utah and western Mesa County, Colorado, where 13 occurrences are documented. Soils are reddish alluvium or colluvium derived from the Permian age Cutler Formation. Many of the occurrences are along roads, and there appear to be fewer plants with increasing distance from the roadside, which led us to include CNHP's Landscape Disturbance Index as an environmental input. Summer precipitation (at least 5 cm) was the most important contributing factor in the model, followed by distance to surface geology of the Cutler Formation, an average last frost date in late April, and April minimum temperatures generally not below freezing. The model was clipped to exclude a small area of modeled habitat in Montrose County and eastern Garfield County, retaining modeled habitat in Mesa and southwestern Garfield County. A probability of 0.34 was chosen for the cut-off value for the binary version of the model for CODEX to include more modeled habitat associated with known EO records.

This model was reviewed by one expert reviewer, who believed the model to be good. The reviewer noted that the model included known occupied habitat for the species but excluded areas of known habitat in the north desert, east of Highway 139. The model was not able to pick that area up well, despite having those locations included as an input. This model should be considered for eventual revision.

Mentzelia paradoxensis (Paradox stickleaf), Tier 2

This is a Colorado endemic of the salt anticline valleys in western Montrose and San Miguel counties, where it is documented from nine occurrences. Substrates are gypsum clay-dominated soils derived from Triassic to Jurassic sedimentary formations of area. This is a warm and dry area. Important climate factors were extreme minimum winter temperatures generally not falling below -20°F (-28.5°C), and April minimum temperatures near freezing. Summer and fall, while dry, are the highest precipitation seasons, and occurrences are in areas receiving precipitation of at least 2.5 inches (6.5 cm) in summer and 4 inches (10 cm) in fall. A threshold value of 0.5 was used for the probability in the CODEX version of the model. Model extent was clipped to habitat within approximately 35 miles of known occurrences.

This model was created in Round II of the project and expert review has not yet been solicited.

Mentzelia rhizomata (Roan Cliffs blazing star), Tier 2

This Colorado endemic species is known from 33 occurrences on the Roan Plateau in Garfield County. Habitats are steep, shaley slopes formed in the Parachute Creek member of the Green River Formation (common both in the Roan Plateau and at the rim edges of the Piceance Basin to the north). Along with distance to the Green River Formation in general, distance to the Parachute Creek member contributed nearly 85% of information in the model. Moderate soil depth and somewhat alkaline soils were apparently sufficient to confine the modeled habitat to the Roan Plateau and a small area of Battlement Mesa. A few pixels of higher probability modeled habitat in upper Rio Blanco County were omitted from the final model as this was well outside the known range. A threshold value of 0.5 was used for the probability in the CODEX version of the model.

This model was reviewed by one expert reviewer, who believed the model to be reasonable. This reviewer felt they had little experience with this species due to the remoteness of the habitat, but believed the elevation, setting and soils covered by the model appear accurate.

Mertensia humilis (Rocky Mountain bluebells), Tier 2

Colorado occurrences of this species are at the southern end of the core range of the species which is primarily in Wyoming. Nine occurrences are documented in north-central Colorado from Jackson and Larimer counties. This species does not appear to be narrowly restricted to particular habitats; it is found in sagebrush shrublands and open areas in montane forests. Vegetation was not a primary contributing factor in the model, which was largely driven by climate factors. The species occurs in high inter-mountain valleys (North Park, Laramie River Valley), where climatic conditions are cold compared to lower elevations, but also drier than surrounding mountain terrain. Extreme winter minimum temperatures below -37°F (-38°C) can be expected, and spring minimum temperatures do not consistently warm above freezing until June. Occurrences are in areas receiving precipitation amounts of at least 3.5 inches (9 cm) in summer, and 3 inches (7.5 cm) in winter. The model threshold was adjusted to the equal training sensitivity and specificity (0.528) and clipped to exclude habitat outside Jackson or Larimer counties.

This model was created in Round II of the project and expert review has not yet been solicited.

Mimulus gemmiparus (Budding monkey flower), Tier 1

This Colorado endemic is found on sheltered granite rock outcrops associated with seeps from Larimer to Park counties. This species was difficult to model due to lack of detailed environmental layers representing rock outcrops and seeps. A new environmental input layer of rock outcrops was created specifically for this model, with the modeler marking outcrops based on aerial photos. Two models were produced and reviewed, with the second model using a layer of rock outcrops marked from aerial photos ultimately chosen. The overwhelmingly important environmental factor for this model was the presence of rock outcrops, with aspect and climatic variables contributing around 4% of importance. A threshold value of 0.5 was used for the probability in the CODEX version of the model and the full extent of the model was retained.

This model was reviewed by two expert reviewers, both who believed the model was reasonable, but too coarse to accurately capture the habitat of this species, which is restricted to microhabitats. One reviewer suggested including landscape position as an environmental factor (mid-lower slopes with a drainage above), but this environmental input layer was not readily available. The second reviewer noted that as well as being underfit in places due to microhabitat requirements, the model also appeared overfit in places, with Horseshoe Park in Rocky Mountain National Park excluded from modeled habitat. They noted this population is ephemeral, and likely dependent on flood events, suggesting a metapopulation structure for this species.

Nuttallia chrysantha (Golden blazing star), Tier 2

This Colorado endemic is known from 28 occurrences in Fremont and Pueblo Counties. The range includes the vicinity of the Cañon City embayment at the junction of the southern Front Range and the Wet Mountains, and along the Arkansas River as far as Pueblo Reservoir. Habitats are typically moderately steep, barren slopes formed in calcareous substrates of the Smoky Hill member of the Niobrara Formation or other upper Cretaceous geology. Distance to shale barrens formed a substantial

portion of the model. A minimum level of fall precipitation around 5 cm, and gentle to moderate slopes were characteristic. Predicted suitable habitat matched the known distribution fairly closely. A threshold value of 0.5 was used for the probability in the CODEX version of the model and the full extent of the model was retained.

This model was reviewed by one expert reviewer, who believed the model to be good, given the known distribution and habitat affinities of the species.

Nuttallia densa (Arkansas Canyon stickleaf), Tier 2

As indicated by its state common name, this Colorado endemic is largely known from the canyon of the Arkansas River between Salida and Cañon City. Twenty-six occurrences are documented from Fremont and Chaffee counties. Habitats are dry open areas in washes, roadsides, and naturally disturbed sites. Important environmental drivers included fall precipitation, distance to water as a surrogate for proximity to steeper drainage areas (i.e., canyon slopes) and degree of slope. The cut-off probability for the CODEX model was set to the medium probability value of 0.112 to include habitat covering highly ranked, large EOs. Consequently, predicted habitat extended up the Arkansas River drainage as far as Buena Vista, and to side drainages near and below Cañon City. The full extent of the model was retained.

This model was reviewed by two expert reviewers, one who believed the model was reasonable and the other good. Both believed the model covered too great of a spatial extent, noting they had never seen the species north of Salida or east of Cañon City. One reviewer also believed the model did not cover enough habitat in the tributaries on the south side of the Arkansas River Canyon. This reviewer also suggested including geology as an environmental factor, as they have only observed the species on Precambrian rocks. A review of known occurrences shows this species is on older rocks, up to the Paleozoic era, but geology was not included in the modeling process, as mapping was not good enough to pick out a pattern. If more fine-scale geology information becomes available, the model should be considered for revision.

Oenothera acutissima (Narrow-leaf evening primrose), Tier 2

In Colorado, this species is restricted to higher elevations in western Moffat County where 15 occurrences are known in the vicinity of Cold Spring Mountain, Douglas Mountain, and Round Top Mountain (areas that essentially form the extreme eastern end of Utah's Uinta Mountains). The species is reported to be associated with seasonally wet areas in this typically dry landscape. These small habitat patches may be connected with the presence of faults and rock joints where seeps and springs form – a poorly mapped environment. Distance to known springs formed an important part of the model, along with summer precipitation of 5-10 cm. Winter extreme minimum temperatures are generally not lower than -40°F (-40°C). Sparse winter (driest season) precipitation and more abundant fall (wettest season) precipitation were characteristic. Modeled habitat matches the known distribution fairly closely. A threshold value of 0.5 was used for the probability in the CODEX version of the model and the full extent of the model was retained.

This model was reviewed by one expert reviewer, who believed the model to be excellent. The reviewer emphasized the importance of a perennial water source for this species, and distance to springs was an important environmental driver in this model.

Oenothera coloradensis ssp. *coloradensis* (Colorado butterfly plant), Tier 1

This formerly federally listed threatened species is limited in range to southeastern Wyoming, western Nebraska, and northeastern Colorado, where it is documented from 14 occurrences. Habitats are generally sub-irrigated alluvial soils. The range of modeled habitat was truncated to only include areas in Douglas County and north. The western boundary of the range was clipped to a contour at the 6560 ft (2000 m) elevation level, which excluded some higher elevation habitat in the vicinity of Estes Park. Important environmental variables included distance to combined REGAP Western Great Plains floodplain and Basin wide herbaceous riparian ecological systems, distance to wetland polygons attributed to Palustrine Emergent Saturated and Palustrine Scrub-Shrub, and extreme maximum summer temperatures. The medium probability value returned by Maxent was used as the classification cut-off for the CODEX binary model to include more EOs covered by modeled habitat.

This model was reviewed by one expert reviewer, who believed the model to be good, given the known distribution and habitat preferences.

Oenopsis foliosa var. *monocephala* (Rayless goldenweed), Tier 2

This Colorado endemic is found in a restricted range in Las Animas County on semi-arid shortgrass steppe on highly eroded soils. The most important environmental drivers of the model were distance to shale barrens, average percent silt in soil, Colorado National Vegetation Classification type (developed areas excluded) and distance to the Niobrara Formation. This model predicted high probability habitat as far north as Denver, and was truncated to Kiowa, Crowley, Pueblo, Huerfano Counties and areas further south. A threshold value of 0.5 was used for the probability in the CODEX version of the model.

This model was reviewed by two expert reviewers. One reviewer believed the model was reasonable, capturing the habitat, but questioned the extent of this model. There are documented occurrences of this species in northern and western Las Animas County and Otero and Huerfano Counties. The reviewer was un-aware of these documented occurrences. The second reviewer returned comments only over email and noted the modeled habitat overlaps with the range of *Oenopsis puebloensis*.

Oenopsis puebloensis (Pueblo goldenweed), Tier 2

Endemic to a small area north and west of Pueblo, this species is believed to be confined to substrates formed by the Smoky Hill member of the Niobrara Formation. This chalky Cretaceous layer forms rounded hilly outcrops supporting sparse but extensive stands of pinyon-juniper over nearly bare, light-colored soil (shale barrens). A number of calciphilic (chalk-loving) species both rare and more common are found on these substrates in south-eastern Colorado. The 28 documented occurrences range from the grounds of Fort Carson south of Colorado Springs down to the area around Pueblo Reservoir, and back up the Arkansas River drainage to the vicinity of Cañon City. Distance to shale barrens and distance to surface geology of the Niobrara Formation were the primary contributing factors in the model. Areas flooded by Pueblo Reservoir were removed from the modeled habitat. Using a cutoff of 0.42, the predicted habitat fits fairly closely with the known distribution, although the southernmost location is not covered.

This model was reviewed by one expert reviewer, who believed the model to be reasonable, showing known areas of suitable habitat, matching their observations.

Oreocarya osterhoutii (Osterhout cat's-eye), Tier 2

This is a species of the iconic regional Colorado Plateau pinyon-juniper landscape, occurring in canyons and mesas of western Colorado, adjacent Utah, and northern Arizona. There are eight documented occurrences in Mesa County. Percent cover of Colorado Plateau Pinyon-Juniper vegetation type was the factor contributing most to the model; occurrences were generally in areas with moderate cover, indicating more open woodland areas. Distance to steep slopes was also important; mesa rims and steep canyon slopes are typical habitat. Areas with fewer than 170 days of frost per year were also an important factor. The model threshold was adjusted to the equal training sensitivity and specificity (0.11). Modeled suitable habitat was not clipped and shows potential habitat in adjacent counties as well as close to the documented locations.

This model was created in Round II of the project and expert review has not yet been solicited.

Oreocarya revealii (Gypsum Valley cat's-eye), Tier 2

As indicated by its common name, this Colorado endemic species is a specialist of gypsum soils derived from Mancos shale. Populations are concentrated in the salt anticline valleys of Montrose, San Miguel, and Dolores counties in southwestern Colorado. Distance to Mancos shale was the most important environmental factor; other key factors were an average last frost day around May 15th, coldest winter temperatures generally not below -18.4°F (-28°C), and winter through summer precipitation averaging just over 1 inch (2.7 cm) per month. Modeled higher likelihood habitat is more-or-less restricted to the southeastern end of Paradox Valley, middle portion of Dry Creek Basin, Big Gypsum Valley, and most of Disappointment Valley. A threshold value of 0.5 was used for the probability in the CODEX version of the model and the model was clipped to exclude habitat predicted in Delta and Montezuma Counties.

This model was reviewed by one expert reviewer, who believed the model to be reasonable, but overly broad, estimating less than 10% of the model area as being suitable habitat. The reviewer questioned the inclusion of one large, lower precision element occurrence (EO) as a model input. There are 17 documented EOs for this species in the BIOTICS database, and points from the polygons for all of them were included as inputs in the modeling process. Including points from a lower precision polygon can distort model results, but in this instance, there were sufficient points from other polygons to balance the lack of precision. Over-prediction of the habitat is largely due to the comparatively coarse scale of the environmental inputs, and lack of more relevant data.

Oxybaphus rotundifolius (Round-leaf four o'clock), Tier 2

The distribution of this calciphilic Colorado endemic species is similar in the main to that of *Oonopsis puebloensis*, but includes additional areas southwest of Pueblo, as well as two occurrences about 90 km further south at the Pinyon Canyon Maneuver Site in Las Animas County. The 39 documented occurrences of this species are generally confined to the Middle Chalk and Upper Chalky shale of the Smoky Hill member of the Niobrara Formation. Distance to shale barrens was the primary contributing factor in the model; areas with a first frost in fall during the first week of October were also characteristic. Areas flooded by Pueblo Reservoir were removed from the modeled habitat. Using a cutoff of 0.275, the predicted habitat fits fairly closely with the known distribution, although a location on the eastern edge of the range in Pueblo County is not covered. Additional habitat on shale hills north of the Huerfano River, and outside the northwestern bounds of PCMS are also included.

This model was reviewed by one expert reviewer, who responded over email that the model was reasonable.

Oxytropis besseyi var. *obnapiformis* (Bessey locoweed), Tier 2

Occurrences of this species are essentially limited to Moffat County in northwestern Colorado, and adjacent areas of Utah and Wyoming. A disjunct record from the western margin of the Piceance Basin 70 km to the south has not been observed since 1978. The majority of the 22 documented occurrences are concentrated in Browns Park and east on similar substrates toward the Axial Basin east of Maybell. In an attempt to include several occurrences from substrates other than the Browns Park Formation, a categorical geology layer was used in the final model. Surface geology type was the most important environmental factor but predicted habitat still did not include an older occurrence record near the Wyoming border or the Piceance Basin location. This is a dry region, but the species appears to require at least 5 cm of precipitation in summer, also an important factor in the model. Outside the known distribution, a few areas of suitable habitat were predicted for Blue Mountain in Dinosaur NM, and Raven Ridge in Rio Blanco County. A threshold value of 0.5 was used for the probability in the CODEX version of the model and full extent of the model was retained.

This model was reviewed by two expert reviewers, who believed the model to be good. One reviewer thought the model captured documented areas well. The second reviewer believed the model included too broad of an area. This reviewer knew of areas identified in the model where presence/absence surveys had been completed and no occupied habitat had been documented, particularly in the Hwy 318/Peck Mesa area south to the Yampa River. However, there is a historical EO known from the Peck Mesa area.

Packera mancosana (Mancos shale packera), Tier 1

This Colorado endemic species is known from a single occurrence record on the dissected plateau south of Lone Mesa in south-central Dolores County. Plants occur in a handful of scattered stands across approximately two kilometers. Although Mancos shale is characteristic of the location, the full mapped geological unit was too broad as an environmental unit, so discrete soil units supporting stands of the species were used. The presence of soil units from mapped stands was the most important factor; additional important contributing environmental factors were higher clay percent and deeper soil on flatter areas. Modeled habitat is limited to an area of about 5 by 3 km in the vicinity of the occurrence, on the uplands above Plateau Creek. A threshold value of 0.5 was used for the probability in the CODEX version of the model and full extent of the model was retained.

This model was reviewed by one expert reviewer, who believed the model to be reasonable. This reviewer noted that the species is an obligate of a special rare type of Manco Shale which is highly eroded. The model strove to capture this using discrete soil units as inputs rather than the broader geological unit. The reviewer believed the elevational range covered by the model may be too broad both in higher and lower elevation and areas which are not Mancos Shale should be excluded from the model.

Pediocactus knowltonii (Knowlton cactus), Tier 1

This extremely rare and Federally Listed Endangered cactus is known from only a single native population in pinyon-juniper/sagebrush vegetation in northern New Mexico, just south of the Colorado border. Maxent models using a handful of points placed near the known location were unsatisfactory, so

a deductive model was constructed using soil type polygons in and immediately adjacent to the occurrence. Corresponding soil units in Colorado were also selected. These were intersected with environmental factor layers to select areas where vegetation, growing season length, and annual precipitation were similar to the known location.

Reviewers were not satisfied with this model. Of two reviewers one ranked the model good and the other poor, but both noted the model was underfit and did not capture refined environmental inputs. So, the deductive model was reconstructed using environmental inputs specified in (Handwerk et al. 2017) but extending the model to some areas outside Southern Ute tribal lands. Inputs were vegetation type of Pinyon-Juniper Woodland or PJ-Sagebrush mix, elevation range of 1865-2057m (6100-6750 ft), surface geology of the San Jose Formation, and a variety of soil types (see model metadata for details).

Penstemon acaulis var. *yampaensis* (Yampa beardtongue), Tier 2

With a distribution adjacent to that of the Plateau penstemon in western Moffat County, Colorado and Daggett County, Utah, this species is documented from 31 locations in Colorado. About a third of these occurrences have not been observed within the past 30 years. The Colorado distribution ranges from north of Cold Spring Mountain southeast to the vicinity of Cross Mountain southwest of Maybell. Occurrences are typically on shaley, sandy, limestone soils derived from Browns Park Formation or the Tipton Tongue (including Wilkins Peak member) of the Green River Formation. Distance to one or both of these two substrates accounted for about 90% of the model predictive ability, and the model including both types was better constrained than models with a single type. Minor contributing factors included spring precipitation generally over 7.5 cm and extreme winter minimum temperatures not lower than -40°F (-40°C). Predicted habitat for this species occupies areas of slightly dryer, lower elevations and younger geologic substrates adjacent to that of *Penstemon scariosus* var. *cyanomontanus* (see below) while overlapping very little with that related species. A threshold value of 0.5 was used for the probability in the CODEX version of the model and full extent of the model was retained.

This model was reviewed by two expert reviewers, who believed the model to be either reasonable or good. The first reviewer thought the general area it encompassed was correct but knew of one location in Irish Canyon not captured by the model. The second reviewer compared recently identified occupied habitat (not yet in the BIOTICS database) to the model and did not identify large data gaps but did question the inclusion of the Vermillion Bluffs.

Penstemon degeneri (Degener beardtongue), Tier 2

This Colorado endemic is documented from 21 occurrences on rocky areas in the vicinity of the Cañon City embayment at the junction of the southern Front Range and the Wet Mountains. Substrates are derived from Precambrian age metamorphic and igneous outcrops. The model incorporating surface geology was better differentiated; distance to the aforementioned types was a primary contributing factor. Dry winters (generally less than 10 cm of precipitation) and comparatively wet summer months (16 cm or more) were also important, as was an average last frost date around the end of May. Slopes were moderate to steep. Modeled habitat extends around most of the slopes of Pikes Peak at elevations up to 8000-8500 feet (2440-2590 m) depending on aspect, extending north to the southern end of the Tarryall Mountains in Park County. Similar elevations in the Wet Mountains of central Fremont, northern Custer, and western Pueblo counties are also included. Disjunct areas modeled in central Jefferson and Park Counties were clipped out. A threshold value of 0.5 was used for the probability in the CODEX version of the model.

This model was reviewed by two expert reviewers, who both believed the model was reasonable. One reviewer thought the model did not perform as well for higher elevations and more western locations. They knew of at least one location which was not captured by the model.

Penstemon fremontii var. *glabrescens* (Fremont's beardtongue), Tier 2

A Colorado endemic documented from 18 locations in the Piceance Basin of Rio Blanco County, this species occurs on sparsely vegetated slopes of soils derived from Green River shale. As expected, distance to Green River surface geology was the primary contributing factor in the model (52.7%). Because this unit is fairly coarsely mapped in the available data, predicted habitat is not highly constrained; the model could be considered under fit. Shallow to moderate depth soils also played a fairly large part, contributing 26.7% to the prediction. Additional important factors were aspect (a tendency to favor more south-facing slopes) and a last frost date around the end of May. Modeled habitat was clipped to remove areas south of the boundary between Rio Blanco and Mesa counties. Remaining higher probability habitat includes scattered areas of central Moffat County, substantial area in the Piceance Basin, extending south to Garfield County with a few drainages in the Roan Plateau, and additional areas in the vicinity of the Grand Hogback to the east, and extending up nearly to the vicinity of Gypsum in the Colorado River Valley. A threshold value of 0.5 was used for the probability in the CODEX version of the model.

This model was reviewed by one expert reviewer, who believed the model to be reasonable. The reviewer noted the model did a good job of capturing known occurrences but was skeptical of modeled habitat around 1-70. The reviewer also noted irrigated fields and other disturbed areas should be excluded.

Penstemon gibbensii (Gibben's beardtongue), Tier 1

This species is documented from three locations in northwestern Moffat County, and also occurs in adjacent Wyoming and Utah counties. Originally reported as occurring on soils derived from the Tertiary age Browns Park formation, it was more recently also found on the substrates of the widespread Wasatch formation. Consequently, although distance to Browns Park formation surface geology was a contributing factor in the model, the most important contribution was a general lack of summer precipitation (<6 cm). Dry winters and winter extreme low temperatures warmer than -40°F (-40°C) were also contributors. Most, but not all, habitat tends to be on south-facing exposures. Despite the lack of key substrate information, modeled habitat was fairly tightly constrained to areas near the known locations, i.e., the floor of Browns Park and the vicinity of the junction of the Little Snake River with Powder Wash. A threshold value of 0.5 was used for the probability in the CODEX version of the model and the full extent of modeled habitat was retained.

This model was reviewed by two expert reviewers. One wrote in an email that the model looked good. The other commented that the model may include too much habitat in the western portion of Powder Wash. Marginal suitable habitat had been identified in this area, but no occupied habitat and appropriate substrate may be restricted to sandy bluffs above the Little Snake River.

Penstemon mensarum (Grand Mesa penstemon), Tier 2

A Colorado endemic of the central west-slope higher elevations, this species is documented from 48 occurrences ranging from the southern part of the White River Plateau in Garfield County southwest to the Uncompahgre Plateau in Montrose County, but with its core distribution on and above the Grand

Mesa. Habitats are open meadow areas in typical montane shrubland or woodland types of the west slope. Winter precipitation was the most important contributing factor in the model; occurrences are in areas receiving 6-14 inches (15-35 cm) of precipitation in this season. Aspen or aspen-mixed conifer forest, montane sagebrush, and oak-mixed mountain shrubland accounted for most of the biophysical setting model contribution. A variety of regional geologic substrates were picked up by the model, but the species is apparently not tightly constrained by surface geology. A threshold value of 0.5 was used for the probability in the CODEX version of the model. The model was clipped to include only predicted habitat within approximately 30 miles of known occurrences.

This model was created in Round II of the project and expert review has not yet been solicited.

Penstemon scariosus var. *albifluvis* (White River penstemon), Tier 1

Known from five occurrences in extreme western Rio Blanco County, from Raven Ridge west of Rangely south to the vicinity of Rabbit Mountain, the species is also found in adjacent Uintah County, Utah. Substrates are derived from the Parachute Creek member of the Green River shale, and distance to this surface geology type provided nearly 90% of the model information. Other important factors were soil depth and an average last frost date around the third week of May. Modeled higher probability habitat follows the documented distribution fairly closely, extending somewhat further north along Raven Ridge to the Utah border, and including an additional area south of Park Canyon at the southern end. A threshold value of 0.5 was used for the probability in the CODEX version of the model and the full extent of modeled habitat was retained.

This model was reviewed by two expert reviewers, both who believed the model was excellent. Both thought the model captured all the appropriate habitat with correct geologic substrate and slope.

Penstemon scariosus var. *cyanomontanus* (Plateau penstemon), Tier 2

This species is documented from seven occurrences in western Moffat County, primarily on slopes of Blue Mountain to the south and Douglas Mountain to the north of the canyon of the Yampa River in Dinosaur National Monument. An additional location is known from Diamond Peak some 32 km to the north, and the range extends into adjacent Uintah County, Utah. Substrates are generally sandy, slickrock crevices, or gravel, derived from older rocks of the Uinta Mountain Group (middle Proterozoic) and adjacent Pennsylvanian age sandstone formations, but are not closely tied to a particular geologic formation. The categorical surface geology layer contributed over 50% of the information in the model; five types were important, and three additional types also supported occurrences. Other important factors included summer precipitation of at least 12.5 cm, and extreme minimum winter temperatures generally above -40°F (-40°C). Vegetation type (as biophysical setting) of pinyon-juniper or sagebrush shrubland was also a contributing factor. Modeled suitable habitat includes extensive middle elevation areas in western Moffat County, extending from Middle Mountain in the north to the slopes below Skull Creek Rim in the south. A threshold value of 0.5 was used for the probability in the CODEX version of the model and the full extent of modeled habitat was retained.

We were unable to identify an expert reviewer for this species.

Phacelia gina-glenneae (Troublesome phacelia), Tier 1

This Colorado endemic is known from a single large occurrence in Middle Park near Kremmling, where it is restricted to weathered volcanic ash substrates of the Troublesome Formation. Naturally, distance to this surface geology was the primary contributing factor in the model. Known stands are generally on

western-facing slopes, and where summer precipitation is at least 8 cm. Similar conditions were predicted for hillsides north and east of Kremmling, additional areas extending north and south of the known location in the Troublesome Creek drainage, as well as the valley of the Colorado River, and narrow hillside areas near the junction of the Colorado and Fraser rivers at Granby (just south of the Troublesome Creek burn of 2020). A threshold value of 0.5 was used for the probability in the CODEX version of the model and the full extent of modeled habitat was retained.

This model was reviewed by one expert reviewer, who believed the model to be poor. They noted the model was overly broad, capturing riparian areas and areas around Granby should not be included. This model should be considered for eventual revision.

Physaria alpina (Avery Peak twinpod), Tier 2

This Colorado endemic is documented from 16 occurrences in alpine turf and fellfield habitats in Park, Lake, Pitkin and Gunnison counties. Due to its alpine nature, elevation was the most important contributing factor in the model. Extreme minimum winter temperatures reaching nearly -55°F (-48.5°C) are possible. In these alpine habitats frost can occur any time of year, precipitation begins to fall as snow during fall months, and snow is likely to remain on the ground for more than half the year. Modeling indicated a tendency for occurrences to be on south-east facing slopes, which could mitigate harsh conditions to some degree. A threshold value of 0.5 was used for the probability in the CODEX version of the model. The model was clipped to include only predicted habitat within the range of known occurrences.

This model was created in Round II of the project and expert review has not yet been solicited.

Physaria bellii (Bell's twinpod), Tier 2

A Colorado endemic, this species is known from 28 occurrences on hogbacks at the mountain front in Boulder and Larimer counties. These are areas where during the Laramide Orogeny the rising mountain terrain faulted and tilted overlying sedimentary layers of generally lower Cretaceous or older origin, forming the Front Range (Dakota) hogback. Occurrences range from the northern edge of Boulder city limits to the vicinity of Livermore in northern Larimer County. The species tolerates disturbance to such an extent that it is found on mine spoil piles and road cuts, as long as the substrate is derived from the appropriate rock type. Distance to shale and sandstone units forming the Front Range hogback north of Colorado Springs was the primary contributing factor in the model. The 0.5 and above model did not capture the northern extent of the range well, so the cutoff value was adjusted to 0.35. Potential habitat was truncated at the Boulder/Jefferson County border.

This model was reviewed by one expert reviewer, who believed the model to be reasonable. The reviewer was familiar with this species on shale soils and outcrops associated with the Niobrara, Fort Hayes, and Pierre members/formations. The reviewer thought the depicted habitat area looked overly broad and should not include habitat south of Boulder Canyon.

Physaria parviflora (Piceance bladderpod), Tier 2

In common with other Piceance Basin endemics, this species is closely associated with shaley soils derived from units of the Green River formation, including the Parachute Creek member surrounding the well-known Mahogany ledge oil shale zone. There are 37 locations documented in Rio Blanco, Garfield, and Mesa counties. Together, distance to Parachute Creek member (83%) and distance to Green River formation (5.3%) were the major contributing factors in the model. This species appears to

have a slightly broader environmental niche than *Thalictrum heliophilum*, which has a nearly identical range. Additional model factors indicate that this species prefers the higher, cooler margins of the basin, where last frost average is in first week of June, and extreme maximum summer temperatures generally below 95°F (35°C). Modeled habitat closely tracks the presence of Parachute Creek substrates on the rim of the Piceance Basin with scattered patches on the western end of Battlement Mesa to the south across the valley of the Colorado River. A threshold value of 0.5 was used for the probability in the CODEX version of the model and the full extent of the model was retained.

This model was reviewed by one expert reviewer, who believed the model to be good, tracking well with known occurrences but including too much habitat off exposed shale barrens.

Physaria pruinoso (Pagosa bladderpod), Tier 2

A narrow endemic known from northern New Mexico and southern Colorado, this species is documented from 23 occurrences in La Plata, Archuleta, and southern Hinsdale counties at lower montane elevations on the southern flank of the San Juan Mountains in Colorado. Distance to substrates derived from Mancos Shale, and high-clay soils were important contributing factors in the model. Occurrences are in areas where winter precipitation at least 5.5 inches (14 cm) but not much higher, and last frost in spring is generally during the first week of June. The model threshold was adjusted to the equal training sensitivity and specificity (0.528) and clipped to exclude predicted habitat in San Miguel County.

This model was created in Round II of the project and expert review has not yet been solicited.

Physaria rollinsii (Rollins twinpod), Tier 1

This Colorado endemic is known from the Gunnison Basin, with 18 documented occurrences ranging from the vicinity of Sargents at the east end, west to the upper end of the Black Canyon of the Gunnison River on dry sagebrush-dominated shrublands. This higher elevation basin is slow to warm in spring with average date of last frost around mid-June. Spring precipitation was the most important factor; winter precipitation was also a primary contributor, probably indicating a minimum tolerable winter/spring total precipitation amount for the species. Average last frost was the most important temperature factor, but other spring minimum temperature factors also contributed to the model. A tendency to occur on more south-facing aspects agrees with the idea that the species favors local conditions that may warm slightly earlier at the beginning of the growing season. Modeled habitat extends up many side drainages and ridges both north and south of the Gunnison River valley, following the distribution of sagebrush shrubland. A threshold value of 0.5 was used for the probability in the CODEX version of the model and was clipped to exclude modeled habitat in Grand County.

This model was reviewed by three expert reviewers, two who believed the model was good and one who believed it was un-usable. The first reviewer noted that the model included the known location where they had observed the species. The second reviewer noted that many of the areas where they had documented the species were captured in this model, but it was not an exact match. One area east of Gunnison where the reviewer had an observation record was not included and areas of un-suitable habitat, including riparian areas of Ohio Creek and Tomichi Creek and Gambel oak woodlands (especially north of Blue Mesa Reservoir) were included in the model. The third reviewer echoed these comments but knew of locations of the species on Sapinero Mesa not included in the model and noted areas of

Douglas fir, riparian areas and irrigated meadows should not be included in the model. This model should be considered for eventual revision.

Physaria scrotiformis (West Silver bladderpod), Tier 1

Documented from four high-elevation locations near the continental divide in San Juan and La Plata counties, this Colorado endemic is a specialist of shallow alpine substrates. Fall precipitation averaging about 29 cm was the most important factor, along with maximum summer temperatures (typically cool), and shallow, alkaline soils, of moderately rough terrain. Although the original occurrence is reported from the lower Mississippian age Leadville limestone, subsequent stands have been documented from younger substrates of lower Permian (Cutler Fm) or Tertiary volcanic origin that are fairly common in the San Juan Mountains. Predicted habitat is concentrated in high elevation areas of the Weminuche Wilderness Area between the Las Animas River and Vallecito Creek. Modeled habitat was truncated to include areas from southern Ouray County to northern La Plata County and adjacent western Hinsdale County, omitting areas further to the east. A threshold value of 0.5 was used for the probability in the CODEX version of the model.

We were unable to identify an expert reviewer for this species.

Physaria vicina (Good-neighbor bladderpod), Tier 2

This Colorado endemic species is found primarily on soils derived from Mancos shale or adjacent sedimentary formations in southwestern Delta County, eastern Montrose County, and northern Ouray County. Two disjunct occurrences are known from the southeastern corner of Garfield County, more than 100 km (62 miles) distant from the main distribution. Distance to Mancos shale was the most important contributing factor in the model, followed by last and first frost dates (a growing season roughly between third week of May and third week of September). Winter precipitation (the driest month) was also an important factor. Modeled habitat is concentrated on rising ground above the Uncompahgre and Gunnison rivers in the vicinity of Montrose. Similar habitat ranges southeast from western Garfield County along the Grand Valley, and the opposite (southern) side of the Uncompahgre Plateau. Disjunct predicted habitat is found near the junction of the Crystal and Roaring Fork rivers on slopes above Carbondale. A threshold value of 0.25 was used for the probability in the CODEX version of the model to include more habitat to cover known locations and modeled habitat in Dolores County was clipped out.

This model was reviewed by two expert reviewers, both who believed the model was poor. The first reviewer believed most observations to be on Mancos shale and Dakota sandstone, in both pinon-juniper woodlands and sagebrush steppe plant communities in shallow to moderately deep loams. This reviewer questioned the findings near Glenwood Springs, which influenced the extent of the model, and believed many of the salt desert shrub ecological sites are not suitable habitat. The second reviewer believed the species could be associated with pinon-juniper, sagebrush, and saltbushes, but that the model was too broad, and the presence of Mancos shale was not very helpful for narrowing down the search areas. This reviewer believed the species is not limited by lack of suitable habitat but is under-documented as it blooms very early in the spring and once the flowers are gone, it is indistinguishable from *Physaria acutifolia*, making it easy to overlook.

Potentilla rupincola (Rocky Mountain cinquefoil), Tier 2

This species had been previously modeled for another CNHP project. However, newly documented locations for this species made it expedient to produce a revised model to replace the original version for CODEX.

A Colorado endemic restricted to north-central Colorado, this species is documented from 34 occurrences in Larimer, Boulder, Clear Creek, and Park counties, where it is found on or near granitic outcrops and crevices. A dataset representing distance to rock outcrops in the Front Range was the primary contributing factor in the model. Most occurrences are in areas receiving 4-6 inches (10-15 cm) of summer precipitation, southern locations receive more. The montane to sub-alpine elevations where this species is found can experience extreme winter minimum temperatures as low as -39°F (-39.5°C). Modeled potential suitable habitat is most common in Larimer and Boulder counties but extends as far south as eastern Park County west of Pikes Peak. A threshold value of 0.5 was used for the probability in the CODEX version of the model and the full extent of the model was retained.

This model was created in Round II of the project and expert review has not yet been solicited.

Ptilagrostis porteri (Porter feathergrass), Tier 2

A Colorado endemic closely linked to fen environments, this species is known from 31 occurrence records, ranging from south-central Lake County east to the vicinity of Woodland Park near the Teller/El Paso County border. Most occurrences are in northern Park County and adjacent Summit County, where rich fens are concentrated in drainages fed by streams originating in calcareous substrates. As expected for a fen indicator species, distance to saturated wetlands and distance to water were the most important environmental factors in the model, followed closely by April minimum temperatures well below freezing. In general, these high-elevation occurrences are cool and moist, in areas well able to support saturated soils. The modeled range was truncated to include only eastern portions of Gunnison, Pitkin, and Eagle counties, Summit, Clear Creek, Lake, Chaffee, Park and Teller counties, and small parts of Gilpin, Jefferson, Douglas, El Paso, and Fremont counties, all within 50 miles of known EOs. A threshold value of 0.5 was used for the probability in the CODEX version of the model.

This model was reviewed by three expert reviewers, who believed the model was either reasonable, good, or poor. One reviewer believed the coverage seemed appropriate, although much of the potential habitat appears degraded and unlikely to support populations. Another reviewer believed the modeled habitat was too broad and more refined environmental inputs, which were not available, were needed to narrow predicted habitat. This reviewer expressed concern that this model could give the impression that the species is much more abundant than it actually is and that the model should contain a caveat that it is a broad, inclusive model of locations where the species may exist and should not be taken as a species distribution map. This is essentially the intent of this model, which is to be used for environmental review, and the purpose of this model is described in this report.

Puccinellia parishii (Parish's alkali grass), Tier 2

Colorado has two documented locations of this rare grass of the southwestern US that lie about 25 km (15 miles) apart in central San Miguel and Dolores counties. Its scattered distribution is connected to its occurrence in moist, seasonally wet habitats within the surrounding arid lands. Colorado occurrences are associated with soils derived from Mancos shale or adjacent formations, and distance to Mancos shale was the most important factor in the model. Distance to palustrine emergent wetland types was

also important. Sufficient winter precipitation, and a last frost date around the first of June were additional contributing conditions. Modeled suitable habitat is concentrated around the two known locations, but small scattered patches occur from southern Montrose County south to east-central Montezuma and west-central La Plata Counties. A threshold value of 0.5 was used for the probability in the CODEX version of the model and the full extent of the model was retained.

This model was reviewed by two expert reviewers, who both believed the model was poor. One reviewer was unsure on the distribution of the species in Colorado. A second reviewer thought the model was too broad and contained unsuitable habitat of areas which are not seasonally saturated. This model should be considered for eventual revision.

Salix arizonica (Arizona willow), Tier 2

This rare willow is a subalpine species of wet meadows, streamsides, and cienegas in Utah, Arizona, New Mexico, and Colorado. The three known Colorado occurrences are in southwestern Conejos County. Winter precipitation of at least 10 inches (26 cm) was the most important contributing factor in the model. Spring precipitation was also important; occurrences are in areas receiving at least 7.5 inches (19 cm) during that season. As might be expected with these precipitation patterns, snow could remain present on the ground for more than half the year. Distance to palustrine emergent wetland types, which represent the habitat in which *Salix arizonica* is found, were also an important contributing factor. Modeled suitable habitat was restricted to Conejos and Archuleta counties and a threshold value of 0.5 was used for the probability in the CODEX version of the model.

This model was created in Round II of the project and expert review has not yet been solicited.

Telesonix jamesii (James telesonix), Tier 2

This species had been previously modeled for another CNHP project. However, newly documented locations for this species made it expedient to produce a revised model to replace the original version for CODEX.

This species of rocky areas is largely endemic to Colorado, although it may extend into northern New Mexico. It is documented from 35 occurrences in the Front Range of Colorado. A dataset representing distance to rock outcrops in the Front Range was the primary contributing factor in the model, as was distance to surface geology of igneous and metamorphic rocks of Precambrian age. Most locations for this species receive more than 20 inches (50 cm) of annual precipitation. Reflecting the mountainous distribution of occurrences, extreme winter low temperatures can be severe, while summer extreme maximum temperatures are generally not above 90°F (32°C). The model threshold was adjusted to 0.45 to better cover documented locations and the full extent of the model was retained.

This model was created in Round II of the project and expert review has not yet been solicited.

Thalictrum heliophilum (Sun-loving meadow rue), Tier 2

This Colorado endemic is known from 33 locations in Rio Blanco, Garfield, and Mesa counties. Occurrences are generally found on moderately steep slopes and are closely tied to shaley soils derived from the Green River formation, especially the Parachute Creek member. As expected, distance to Parachute Creek member was by far the most important factor in the model, contributing over 90% of the information; slope was the next greatest contributor to the model. A slight tendency to occur on south to west facing slopes was also seen, but other environmental factors were not major contributors.

Modeled habitat closely tracks the presence of Parachute Creek substrates on the southern and western portions of the Piceance Basin and the western ends of Battlement Mesa and Grand Mesa to the south across the valley of the Colorado River. The model threshold was adjusted to 0.35 to include more of a documented occurrence on Battlement Mesa. The full extent of the model was retained.

This model was reviewed by one expert reviewer, who had little experience with this species. The reviewer believed the model was reasonable based on elevation and soils.

Thelypodopsis juniperorum (Juniper tumble mustard), Tier 2

A west-slope Colorado endemic, this species is documented from 19 occurrences in Mesa, Delta, Montrose, and Gunnison counties where it is found in typical habitats of the northern Colorado Plateau including pinyon-juniper, sagebrush, and oak-mixed mountain shrublands. Biophysical settings selected in the model were dominated by pinyon-juniper woodland and a variety of shrubland types such as oak-montane shrub, mixed salt-desert scrub, lower montane foothill shrubland, and sagebrush. Surface geology was variable, but largely formed in sedimentary units of Jurassic to Cretaceous age or overlying Quaternary substrates. Soils generally have high clay content. Slopes are generally moderate but not flat, and extreme minimum winter temperatures generally warmer than -26°F (-32°C). A threshold value of 0.5 was used for the probability in the CODEX version of the model and modeled suitable habitat was clipped to include only west-slope areas.

This model was created in Round II of the project and expert review has not yet been solicited.

Thelypodium paniculatum (Northwestern thelypod), Tier 2

There are two poorly documented occurrences of this species from the first half of the 20th century in Colorado. Colorado occurrences would be peripheral to the central distribution of the species in Wyoming, where it is reported as a species of mesic or wet meadows and riparian areas. A deductive model was constructed, based on vegetation type and elevation. Modeled habitat included areas with elevations of 7,500-9,500 ft (2286-2895.6 m) and existing vegetation type (LANDFIRE Remap 2016) Rocky Mountain Alpine-Montane Wet Meadow, Rocky Mountain Lower Montane-Foothill Riparian Shrubland, Rocky Mountain Lower Montane-Foothill Riparian Woodland, Rocky Mountain Subalpine-Montane Mesic Meadow, Rocky Mountain Subalpine-Montane Riparian Shrubland, or Rocky Mountain Subalpine-Montane Riparian Woodland. The intersection of these two datasets was clipped to include only Moffat, northern Routt, and Jackson counties, and the lower Laramie River valley of Larimer County.

This model was created in Round II of the project and expert review has not yet been solicited.

Townsendia fendleri (Fendler's townsend-daisy), Tier 2

In Colorado, this species is documented from 26 locations in the south-central part of the state. Occurrences in the upper Arkansas River valley are typically on eroded badland outcrops in pinyon-juniper woodlands derived from Tertiary or Cretaceous sedimentary formations, but those in more southern stands do not share this substrate affinity. Distance to Southern Rocky Mountain Pinyon-Juniper Woodland vegetation type was the primary contributing factor in the model. The model also picked up foothill and montane grassland biophysical settings as contributing to the habitat. Fall precipitation is generally low in the range of this species, ranging from 2-3 inches (5-8 cm). A subset of occurrences in the Upper Arkansas Valley were used for modeling to decrease the influence of this more densely populated region in model results. The model threshold was adjusted to 0.25 to better cover

documented locations. Modeled suitable habitat was clipped to exclude areas west or south of Poncha Pass at the north end of the San Luis Valley.

This model was created in Round II of the project and expert review has not yet been solicited.

Townsendia glabella (Gray's Townsend-daisy), Tier 2

This endemic of southwestern Colorado is documented from 22 locations, nine of which are considered historic (not observed during the past 30 years). Although reported as occurring on “the Smokey Hill member of Mancos shale”, geologic sources warn that the difficulty of mapping units corresponding to Niobrara formation members in the Mancos shale between the Dakota Sandstone/Burro formation and the Mesa Verde group in this part of Colorado is extreme. Consequently, the species was modeled using distance to units of the Mancos shale as this substrate is currently mapped on 1x2 degree maps for the area. In addition to distance to Mancos shale surface geology, moderately deep soils with comparatively high clay content at elevations generally below 7550 ft (2300 m) were characteristic. May minimum temperatures above freezing may also contribute an important isoline in the distribution. Modeled higher probability suitable habitat omits some historical occurrence records and was clipped to limit the final extent to Montezuma, La Plata and Archuleta counties. A threshold value of 0.5 was used for the probability in the CODEX version of the model.

This model was reviewed by two expert reviewers, one who believed the model was reasonable and the other poor. The first reviewer believed the model was coarse and should exclude wetlands and the town. This reviewer would like to see the model be more closely aligned with platy shale barrens. A second reviewer agreed that this model was too broad, picking up area in the forest with shale geology but not shale soils. This model should be considered for eventual revision.

Trifolium dasyphyllum ssp. *anemophilum* (Whip-root clover), Tier 2

This species is currently only documented in Colorado from a single occurrence in Weld County; it is primarily known from south central Wyoming and the Laramie foothills. A deductive model was constructed from soil types only (NRCS 2012) and limited to Weld and Logan counties. Soil map units selected to represent the habitat were Ustic Torriorthents-Rock outcrop complex, 9 to 40 percent slopes (MU Key 95166) and Badland (MU Key 95101).

This model was created in Round II of the project and expert review has not yet been solicited.

Discussion

Coverage and use of available species models

The Colorado Natural Heritage Program tracks roughly 540 plant species, with 117 of these, the Plants of Greatest Conservation Need, ranked globally critically imperiled (G1) or imperiled (G2). These species are at risk throughout their range and under threat of extinction. Pressures on these species include oil and gas development, recreation, and suburban or exurban development. Many of these species are under-surveyed and little is known about their life history and environmental needs. This project defines both mapped locations of potentially suitable habitat and identifies environmental drivers to give a better understanding of species most important needs.

Models produced during this project are suitable for use in identifying field survey target areas, and for landscape scale spatial analysis or to aid in management of and avoidance of impacts to the species.

Because the primary use of these models in CODEX is to assist landowners and managers in identifying which species of concern are most likely to occur in an area of interest, we were not concerned that models would be overly constrained by using known typical substrates as primary input, as long as predicted habitat did not exactly outline individual occurrences. Binary versions are easily exported to kml/kmz format for use in Google Maps and Google Earth and are smaller files for use on other GIS devices.

For some species, a few element occurrence (EO) records were excluded from use in habitat modeling. Excluded records were typically very old historical or extant EOs, and those with low spatial precision (mapped as covering very large spatial areas). This exclusion can result in such locations falling outside high probability modeled habitat. Our modeling process aims to define areas of most likely habitat for the species, not simply to buffer all known EO locations. Higher probability modeled areas include habitats with environmental conditions most similar to the greatest number of known occurrences and may exclude EOs which do not meet these criteria. These EOs, therefore, will fall within lower probability areas of the model. We recognize that the binary model in CODEX may exclude some documented occurrences of the species; however, our intent is to delineate areas most likely to harbor the species, striking a balance of including the most similar areas near documented locations while not excluding additional reasonable habitat.

With the addition of the 80 species modeled in this project, modeled distribution for a total of 107 Plants of Greatest Conservation Need will be represented in the CODEX. The spatial display of the modeled distribution will be visible to the user in CODEX beginning in April 2022 and the modeled distribution will be used in the analysis for environmental review. Results will be returned in tabular form. The models used in CODEX will be a binary version (yes/no) of the full probability model delivered to CNAP which includes a likelihood from 0-1 over the modeled area. Full spatial models have been delivered to CNAP for use in species surveys or other conservation work. Portions of these full spatial models could be shared with agency partners with a signed data-sharing agreement and with CNAP permission as needed.

Additional modeling needs

Although nearly all SWAP Tier 1 and 2 species have been modeled, there remain two SWAP Tier 2 Plants of Greatest Conservation Need lacking a species distribution model (Table 2). We chose not to model these species at this time due to unresolved taxonomic issues. There are also 27 of the older Tier 1 models produced as rapid assessment deductive models produced as binary surfaces; most would be improved by remodeling using better techniques. CNHP continuously re-evaluates species rankings as new species are described, known species revised, or additional occurrences found, so that these totals are likely to change over time. New occurrence records are regularly submitted to or requested by CNHP. This data can be checked against existing models and used to determine when a new model is needed. In addition, a number of the models produced during this project could benefit from additional work to identify more useful model inputs that might improve the predictive ability of the model. Many of these were identified in the expert review process, which will be completed in Fall 2022, and used to prioritize model revision. Three species distribution models were revised in winter 2022 based on model reviewer feedback. For a discussion of model review results and potential processes to improve models, see Future data development section below and Appendix C.

There are an additional 28 BLM sensitive species in Colorado not included in SWAP that lack models, and 23 USFS Region 2 sensitive species occurring in Colorado not included on either the SWAP or BLM list that have not been modeled (Table 2). There are also perhaps 110 or so fully tracked G3 (rounded rank) species without special status which could be modeled if occurrence data is available.

Finally, all our models would benefit from some form of ground-truthing. Because statistically rigorous model validation is highly cost/labor intensive, field verification efforts should be encouraged for crews who are surveying a particular area and are able to check survey locations with GPS against the predicted habitat to confirm presence or absence.

Table 2. Colorado special status plant species lacking species distribution models.

Scientific Name	Common Name	USFS	BLM	PGCN
<i>Amsonia jonesii</i>	Jones' bluestar		X	
<i>Aquilegia chrysantha</i> var. <i>rydbergii</i>	Rydberg's golden columbine	X	X	
<i>Astragalus detritalis</i>	debris milkvetch		X	
<i>Astragalus duchesnensis</i>	Duchesne milkvetch		X	
<i>Astragalus leptaleus</i>	park milkvetch	X		
<i>Astragalus musiniensis</i>	Ferron's milkvetch		X	
<i>Astragalus proximus</i>	Aztec milkvetch	X		
<i>Astragalus ripleyi</i>	Ripley's milkvetch	X	X	
<i>Astragalus sesquiflorus</i>	sandstone milkvetch		X	
<i>Botrychium campestre</i>	Iowa moonwort, prairie moonwort	X		
<i>Botrychium lineare</i>	Narrowleaf grape fern			Tier 2
<i>Calochortus flexuosus</i>	winding mariposa lily	X		
<i>Carex diandra</i>	lesser paniced sedge	X		
<i>Chenopodium cycloides</i>	sandhill goosefoot	X		
<i>Cryptantha caespitosa</i>	tufted cryptantha		X	
<i>Cryptantha osterhoutii</i>	Osterhout's cryptantha		X	
<i>Cryptogramma stelleri</i>	fragile rockbrake		X	
<i>Cymopterus duchesnensis</i>	Uinta Basin springparsley		X	
<i>Cypripedium parviflorum</i>	lesser yellow lady's slipper	X		
<i>Descurainia torulosa</i>	mountain tansymustard	X		
<i>Drosera anglica</i>	English sundew	X		
<i>Drosera rotundifolia</i>	roundleaf sundew	X		
<i>Epipactis gigantea</i>	stream orchid, giant helleborine	X		
<i>Eriogonum acaule</i>	singlestem buckwheat		X	
<i>Eriogonum contortum</i>	grand buckwheat		X	
<i>Eriogonum ephedroides</i>	ephedra buckwheat		X	
<i>Eriogonum exilifolium</i>	dropleaf buckwheat	X		
<i>Eriogonum tumulosum</i>	Woodside buckwheat		X	
<i>Eriogonum viridulum</i>	clay hill buckwheat		X	
<i>Eriophorum chamissonis</i>	Chamisso's cottongrass	X		

Scientific Name	Common Name	USFS	BLM	PGCN
<i>Frasera paniculata</i>	tufted frasera		X	
<i>Gentianella tortuosa</i>	Cathedral Bluff dwarf gentian		X	
<i>Gilia (Aliciella) stenothyrsa</i>	Uinta Basin gilia		X	
<i>Ipomopsis aggregata ssp. weberi</i>	scarlet gilia	X		Tier 2
<i>Kobresia simpliciuscula</i>	simple bog sedge	X		
<i>Lomatium latilobum</i>	Canyonlands biscuitroot		X	
<i>Malaxis monophyllos var. brachypoda</i>	white adder's-mouth orchid	X		
<i>Neoparrya lithophila</i>	Bill's neoparrya	X	X	
<i>Parthenium ligulatum</i>	Colorado feverfew		X	
<i>Pediomelum aromaticum</i>	aromatic Indian breadroot		X	
<i>Penstemon harringtonii</i>	Harrington's beardtongue	X	X	
<i>Rubus arcticus ssp. acaulis</i>	dwarf raspberry	X		
<i>Salix myrtilifolia</i>	blueberry willow	X		
<i>Selaginella selaginoides</i>	club spikemoss	X		
<i>Sisyrinchium pallidum</i>	pale blue-eyed grass		X	
<i>Sphaeromeria capitata</i>	rock tansy		X	
<i>Sphagnum angustifolium</i>	sphagnum	X		
<i>Sphagnum balticum</i>	Baltic sphagnum	X		
<i>Townsendia strigosa</i>	hairy Townsend daisy		X	
<i>Trichophorum pumilum</i>	Rolland's bulrush		X	
<i>Triteleia grandiflora</i>	largeflower triteleia	X		
<i>Utricularia minor</i>	lesser bladderwort	X		
<i>Viola selkirkii</i>	Selkirk's violet	X		

Future data development

Additional or improved data

For species which proved difficult to model satisfactorily, more detailed environmental data layers could help refine modeled habitat. For example, an expanded detailed rock outcrop layer was developed for *Mimulus gemmiparus*, but also proved useful for two other species. More challenging to produce, a high-quality dataset depicting areas where groundwater comes to the surface (small seeps and springs) would be useful for *M. gemmiparus* as well *Oenothera acutissima*, *Draba weberi*, and species of hanging garden or alcove environments. Production of these and similar enhanced data layers was outside of the scope of this project.

Environmental inputs used in this project can be grouped into a few basic types:

Substrate (geologic and soil factors)

Substrate datasets are derived from ground-based mapping, and the original data is often quite old. There is essentially only a single soils dataset available, which has been revised and manipulated, but retains much of the coarseness of the original effort. Surface geology has somewhat better quality, but available data is often incomplete within a species range. Statewide maps digitized from

original hand-drawn publications are the primary source of coverage at a regional or national scale. Many smaller areas (1:24000 to 1:500000 scale quads) have been mapped, but not all are available in digital format, and those that have been digitized are often not edge matched with adjoining quads.

For some future models that cover a limited range, it may be worthwhile to digitize particular geologic units from fine-scale mapping within a limited study area, as was done for the two listed Piceance Basin *Physaria* species (Decker et al. 2013). Or, as with the rock outcrop layer for *M. gemmiparus*, and the CNHP-developed shale barrens layer, identify important factors which can be fairly quickly mapped from aerial imagery over large areas.

New soils data is unlikely to be readily available in the foreseeable future and would be difficult to map. Modelers should keep an eye out for new interpretations of the older data that might prove useful.

Climatic (temperature and precipitation patterns)

Climate datasets are plentiful; the ways in which precipitation and temperature models can be partitioned into time slices ranging from minutes to millennia seem endless, and a number of different observational datasets have been used in climate modeling (e.g., tree-ring or midden data in addition to historical observations). It is important, however, to remember that full coverage datasets are interpolated from point observations, and elevation is a primary component of the process. Areas of complex topography, including much of western Colorado, have highly variable patterns of precipitation, making accurate interpolation difficult. Microclimatic patterns will remain nearly impossible to model over large areas for the foreseeable future.

Topographic (elevation and related factors including slope, aspect, and other terrain descriptors)

Topographic datasets are generally derived from the Digital Elevation Model and are consequently highly correlated with each other and with climate models. If a particular topographic pattern associated with a species distribution can be identified and quantified, additional datasets can be generated with comparative ease. Many researchers develop new algorithms to calculate topographic indices, and these are widely available online.

Biophysical (vegetation and hydrology related)

Landcover mapping of widespread vegetation types is typically done by classification of aerial or satellite imagery. Available statewide vegetation datasets have acceptable accuracy at landscape scales but are very often incorrect at the very fine scale level pertinent to small plant populations. Similarly, small-patch vegetation types (e.g., wetlands) are poorly mapped by satellite image classification, and are best represented by hand-mapped polygons on high resolution aerial imagery, or ground-based mapping. Categorical versions of landcover datasets are useful to identify association with particular vegetation types or patterns of occurrence on the landscape that can then be addressed with additional data development.

Most hydrologic data is developed and distributed through the National Hydrography Dataset and related products. The very large vector digital datasets can be challenging to manipulate, and errors in the data are common. Again, this data is acceptable at landscape scales but may not reflect actual conditions near rare plant populations in a useful way.

Disturbance (both natural and anthropogenic)

A single dataset incorporating many types of anthropogenic disturbance was used during this project, and individual disturbance types could be broken out of the index if needed. Natural disturbances such as fire, flooding, drought, and so forth are typically addressed through their effects on landcover or hydrologic patterns.

An ever-present challenge in modeling rare plant species is the relative size of the plant itself in comparison with the resolution and precision of environmental data. Our models were produced using a resolution of 30 x 30 m cells. For some species, the entire known population could fit in a single cell this size. For regional species-of-concern survey and landscape-scale planning, this is an adequate resolution. However, further refinement of local habitat extent is frequently desired for locations where management decision will affect potential habitat. In such instances, our models can serve as a baseline for re-running a restricted area model using data at a finer resolution (e.g., 10 m cells) that will help resource managers narrow the area of interest. Of course, the production of 10 m resolution data does not mean that values at a particular point on the ground are more accurate than those of coarser data. Any model is only as good as its poorest input. Finer interpolation of values measured at selected points will never replace the expert botanist's search image in the real world but can suggest that areas never before considered as suitable might be worth a look.

In general, modelers should always consider how to represent micro-habitat factors that are important to individual species at a landscape scale. The distance-to-substrate is one such method. Discussions between the modeler and botanists or others familiar with the species are central to the model development process.

Refined models

Although many species were modeled with a single run, nearly all species would be better served with one or more additional model iterations; the initial run often suggests environmental factors that might be improved with additional data. Most rare plant species are little studied, so interpretation of multiple model results could also point to important, but previously unsuspected, factors controlling a species distribution. For example, the modeled pattern of immediately adjacent but rarely overlapping local habitat between *Penstemon acaulis* var. *yampaensis* and *Penstemon scariosus* var. *cyanomontanus* in Moffat County (Figure 1) is an interesting ecological insight at a scale not often considered by botanists. Elevational separation between these two penstemon species means that *Penstemon scariosus* var. *cyanomontanus* experiences higher, generally cooler, and slightly wetter conditions in comparison with its near neighbor *P. yampaensis*, found in lower, drier habitat where temperature extremes are slightly more pronounced.




Figure 1. Modeled suitable habitat for two rare penstemon species in Moffat County, Colorado.

Future conditions

Climate change is an immediate concern in the management of rare plant species. Previous evaluations of Colorado’s individual rare plant species vulnerability to changing climatic conditions have largely concluded that virtually all are highly vulnerable (CPW 2015, CNHP 2015). These vulnerability assessments were produced using generalized techniques that were not able to assess more detailed species-specific information. Unquestionably conditions are changing, and it would be expedient to generate models of species distribution under future conditions. Maxent includes options for using projected future climate data that can be used to investigate the effects of changing climate on species distributions. A test of the procedure for making a projected model was completed for *Draba smithii* (Appendix D).

Conclusion

During this project, CNHP staff developed an efficient and repeatable method for producing high quality predictive habitat models for rare plant species in Colorado. Many of the environmental input layers at the statewide level developed for this effort can be used in future modeling efforts. We now have a much better idea of what it takes to develop a collection of models, factors that might make modeling a single species difficult, and where cost-savings from production of multiple models can be realized. With the addition of these models into CODEX, the modeled distribution for almost all of the plant species listed as Plants of Greatest Conservation Need will be represented in the statewide conservation data

sharing platform, improving environmental review. The binary models, along with the full probability models provided to CNAP, will aid in the conservation of these species through their use in prioritizing and planning for conservation activities. Colorado's Plants of Greatest Conservation Need are critically under-studied and under-surveyed. This modeling work amplifies our knowledge, building upon decades of field work preserved in CNHP's database, and advances our understanding of species habitat.

References

- Colorado Natural Heritage Program [CNHP]. 2015. Climate Change Vulnerability Assessment for Colorado Bureau of Land Management. K. Decker, L. Grunau, J. Handwerk, and J. Siemers, editors. Colorado Natural Heritage Program, Colorado State University, Fort Collins, Colorado.
- Colorado Natural Heritage Program [CNHP]. 2021. Biodiversity Tracking and Conservation System (BIOTICS). Colorado Natural Heritage Program, Colorado State University, Fort Collins, Colorado.
- Colorado Parks and Wildlife [CPW]. 2015. State Wildlife Action Plan: A Strategy for Conserving Wildlife in Colorado. Appendix A: Rare Plants. Denver, Colorado.
- Decker, K. M. Fink, J. Handwerk, G. Smith. 2013. Iterative distribution modeling for two endemic plants of the northern Piceance Basin. Colorado Natural Heritage Program, Colorado State University, Fort Collins, Colorado.
- Elith, J., S.J. Phillips, T. Hastie, M. Dudik, Y.E. Chee, and C.J. Yates. 2011. A statistical explanation of MaxEnt for ecologists. *Diversity and Distributions* 17:43-57
- ESRI Inc. 1999-2015. ArcGIS 10.4.1 for Desktop. Esri, Redlands, California, USA.
- Handwerk, J.E., D. G. Malone, and J. C. Emerick. 2017. Surveys for *Pediocactus knowltonii* on Southern Ute Indian Tribal lands, Colorado. Colorado Natural Heritage Program, Colorado State University, Fort Collins, Colorado.
- New Mexico Rare Plant Technical Council. 1999. New Mexico Rare Plants. Albuquerque, NM: New Mexico Rare Plants Home Page. <https://nmrareplants.unm.edu>
- NRCS. 2012. Gridded Soil Survey Geographic (gSSURGO) Database for Colorado. United States Department of Agriculture, Natural Resources Conservation Service. Available at <http://datagateway.nrcs.usda.gov>. November 20, 2012 (FY2013 official release).
- Pearson, R.G. 2010. Species' Distribution Modeling for Conservation Educators and Practitioners. Network of Conservation Educators and Practitioners, Center for Biodiversity and Conservation, American Museum of Natural History *Lessons in Conservation*, Vol. 3, pp. 54-89
<https://www.amnh.org/content/download/141368/2285424/file/ncep.amnh.org/linc/>
- Phillips, S.J., M. Dudik, and R.E. Schapire. 2004. A maximum entropy approach to species distribution modeling. Pages 655-662 in *Proceedings of the 21st International Conference on Machine Learning*. ACM Press, New York.
- Phillips, S.J., R.P. Anderson, and R.E. Schapire. 2006. Maximum entropy modeling of species geographic distributions. *Ecological Modelling* 190:231-259.
- Phillips, S.J., M. Dudík, R.E. Schapire. 2020. Maxent software for modeling species niches and distributions (Version 3.4.3). Available from:
http://biodiversityinformatics.amnh.org/open_source/maxent/

SEINet Network of North American Plant Herbaria. 2021. Arizona-New Mexico Chapter Portal. Arizona State University Biodiversity Knowledge Integration Center, Tempe, Arizona; University of New Mexico Herbarium, Albuquerque, New Mexico. <http://swbiodiversity.org/seinet/index.php>

Appendix A: Data sources

Data layers used as environmental input factors in Maxent and deductive models. Raster names are as shown in model result outputs and metadata. Metadata entry gives the full layer name, source or sources, and a brief explanation of data processing and interpretation.

Raster name	Metadata entry
aprilmintemp (apr_mintemp)	<p>Environmental Input layer: April Minimum Temperature Source citation: Peter E. Thornton, National Center for Atmospheric Research. 2002. Daymet: Climatological Summaries for the Conterminous United States, 1980-1997. Monthly Minimum Temperature; April. Raster digital data, 1 km resolution. http://www.daymet.org</p> <p>Daymet Monthly Minimum Temperature in April for Colorado. Units are degrees Celsius. Daymet represents an average from 1980 - 1997, at 1 kilometer resolution. Raster was down sampled to 30m, re-projected and snapped to be compatible with other environmental inputs.</p>
average_clay	<p>Environmental Input layer: Average % clay in soil Source citations: Miller, D.A. and R.A. White. 1998. A Conterminous United States Multi-Layer Soil Characteristics Data Set for Regional Climate and Hydrology Modeling. Data derived from Natural Resources Conservation Service (NRCS) State Soil Geographic database (STATSGO). Tabular digital data. http://www.essc.psu.edu/soil_info/index.cgi?soil_data&conus NRCS. 1994. State Soil Geographic (STATSGO) data base for Colorado. United States Department of Agriculture, Natural Resources Conservation Service. Available at http://datagateway.nrcs.usda.gov NRCS. 2012. Gridded Soil Survey Geographic (gSSURGO) Database for Colorado. United States Department of Agriculture, Natural Resources Conservation Service. Available at http://datagateway.nrcs.usda.gov. November 20, 2012 (FY2013 official release).</p> <p>Values are supplied for each of 11 standard soil levels, down to 2.5m. Values of 0 are really NoData. Non-zero values were averaged from layers 1 - 6 as a proxy for percent clay composition down to 60cm soil depth. Due to the coarse scale of STATSGO (NRCS 1994) and the incomplete nature of SSURGO (NRCS 2012) in Colorado, all soil inputs used in CODEX PGCN models were based on the combined STATSGO-SSURGO version. Tabular data was joined to the combined STATSGO-SSURGO vector digital dataset for Colorado and exported as a 30m raster.</p>

Raster name	Metadata entry
average_sand	<p>Environmental Input layer: Average % sand in soil</p> <p>Source citations: Miller, D.A. and R.A. White. 1998. A Conterminous United States Multi-Layer Soil Characteristics Data Set for Regional Climate and Hydrology Modeling. Data derived from Natural Resources Conservation Service (NRCS) State Soil Geographic database (STATSGO). Tabular digital data. http://www.essc.psu.edu/soil_info/index.cgi?soil_data&conus</p> <p>NRCS. 1994. State Soil Geographic (STATSGO) data base for Colorado. United States Department of Agriculture, Natural Resources Conservation Service. Available at http://datagateway.nrcs.usda.gov</p> <p>NRCS. 2012. Gridded Soil Survey Geographic (gSSURGO) Database for Colorado. United States Department of Agriculture, Natural Resources Conservation Service. Available at http://datagateway.nrcs.usda.gov. November 20, 2012 (FY2013 official release).</p> <p>Values are supplied for each of 11 standard soil levels, down to 2.5m. Values of 0 are really NoData. Non-zero values were averaged from layers 1 - 6 as a proxy for percent sand composition down to 60cm soil depth. Due to the coarse scale of STATSGO (NRCS 1994) and the incomplete nature of SSURGO (NRCS 2012) in Colorado, all soil inputs used in CODEX PGCN models were based on the combined STATSGO-SSURGO version. Tabular data was joined to the combined STATSGO-SSURGO vector digital dataset for Colorado and exported as a 30m raster.</p>
average_silt	<p>Environmental Input layer: Average % silt in soil</p> <p>Source citations: Miller, D.A. and R.A. White. 1998. A Conterminous United States Multi-Layer Soil Characteristics Data Set for Regional Climate and Hydrology Modeling. Data derived from Natural Resources Conservation Service (NRCS) State Soil Geographic database (STATSGO). Tabular digital data. http://www.essc.psu.edu/soil_info/index.cgi?soil_data&conus</p> <p>NRCS. 1994. State Soil Geographic (STATSGO) data base for Colorado. United States Department of Agriculture, Natural Resources Conservation Service. Available at http://datagateway.nrcs.usda.gov</p> <p>NRCS. 2012. Gridded Soil Survey Geographic (gSSURGO) Database for Colorado. United States Department of Agriculture, Natural Resources Conservation Service. Available at http://datagateway.nrcs.usda.gov. November 20, 2012 (FY2013 official release).</p> <p>Values are supplied for each of 11 standard soil levels, down to 2.5m. Values of 0 are really NoData. Non-zero values were averaged from layers 1 - 6 as a proxy for percent silt composition down to 60cm soil depth. Due to the coarse scale of STATSGO (NRCS 1994) and the incomplete nature of SSURGO (NRCS 2012) in Colorado, all soil inputs used in CODEX PGCN models were based on the combined STATSGO-SSURGO version. Tabular data was joined to the combined STATSGO-SSURGO vector digital dataset for Colorado and exported as a 30m raster.</p>

Raster name	Metadata entry
avg_firstfrost	<p>Environmental Input layer: Average First Frost</p> <p>Source citations: Thornton, PE, MM Thornton, BW Mayer, N Wilhelmi, Y Wei, RB Cook. 2012. Daymet: Daily surface weather on a 1km grid for North America,1980-2012. Acquired online (http://daymet.ornl.gov/) on 02/20/2014 from Oak Ridge National Laboratory Distributed Active Archive Center, Oak Ridge, Tennessee, U.S.A. http://dx.doi.org/doi:10.3334/ORNLDAAC/Daymet_V2 via the USGS Geo Data Portal (http://cida.usgs.gov/gdp/).</p> <p>Daymet Daily surface weather for Colorado. Units are degrees Celsius. Daymet represents an average from 1980 - 2012, at 1 kilometer resolution. The earliest (Julian) day of each year during summer/fall on which the minimum temperature was <= 0°C was averaged. Raster was down sampled to 30m, re-projected and snapped to be compatible with other environmental inputs.</p>
avg_lastfrost	<p>Environmental Input layer: Average Last Frost</p> <p>Source citations: Thornton, PE, MM Thornton, BW Mayer, N Wilhelmi, Y Wei, RB Cook. 2012. Daymet: Daily surface weather on a 1km grid for North America,1980-2012. Acquired online (http://daymet.ornl.gov/) on 02/20/2014 from Oak Ridge National Laboratory Distributed Active Archive Center, Oak Ridge, Tennessee, U.S.A. http://dx.doi.org/doi:10.3334/ORNLDAAC/Daymet_V2 via the USGS Geo Data Portal (http://cida.usgs.gov/gdp/).</p> <p>Daymet Daily surface weather for Colorado. Units are degrees Celsius. Daymet represents an average from 1980 - 2012, at 1 kilometer resolution. The latest (Julian) day of each year during spring/summer on which the minimum temperature was <= 0°C was averaged. Raster was down sampled to 30m, re-projected and snapped to be compatible with other environmental inputs.</p>
co_ned30m	<p>Environmental Input layer: 30m Digital Elevation Model for Colorado</p> <p>Source citations: U.S. Geological Survey. 2006. 30m Digital Elevation Model for Colorado. Raster digital data. http://seamless.usgs.gov/website/seamless/viewer.php</p> <p>Raster was re-projected, clipped to the Colorado state boundary extent with a minimum border of 8.5km, and used as base extent and snap reference for all environmental inputs.</p>
colo_bps	<p>Environmental Input layer: Biophysical Settings (BPS)</p> <p>Source citation: LANDFIRE, Earth Resources Observation and Science Center (EROS), U.S. Geological Survey. 2020. Biophysical Settings. LANDFIRE Remap 2016, CONUS. Raster digital data. www.landfire.gov</p> <p>BPS represents the vegetation system that may have been dominant on the landscape prior to Euro-American settlement. Data for continental US was downloaded, reprojected, clipped to reference extent, and snapped to be compatible with other environmental inputs. This is a categorial dataset.</p>

Raster name	Metadata entry
colo_evt	<p>Environmental Input layer: Existing Vegetation Type (EVT) Source citation: LANDFIRE, Earth Resources Observation and Science Center (EROS), U.S. Geological Survey. 2020. Existing Vegetation Type. LANDFIRE Remap 2016, CONUS. Raster digital data. www.landfire.gov</p> <p>EVT represents the current distribution of the terrestrial ecological systems classification, developed by NatureServe for the western hemisphere, through 2016. A terrestrial ecological system is defined as a group of plant community types (associations) that tend to co-occur within landscapes with similar ecological processes, substrates, and/or environmental gradients. Data for continental US was downloaded, reprojected, clipped to reference extent, and snapped to be compatible with other environmental inputs. This is a categorical dataset.</p>
colo_geol	<p>Environmental Input layer: Colorado Surface Geology Source citation: Tweto, Ogden. 1979. Geologic map of Colorado. Vector digital data, 1:500,000. U.S. Geological Survey. ftp://greenwood.cr.usgs.gov/pub/open-file-reports/ofr-92-0507/</p> <p>Original ARC/INFO data was converted to ArcGIS shapefile and projected to UTM_NAD83_Zone13. Shapefile was converted to a 30m raster, using the CELL_CENTER cell assignment type, and snapped to be compatible with other environmental inputs. Formation name abbreviation was retained. This is a categorical dataset.</p>
colo_nvc_veg	<p>Environmental Input layer: National Vegetation Classification (NVC) Source citation: LANDFIRE, Earth Resources Observation and Science Center (EROS), U.S. Geological Survey. 2020. National Vegetation Classification (NVC). LANDFIRE Remap 2016, CONUS. Raster digital data. www.landfire.gov</p> <p>National Vegetation Classification (NVC) represents the current distribution of vegetation groups within the U.S. National Vegetation Classification System ([version 2.0] http://usnvc.org/). Data for continental US was downloaded, reprojected, clipped to reference extent, and snapped to be compatible with other environmental inputs. This is a categorical dataset.</p>
colo_nvcveg_nodev	<p>Environmental Input layer: National Vegetation Classification (NVC), not including developed areas Source citation: LANDFIRE, Earth Resources Observation and Science Center (EROS), U.S. Geological Survey. 2020. National Vegetation Classification (NVC). LANDFIRE Remap 2016, CONUS. Raster digital data. www.landfire.gov</p> <p>National Vegetation Classification (NVC) represents the current distribution of vegetation groups within the U.S. National Vegetation Classification System ([version 2.0] http://usnvc.org/). Data for continental US was downloaded, reprojected, clipped to reference extent, and snapped to be compatible with other environmental inputs. Cells with attributes of Developed-Low Intensity, Developed-Medium Intensity, Developed-High Intensity, and Developed-Roads were reclassified to NoData. This is a categorical dataset.</p>

Raster name	Metadata entry
dist_30plus_slope	<p>Environmental Input layer: Distance to slopes of more than 30 degrees Source citation: Colorado Natural Heritage Program. 2021. Unpublished data using USGS 30m DEM. Raster digital data.</p> <p>Slope in degrees raster (derived from U.S. Geological Survey. 2006. 30m Digital Elevation Model for Colorado) was reclassified to identify cells greater than or equal to 30 degrees. Cells meeting this criteria were used as the input for generating a Euclidian Distance raster.</p>
dist_badlands	<p>Environmental Input layer: Distance to Intermountain Basins Cliff Scree & Badland Sparse Vegetation Source citation: LANDFIRE, Earth Resources Observation and Science Center (EROS), U.S. Geological Survey. 2020. National Vegetation Classification (NVC). LANDFIRE Remap 2016, CONUS. Raster digital data. www.landfire.gov</p> <p>National Vegetation Classification (NVC) represents the current distribution of vegetation groups within the U.S. National Vegetation Classification System ([version 2.0] http://usnvc.org/). Original raster data was re-projected, then clipped and snapped to be compatible with other environmental inputs. This dataset was then reclassified to retain NVC_NAME = Inter-Mountain Basins Cliff Scree & Badland Sparse Vegetation, while all other types were classified as NoData. This raster was then used as the input feature source data for generating a Euclidian Distance raster.</p>
dist_brownsPk	<p>Environmental Input layer: Distance to Browns Park Formation Source citation: Tweto, Ogden. 1979. Geologic map of Colorado. Vector digital data, 1:500,000. U.S. Geological Survey. ftp://greenwood.cr.usgs.gov/pub/open-file-reports/ofr-92-0507/</p> <p>Original ARC/INFO data was converted to ArcGIS shapefile and projected to UTM_NAD83_Zone13.</p> <p>Polygons with NAME attribute Tbp were selected, exported, and used as the input feature source data for generating a Euclidian Distance raster.</p>
dist_carlgrhngnan	<p>Environmental Input layer: Distance to Carlile Shale, Greenhorn Limestone, and Graneros Shale Source citation: Tweto, Ogden. 1979. Geologic map of Colorado. Vector digital data, 1:500,000. U.S. Geological Survey. ftp://greenwood.cr.usgs.gov/pub/open-file-reports/ofr-92-0507/</p> <p>Original ARC/INFO data was converted to ArcGIS shapefile and projected to UTM_NAD83_Zone13.</p> <p>Polygons with NAME attribute Kcg were selected, exported, and used as the input feature source data for generating a Euclidian Distance raster.</p>

Raster name	Metadata entry
dist_copl_pj	<p>Environmental Input layer: Distance to Colorado Plateau Pinyon-Juniper Woodland vegetation type vegetation type Source citation: LANDFIRE, Earth Resources Observation and Science Center (EROS), U.S. Geological Survey. 2020. Existing Vegetation Type (EVT). LANDFIRE Remap 2016, CONUS. Raster digital data. www.landfire.gov</p> <p>Existing Vegetation Type represents complexes of plant communities representing NatureServe's terrestrial Ecological Systems classification. Data for continental US was downloaded, reprojected, clipped to reference extent, and snapped to be compatible with other environmental inputs. Cells attributed as EVT_Name = Distance to Colorado Plateau Pinyon-Juniper Woodland were used as the input feature source data for generating a Euclidian Distance raster.</p>
dist_cutler	<p>Environmental Input layer: Distance to Cutler Formation Source citation: Originator: Day, W.C., Green, G.N., Knepper, D.H., and Phillips, R.C. 2000. Spatial Geologic Data Model for the Gunnison, Grand Mesa, Uncompahgre National Forests Mineral Resource Assessment Area, Southwestern Colorado and Digital Data for the Leadville, Montrose, Durango, and the Colorado Parts of the Grand Junction, Moab, and Cortez 1° x 2° Geologic Maps. Vector digital data, 1:250,000. U.S. Geological Survey. https://pubs.usgs.gov/of/1999/ofr-99-0427/</p> <p>Original ARC/INFO data was converted to ArcGIS shapefile and projected to UTM_NAD83_Zone13. Units sharing boundaries across quads were dissolved.</p> <p>Polygons with NAME attribute Pc were selected, exported, and used as the input feature source data for generating a Euclidian Distance raster.</p>
dist_dry_union	<p>Environmental Input layer: Distance to Dry Union Formation Source citation: Tweto, Ogden. 1979. Geologic map of Colorado. Vector digital data, 1:500,000. U.S. Geological Survey. ftp://greenwood.cr.usgs.gov/pub/open-file-reports/ofr-92-0507/</p> <p>Original ARC/INFO data was converted to ArcGIS shapefile and projected to UTM_NAD83_Zone13.</p> <p>Polygons with NAME attribute Td were selected, exported, and used as the input feature source data for generating a Euclidian Distance raster.</p>
dist_fault	<p>Environmental Input layer: Distance to fault Source citation: Green, G.N., 1992, CO_Geology_Faults, The Digital Geologic Map of Colorado in ARC/INFO Format. Vector digital data, 1:500,000.U.S. Geological Survey Open-File Report 92-0507A-O, 9 p. and 14 magnetic disks; online at http://pubs.usgs.gov/of/1992/ofr-92-0507/.</p> <p>Original ARC/INFO data was converted to ArcGIS shapefile and projected to UTM_NAD83_Zone13 and used as the input feature source data for generating a Euclidian Distance raster.</p>

Raster name	Metadata entry
dist_fr_hogback_shales	<p>Environmental Input layer: Distance to shale and sandstone units forming the Front Range (Dakota) hogback north of Colo Spgs. Source citation: Tweto, Ogden. 1979. Geologic map of Colorado. Vector digital data, 1:500,000. U.S. Geological Survey. ftp://greenwood.cr.usgs.gov/pub/open-file-reports/ofr-92-0507/</p> <p>Original ARC/INFO data was converted to ArcGIS shapefile and projected to UTM_NAD83_Zone13.</p> <p>Polygons with NAME attribute Kc, Kpl, KJdr, KJds, P&if, @Pll, @&lf, @Pjs were selected, exported, and used as the input feature source data for generating a Euclidian Distance raster.</p>
dist_fr_outcrops	<p>Environmental Input layer: Distance to Front Range Rock Outcrops, third (expanded) version (150m) Source citation: Decker, Karin. 2021. Rock outcrops in the Front Range, but not including the mountain front hogbacks. Vector digital data. Approximately 1:15,000. Colorado Natural Heritage Program, unpublished data.</p> <p>Points marking the approximate location of (primarily granitic) rock outcrops were digitized using high (but variable) resolution World Imagery from Environmental Systems Research Institute (ESRI) and checked against Google Earth views as needed. The point shapefile was buffered to 150m radius, with overlaps dissolved, and used as the input feature source data for generating a Euclidian Distance raster.</p>
dist_granitic_Yg	<p>Environmental Input layer: Distance to granitic rocks of 1,400-MY age group Source citation: Tweto, Ogden. 1979. Geologic map of Colorado. Vector digital data, 1:500,000. U.S. Geological Survey. ftp://greenwood.cr.usgs.gov/pub/open-file-reports/ofr-92-0507/</p> <p>Original ARC/INFO data was converted to ArcGIS shapefile and projected to UTM_NAD83_Zone13.</p> <p>Polygons with NAME attribute Yg were selected, exported, and used as the input feature source data for generating a Euclidian Distance raster.</p>
dist_greasewood	<p>Environmental Input layer: Distance to Inter-Mountain Basins Greasewood Flat ecological system Source citation: US Geological Survey. 2011. GAP/LANDFIRE National Terrestrial Ecosystems. Raster digital data. http://gis1.usgs.gov/csas/gap/viewer/land_cover/Map.aspx</p> <p>Original raster data was re-projected, then clipped and snapped to be compatible with other environmental inputs. This dataset was then reclassified to retain Ecolsys_LU = Inter-Mountain Basins Greasewood Flat, while all other types were classified as NoData. This raster was then used as the input feature source data for generating a Euclidian Distance raster.</p>

Raster name	Metadata entry
dist_greenriv	<p>Environmental Input layer: Distance to Green River Formation Source citation: Tweto, Ogden. 1979. Geologic map of Colorado. Vector digital data, 1:500,000. U.S. Geological Survey. ftp://greenwood.cr.usgs.gov/pub/open-file-reports/ofr-92-0507/</p> <p>Original ARC/INFO data was converted to ArcGIS shapefile and projected to UTM_NAD83_Zone13.</p> <p>Polygons with NAME attribute Tg were selected, exported, and used as the input feature source data for generating a Euclidian Distance raster.</p>
dist_herb_riparian	<p>Environmental Input layer: Distance to combined WGP floodplain and herbaceous riparian Source citations: US Geological Survey. 2011. GAP/LANDFIRE National Terrestrial Ecosystems. Raster digital data. http://gis1.usgs.gov/csas/gap/viewer/land_cover/Map.aspx Colorado Division of Wildlife. 2004. Colorado Vegetation Classification Project; Statewide Mosaic. Raster digital data.</p> <p>Original raster data was re-sampled, re-projected then clipped and snapped to be compatible with other environmental inputs. The GAP dataset was then reclassified to retain Ecolsys_LU = Western Great Plains Floodplain, while all other types were classified as NoData. The CDOW dataset was reclassified to retain CLASS_NM = Herbaceous Riparian. The two rasters were added, and all non-zero values retained. This raster was then used as the input feature source data for generating a Euclidian Distance raster.</p>
dist_hermosa	<p>Environmental Input layer: Distance to Hermosa Formation Source citation: Originator: Day, W.C., Green, G.N., Knepper, D.H., and Phillips, R.C. 2000. Spatial Geologic Data Model for the Gunnison, Grand Mesa, Uncompahgre National Forests Mineral Resource Assessment Area, Southwestern Colorado and Digital Data for the Leadville, Montrose, Durango, and the Colorado Parts of the Grand Junction, Moab, and Cortez 1° x 2° Geologic Maps. Vector digital data, 1:250,000. U.S. Geological Survey. https://pubs.usgs.gov/of/1999/ofr-99-0427/</p> <p>Original ARC/INFO data was converted to ArcGIS shapefile and projected to UTM_NAD83_Zone13. Units sharing boundaries across quads were dissolved.</p> <p>Polygons with NAME attribute Ph, Php, and Phu were selected, exported, and used as the input feature source data for generating a Euclidian Distance raster.</p>

Raster name	Metadata entry
dist_ignmet_xy	<p>Environmental Input layer: Distance to Igneous and Metamorphic Rocks of Precambrian Age Source citation: Tweto, Ogden. 1979. Geologic map of Colorado. Vector digital data, 1:500,000. U.S. Geological Survey. ftp://greenwood.cr.usgs.gov/pub/open-file-reports/ofr-92-0507/</p> <p>Original ARC/INFO data was converted to ArcGIS shapefile and projected to UTM_NAD83_Zone13.</p> <p>Polygons with NAME attribute Xb, Xfh, Xq, Yp, Yg, Xg, Xm, or Yxg were selected, exported, and used as the input feature source data for generating a Euclidian Distance raster.</p>
dist_IMB_mont_SB	<p>Environmental Input layer: Distance to Inter-Mountain Basins Montane Sagebrush Steppe vegetation type Source citation: LANDFIRE, Earth Resources Observation and Science Center (EROS), U.S. Geological Survey. 2020. Existing Vegetation Type (EVT). LANDFIRE Remap 2016, CONUS. Raster digital data. www.landfire.gov</p> <p>Existing Vegetation Type represents complexes of plant communities representing NatureServe's terrestrial Ecological Systems classification. Data for continental US was downloaded, reprojected, clipped to reference extent, and snapped to be compatible with other environmental inputs. Cells attributed as EVT_Name = Inter-Mountain Basins Montane Sagebrush Steppe were used as the input feature source data for generating a Euclidian Distance raster.</p>
dist_IMB_sage	<p>Environmental Input layer: Distance to Inter-Mountain Basins Big Sagebrush Shrubland vegetation type Source citation: LANDFIRE, Earth Resources Observation and Science Center (EROS), U.S. Geological Survey. 2020. Existing Vegetation Type (EVT). LANDFIRE Remap 2016, CONUS. Raster digital data. www.landfire.gov</p> <p>Existing Vegetation Type represents complexes of plant communities representing NatureServe's terrestrial Ecological Systems classification. Data for continental US was downloaded, reprojected, clipped to reference extent, and snapped to be compatible with other environmental inputs. Cells attributed as EVT_Name = Inter-Mountain Basins Big Sagebrush Shrubland were used as the input feature source data for generating a Euclidian Distance raster.</p>
dist_leadville	<p>Environmental Input layer: Distance to units containing Leadville (& Manitou) Limestone Source citation: Tweto, Ogden. 1979. Geologic map of Colorado. Vector digital data, 1:500,000. U.S. Geological Survey. ftp://greenwood.cr.usgs.gov/pub/open-file-reports/ofr-92-0507/</p> <p>Original ARC/INFO data was converted to ArcGIS shapefile and projected to UTM_NAD83_Zone13.</p> <p>Polygons with NAME attribute M_, MD, MD_, MDO, and O_ were selected, exported, and used as the input feature source data for generating a Euclidian Distance raster.</p>

Raster name	Metadata entry
dist_mancos	<p>Environmental Input layer: Distance to Mancos Shale Source citation: Tweto, Ogden. 1979. Geologic map of Colorado. Vector digital data, 1:500,000. U.S. Geological Survey. ftp://greenwood.cr.usgs.gov/pub/open-file-reports/ofr-92-0507/</p> <p>Original ARC/INFO data was converted to ArcGIS shapefile and projected to UTM_NAD83_Zone13.</p> <p>Polygons with NAME attribute Km were selected, exported, and used as the input feature source data for generating a Euclidian Distance raster.</p>
dist_mancos_sixqd	<p>Environmental Input layer: Distance to Mancos Formation in six quad area Source citation: Originator: Day, W.C., Green, G.N., Knepper, D.H., and Phillips, R.C. 2000. Spatial Geologic Data Model for the Gunnison, Grand Mesa, Uncompahgre National Forests Mineral Resource Assessment Area, Southwestern Colorado and Digital Data for the Leadville, Montrose, Durango, and the Colorado Parts of the Grand Junction, Moab, and Cortez 1° x 2° Geologic Maps. Vector digital data, 1:250,000. U.S. Geological Survey. https://pubs.usgs.gov/of/1999/ofr-99-0427/</p> <p>Original ARC/INFO data was converted to ArcGIS shapefile and projected to UTM_NAD83_Zone13. Units sharing boundaries across quads were dissolved.</p> <p>Polygons with NAME attribute Km were selected, exported, and used as the input feature source data for generating a Euclidian Distance raster.</p>
dist_mixed_forest	<p>Environmental Input layer: Distance to mixed conifer forests vegetation type Source citation: LANDFIRE, Earth Resources Observation and Science Center (EROS), U.S. Geological Survey. 2020. Existing Vegetation Type (EVT). LANDFIRE Remap 2016, CONUS. Raster digital data. www.landfire.gov</p> <p>Existing Vegetation Type represents complexes of plant communities representing NatureServe's terrestrial Ecological Systems classification. Data for continental US was downloaded, reprojected, clipped to reference extent, and snapped to be compatible with other environmental inputs. Cells attributed as EVT_Name = Southern Rocky Mountain Dry-Mesic Montane Mixed Conifer Forest and Woodland, Southern Rocky Mountain Mesic Montane Mixed Conifer Forest and Woodland, or Inter-Mountain Basins Aspen-Mixed Conifer Forest and Woodland were used as the input feature source data for generating a Euclidian Distance raster.</p>

Raster name	Metadata entry
dist_morrison	<p>Environmental Input layer: Distance to Morrison Formation, all units Source citation: Tweto, Ogden. 1979. Geologic map of Colorado. Vector digital data, 1:500,000. U.S. Geological Survey. ftp://greenwood.cr.usgs.gov/pub/open-file-reports/ofr-92-0507/</p> <p>Original ARC/INFO data was converted to ArcGIS shapefile and projected to UTM_NAD83_Zone13.</p> <p>Polygons with NAME attribute J@mc, Jm, Jmc, Jmce, Jme, Jmj, Jmr, Jmre, Jms, Jmse, Jmw, or Jmwe were selected, exported, and used as the input feature source data for generating a Euclidian Distance raster.</p>
dist_niobrara	<p>Environmental Input layer: Distance to Niobrara Formation Source citation: Tweto, Ogden. 1979. Geologic map of Colorado. Vector digital data, 1:500,000. U.S. Geological Survey. ftp://greenwood.cr.usgs.gov/pub/open-file-reports/ofr-92-0507/</p> <p>Original ARC/INFO data was converted to ArcGIS shapefile and projected to UTM_NAD83_Zone13.</p> <p>Polygons with NAME attribute Kn were selected, exported, and used as the input feature source data for generating a Euclidian Distance raster.</p>
dist_nlcd_shrub	<p>Environmental Input layer: Distance to Shrub/Scrub and Woody Wetlands landcover type Source citation: National Land Cover Database 2011 (NLCD) 2011. Multi-Resolution Land Characteristics Consortium (MRLC) dataset. https://data.nal.usda.gov/dataset/national-land-cover-database-2011-nlcd-2011</p> <p>Data for continental US was downloaded, reprojected, clipped to reference extent, and snapped to be compatible with other environmental inputs. Cells attributed as NLCD_2011 = Shrub/Scrub or Woody Wetlands were used as the input feature source data for generating a Euclidian Distance raster.</p>
dist_nvcm06c02	<p>Environmental Input layer: Distance to Semi-Desert Nonvascular & Sparse Vascular Vegetation NVC formation Source citation: US Geological Survey. 2011. GAP/LANDFIRE National Terrestrial Ecosystems. Raster digital data. http://gis1.usgs.gov/csas/gap/viewer/land_cover/Map.aspx</p> <p>Original raster data was re-projected, then clipped and snapped to be compatible with other environmental inputs. This dataset was then reclassified to retain Formation = Semi-Desert Nonvascular & Sparse Vascular Vegetation (06.C.02), while all other types were classified as NoData. This raster was then used as the input feature source data for generating a Euclidian Distance raster.</p>

Raster name	Metadata entry
dist_parachute	<p>Environmental Input layer: Distance to Parachute Creek member of Green River Formation Source citation: Tweto, Ogden. 1979. Geologic map of Colorado. Vector digital data, 1:500,000. U.S. Geological Survey. ftp://greenwood.cr.usgs.gov/pub/open-file-reports/ofr-92-0507/</p> <p>Original ARC/INFO data was converted to ArcGIS shapefile and projected to UTM_NAD83_Zone13.</p> <p>Polygons with NAME attribute Tgp were selected, exported, and used as the input feature source data for generating a Euclidian Distance raster.</p>
dist_pb_glac	<p>Environmental Input layer: Distance to Glacial drift of Pinedale and Bull Lake age Source citation: Tweto, Ogden. 1979. Geologic map of Colorado. Vector digital data, 1:500,000. U.S. Geological Survey. ftp://greenwood.cr.usgs.gov/pub/open-file-reports/ofr-92-0507/</p> <p>Original ARC/INFO data was converted to ArcGIS shapefile and projected to UTM_NAD83_Zone13.</p> <p>Polygons with NAME attribute Qd were selected, exported, and used as the input feature source data for generating a Euclidian Distance raster.</p>
dist_pacman_soils	<p>Environmental Input layer: Distance to soils with Packera mancosana occurrences Source citations: NRCS. 2012. Gridded Soil Survey Geographic (gSSURGO) Database for Colorado. United States Department of Agriculture, Natural Resources Conservation Service. Available at http://datagateway.nrcs.usda.gov. November 20, 2012 (FY2013 official release).</p> <p>Soil units with SSURGO attributes MUKEY 507319, 507229, 502153, 501969 were selected and used as the input feature source data for generating a Euclidian Distance raster.</p>
dist_pem	<p>Environmental Input layer: Distance to Palustrine Emergent wetland types Source citation: U.S. Fish and Wildlife Service. 2016. CONUS_wet_poly_West; CO_Wetlands, National Wetlands Inventory - Version 2. Vector digital data, 1:12,000. https://www.fws.gov/wetlands/Data/Data-Download.html</p> <p>Polygons with attributes beginning with PEM were selected (query: Like PEM%) and used as the input feature source data for generating a Euclidian Distance raster.</p>

Raster name	Metadata entry
dist_qae	<p>Environmental Input layer: Distance to Quaternary alluvium and eolian deposits Source citation: Originator: Day, W.C., Green, G.N., Knepper, D.H., and Phillips, R.C. 2000. Spatial Geologic Data Model for the Gunnison, Grand Mesa, Uncompahgre National Forests Mineral Resource Assessment Area, Southwestern Colorado and Digital Data for the Leadville, Montrose, Durango, and the Colorado Parts of the Grand Junction, Moab, and Cortez 1° x 2° Geologic Maps. Vector digital data, 1:250,000. U.S. Geological Survey. https://pubs.usgs.gov/of/1999/ofr-99-0427/</p> <p>Original ARC/INFO data was converted to ArcGIS shapefile and projected to UTM_NAD83_Zone13. Units sharing boundaries across quads were dissolved.</p> <p>Polygons with NAME attribute Qae were selected, exported, and used as the input feature source data for generating a Euclidian Distance raster.</p>
dist_raton	<p>Environmental Input layer: Distance to Raton Formation Source citations: Tweto, Ogden. 1979. Geologic map of Colorado. Vector digital data, 1:500,000. U.S. Geological Survey. ftp://greenwood.cr.usgs.gov/pub/open-file-reports/ofr-92-0507/ Green, G.N., and Jones, G.E. 1997. The Digital Geologic Map of New Mexico in ARC/INFO Format. Vector digital data, 1:500,000. U.S. Geological Survey. http://pubs.usgs.gov/of/2005/1351/data/NMgeol_dd.zip</p> <p>Original ARC/INFO data was converted to ArcGIS shapefile and projected to UTM_NAD83_Zone13.</p> <p>Polygons with NAME attribute Tkr were selected, exported, merged, and used as the input feature source data for generating a Euclidian Distance raster.</p>
dist_RM_aspen	<p>Environmental Input layer: Rocky Mountain Aspen Forest and Woodland vegetation type Source citation: LANDFIRE, Earth Resources Observation and Science Center (EROS), U.S. Geological Survey. 2020. Existing Vegetation Type (EVT). LANDFIRE Remap 2016, CONUS. Raster digital data. www.landfire.gov</p> <p>Existing Vegetation Type represents complexes of plant communities representing NatureServe's terrestrial Ecological Systems classification. Data for continental US was downloaded, reprojected, clipped to reference extent, and snapped to be compatible with other environmental inputs. Cells attributed as EVT_Name = Rocky Mountain Aspen Forest and Woodland were used as the input feature source data for generating a Euclidian Distance raster.</p>

Raster name	Metadata entry
dist_RM_oakshrub	<p>Environmental Input layer: Distance to Rocky Mountain Gambel Oak-Mixed Montane Shrubland vegetation type Source citation: LANDFIRE, Earth Resources Observation and Science Center (EROS), U.S. Geological Survey. 2020. Existing Vegetation Type (EVT). LANDFIRE Remap 2016, CONUS. Raster digital data. www.landfire.gov</p> <p>Existing Vegetation Type represents complexes of plant communities representing NatureServe's terrestrial Ecological Systems classification. Data for continental US was downloaded, reprojected, clipped to reference extent, and snapped to be compatible with other environmental inputs. Cells attributed as EVT_Name = Rocky Mountain Gambel Oak-Mixed Montane Shrubland were used as the input feature source data for generating a Euclidian Distance raster.</p>
dist_rock_outcrops2	<p>Environmental Input layer: Distance to Rock Outcrops, second version (180m) Source citation: Decker, Karin. 2021. Rock outcrops in general range of Mimulus gemmiparus. Vector digital data. Approximately 1:15,000. Colorado Natural Heritage Program, unpublished data.</p> <p>Points marking the approximate location of granitic rock outcrops were digitized using high (but variable) resolution World Imagery from Environmental Systems Research Institute (ESRI) and checked against Google Earth views as needed. The point shapefile was converted to a 180m raster and used as the input feature source data for generating a Euclidian Distance raster.</p>
dist_saltbush	<p>Environmental Input layer: Distance to Intermountain Shadscale-Saltbush Scrub or Intermountain Dwarf Saltbush-Sagebrush Scrub Source citation: LANDFIRE, Earth Resources Observation and Science Center (EROS), U.S. Geological Survey. 2020. National Vegetation Classification (NVC). LANDFIRE Remap 2016, CONUS. Raster digital data. www.landfire.gov</p> <p>National Vegetation Classification (NVC) represents the current distribution of vegetation groups within the U.S. National Vegetation Classification System ([version 2.0] http://usnvc.org/). Original raster data was re-projected, then clipped and snapped to be compatible with other environmental inputs. This dataset was then reclassified to retain NVC_NAME = Intermountain Shadscale-Saltbush Scrub or Intermountain Dwarf Saltbush-Sagebrush Scrub, while all other types were classified as NoData. This raster was then used as the input feature source data for generating a Euclidian Distance raster.</p>
dist_sandstone	<p>Environmental Input layer: Distance to sandstone lithology Source citation: Horton, J.D., San Juan, C.A., and Stoesser, D.B, 2017, The State Geologic Map Compilation (SGMC) geodatabase of the conterminous United States (ver. 1.1, August 2017): U.S. Geological Survey Data Series 1052, 46 p., https://doi.org/10.3133/ds1052. The State Geologic Map Compilation (SGMC) Geodatabase of the Conterminous United States (ver. 1.1, August 2017) - SGMC_Geology. Colorado polygons are based on Tweto 1979, with additional classification fields added.</p> <p>Original downloaded polygon data was reprojected to UTM NAD83, zone 13, and clipped to the project extent. Polygons with MAJOR1 attribute = Sandstone were selected, exported, and used as the input feature source data for generating a Euclidian Distance raster.</p>

Raster name	Metadata entry
dist_saturated	<p>Environmental Input layer: Distance to saturated wetlands Source citation: U.S. Fish and Wildlife Service. 2016. CONUS_wet_poly_West; CO_Wetlands, National Wetlands Inventory - Version 2. Vector digital data, 1:12,000. https://www.fws.gov/wetlands/Data/Data-Download.html</p> <p>Polygons with attributes Palustrine Emergent Saturated (PEMB) and Palustrine Scrub-Shrub Saturated (PSSB) were selected and used as the input feature source data for generating a Euclidian Distance raster.</p>
dist_sg_bps	<p>Environmental Input layer: Distance to Western Great Plains Shortgrass Prairie BPS Source citation: LANDFIRE, Earth Resources Observation and Science Center (EROS), U.S. Geological Survey. 2020. Biophysical Settings. LANDFIRE Remap 2016, CONUS. Raster digital data. www.landfire.gov</p> <p>BPS represents the vegetation system that may have been dominant on the landscape prior to Euro-American settlement. Data for continental US was downloaded, reprojected, clipped to reference extent, and snapped to be compatible with other environmental inputs. Cells attributed as BPS Name = Western Great Plains Shortgrass Prairie were used as the input feature source data for generating a Euclidian Distance raster.</p>
dist_shale_barren	<p>Environmental Input layer: Distance to Shale Barrens Source citation: Decker, Karin. 2021. Shale Barrens of Southeastern Colorado. Vector digital data, 1:12,000. Colorado Natural Heritage Program, unpublished data.</p> <p>Polygons were digitized using high (but variable) resolution World Imagery from Environmental Systems Research Institute (ESRI) and checked against 1 x 2-degree geology quad maps (georeferenced tif images). Shale barren polygons were hand-drawn in ArcGIS 10.4 (ESRI 2015) by the photo-interpreter based on the best estimation of sparsely vegetated boundaries and used as the input feature source data for generating a Euclidian Distance raster.</p>
dist_shortgrass	<p>Environmental Input layer: Distance to Great Plains Shortgrass Prairie Source citation: LANDFIRE, Earth Resources Observation and Science Center (EROS), U.S. Geological Survey. 2020. National Vegetation Classification (NVC). LANDFIRE Remap 2016, CONUS. Raster digital data. www.landfire.gov</p> <p>National Vegetation Classification (NVC) represents the current distribution of vegetation groups within the U.S. National Vegetation Classification System ([version 2.0] http://usnvc.org/). Data for continental US was downloaded, reprojected, clipped to reference extent, and snapped to be compatible with other environmental inputs. Cells attributed as NVC Name = Great Plains Shortgrass Prairie were used as the input feature source data for generating a Euclidian Distance raster.</p>

Raster name	Metadata entry
dist_springs	<p>Environmental Input layer: Distance to springs Source citation: Ledbetter, Jeri D., MGIS, Lawrence E. Stevens, PhD, Abraham Springer, PhD, and Benjamin Brandt, MGIS. 2014. Springs Inventory Database. Online Database. Springs and Springs-Dependent Species Database. Vers. 1.0. Springs Stewardship Institute, springsdata.org.</p> <p>Original kml data was converted to ArcGIS geodatabase and points were projected to UTM_NAD83_Zone13. Points were used as the input feature source data for generating a Euclidian Distance raster.</p>
dist_srm_pj	<p>Environmental Input layer: Distance to Southern Rocky Mountain Pinyon-Juniper Woodland Source citation: LANDFIRE, Earth Resources Observation and Science Center (EROS), U.S. Geological Survey. 2020. Existing Vegetation Type (EVT). LANDFIRE Remap 2016, CONUS. Raster digital data. www.landfire.gov</p> <p>Existing Vegetation Type represents complexes of plant communities representing NatureServe's terrestrial Ecological Systems classification. Data for continental US was downloaded, reprojected, clipped to reference extent, and snapped to be compatible with other environmental inputs. Cells attributed as EVT_Name = Southern Rocky Mountain Pinyon-Juniper Woodland were used as the input feature source data for generating a Euclidian Distance raster.</p>
dist_tiptwilk	<p>Environmental Input layer: Distance to Tipton Tongue of Green River Formation (includes Wilkins Peak member) Source citations: Tweto, Ogden. 1979. Geologic map of Colorado. Vector digital data, 1:500,000. U.S. Geological Survey. ftp://greenwood.cr.usgs.gov/pub/open-file-reports/ofr-92-0507/</p> <p>Original ARC/INFO data was converted to ArcGIS shapefile and projected to UTM_NAD83_Zone13.</p> <p>Polygons with NAME attribute Tgt were selected, exported, merged, and used as the input feature source data for generating a Euclidian Distance raster.</p>
dist_td_kjdr	<p>Environmental Input layer: Distance to Dry Union Formation (Td) and part of Dakota/Purgatoire/Morrison/Ralston Creek formations (Kjdr) Source citations: Tweto, Ogden. 1979. Geologic map of Colorado. Vector digital data, 1:500,000. U.S. Geological Survey. ftp://greenwood.cr.usgs.gov/pub/open-file-reports/ofr-92-0507/</p> <p>Original ARC/INFO data was converted to ArcGIS shapefile and projected to UTM_NAD83_Zone13.</p> <p>Polygons with NAME attribute Td and Kjdr were selected, exported, merged, and used as the input feature source data for generating a Euclidian Distance raster.</p>

Raster name	Metadata entry
dist_troublesome	<p>Environmental Input layer: Distance to Troublesome Formation Source citations: Tweto, Ogden. 1979. Geologic map of Colorado. Vector digital data, 1:500,000. U.S. Geological Survey. ftp://greenwood.cr.usgs.gov/pub/open-file-reports/ofr-92-0507/</p> <p>Original ARC/INFO data was converted to ArcGIS shapefile and projected to UTM_NAD83_Zone13.</p> <p>Polygons with NAME attribute Tt were selected, exported, merged, and used as the input feature source data for generating a Euclidian Distance raster.</p>
dist_tuff2	<p>Environmental Input layer: Distance to selected Tertiary volcanic tuffs Source citation: Originator: Day, W.C., Green, G.N., Knepper, D.H., and Phillips, R.C. 2000. Spatial Geologic Data Model for the Gunnison, Grand Mesa, Uncompahgre National Forests Mineral Resource Assessment Area, Southwestern Colorado and Digital Data for the Leadville, Montrose, Durango, and the Colorado Parts of the Grand Junction, Moab, and Cortez 1° x 2° Geologic Maps. Vector digital data, 1:250,000. U.S. Geological Survey. https://pubs.usgs.gov/of/1999/ofr-99-0427/</p> <p>Original ARC/INFO data was converted to ArcGIS shapefile and projected to UTM_NAD83_Zone13. Units sharing boundaries across quads were dissolved.</p> <p>Polygons with NAME attribute Tbm, Tev, Tfg, Theb, Tq, and Tur were selected, exported, and used as the input feature source data for generating a Euclidian Distance raster.</p>
dist_tvolec_sel	<p>Environmental Input layer: Distance to selected Tertiary volcanic formations in south-central Colorado Source citation: Tweto, Ogden. 1979. Geologic map of Colorado. Vector digital data, 1:500,000. U.S. Geological Survey. ftp://greenwood.cr.usgs.gov/pub/open-file-reports/ofr-92-0507/</p> <p>Original ARC/INFO data was converted to ArcGIS shapefile and projected to UTM_NAD83_Zone13.</p> <p>Polygons with NAME attribute Taf, Tbb, Tiql, Tpl were selected, exported, and used as the input feature source data for generating a Euclidian Distance raster.</p>
dist_wasatch_all	<p>Environmental Input layer: Distance to Wasatch Formation units in northwest Colorado Source citation: Tweto, Ogden. 1979. Geologic map of Colorado. Vector digital data, 1:500,000. U.S. Geological Survey. ftp://greenwood.cr.usgs.gov/pub/open-file-reports/ofr-92-0507/</p> <p>Original ARC/INFO data was converted to ArcGIS shapefile and projected to UTM_NAD83_Zone13.</p> <p>Polygons with NAME attribute Twn, Twc, Tw, Two were selected, exported, and used as the input feature source data for generating a Euclidian Distance raster.</p>

Raster name	Metadata entry
dist_water	<p>Environmental Input layer: Distance to water Source citation: U.S. Geological Survey. 2010. High Resolution National Hydrography Dataset. File-based geodatabase, vector digital data 12,000 - 24,000. http://nhd.usgs.gov/index.html</p> <p>USGS High Resolution National Hydrography Dataset (NHD) for Colorado was queried for permanent water (polygon, line, and point). Results were converted to 30m raster and a distance raster calculated.</p> <p>NHDFlowline: ("FType" = 460 OR "FType" = 558) AND (("FCode" = 46000 OR "FCode" = 46006) OR ("GNIS_Name" IS NOT Null))</p> <p>NHDWaterbody: "FCode" = 39000 OR "FCode" = 39004 OR "FCode" = 39009 OR "FCode" = 39010 OR "FCode" = 39011 OR "FCode" = 39012 OR "FCode" = 43600 OR "FCode" = 43617 OR "FCode" = 43618 OR "FCode" = 43621</p> <p>NHDPoint: "FType" = 458</p>
dry_days_fall	<p>Environmental Input layer: Average number of days during fall (Sep-Oct-Nov) with precipitation <=5mm Source citations: Thornton, PE, MM Thornton, BW Mayer, N Wilhelmi, Y Wei, RB Cook. 2012. Daymet: Daily surface weather on a 1km grid for North America,1980-2012. Acquired online (http://daymet.ornl.gov/) on 02/20/2014 from Oak Ridge National Laboratory Distributed Active Archive Center, Oak Ridge, Tennessee, U.S.A. http://dx.doi.org/doi:10.3334/ORNLDAAC/Daymet_V2 via the USGS Geo Data Portal (http://cida.usgs.gov/gdp/).</p> <p>Daymet Daily surface weather for Colorado. Units are degrees Celsius. Daymet represents an average from 1980 - 2012, at 1 kilometer resolution. The number of days during the period covering September, October, and November where precipitation was less than or equal to 5mm was averaged. Raster was down sampled to 30m, re-projected and snapped to be compatible with other environmental input.</p>
eastness	<p>Environmental Input layer: Eastness (aspect) Source citation: Colorado Natural Heritage Program. 2011. Unpublished data using USGS 30m DEM. Raster digital data.</p> <p>The Elevation raster was used to create an Aspect raster, which was then used to create two separate rasters representing northness and eastness. northness = cos(aspect) eastness = sin(aspect)</p> <p>Values range from -1 to +1. Northness will take values close to 1 if the aspect is generally northward, close to -1 if the aspect is southward, and close to 0 if the aspect is either east or west. Eastness behaves similarly, except that values close to 1 represent east-facing slopes. For more information: http://ordination.okstate.edu/envvar.htm</p>

Raster name	Metadata entry
frostday_yr	<p>Environmental Input layer: Number of days per year with minimum temperature at or below freezing Source citation: Peter E. Thornton, National Center for Atmospheric Research. 2002. Daymet: Climatological Summaries for the Conterminous United States, 1980-1997. Frost days annual. Raster digital data, 1 km resolution. http://www.daymet.org</p> <p>Daymet annual number of frost days for Colorado. Units are days. Daymet represents an average from 1980 - 1997, at 1 kilometer resolution. Raster was down sampled to 30m, re-projected and snapped to be compatible with other environmental inputs.</p>
geol_sgmc_maj1	<p>Environmental Input layer: First-listed major lithology of geologic unit Source citation: Horton, J.D., San Juan, C.A., and Stoesser, D.B, 2017, The State Geologic Map Compilation (SGMC) geodatabase of the conterminous United States (ver. 1.1, August 2017): U.S. Geological Survey Data Series 1052, 46 p., https://doi.org/10.3133/ds1052. The State Geologic Map Compilation (SGMC) Geodatabase of the Conterminous United States (ver. 1.1, August 2017) - SGMC_Geology.</p> <p>Original downloaded data was reprojected to UTM NAD83, zone 13, and clipped to the project extent, then converted to 30 m raster using the "MAJOR1" field as grid value, snapped to be compatible with other environmental inputs. Colorado polygons are based on Tweto 1979, with additional classification fields added.</p>
jan_mintemp	<p>Environmental Input layer: January Minimum Temperature Source citation: Peter E. Thornton, National Center for Atmospheric Research. 2002. Daymet: Climatological Summaries for the Conterminous United States, 1980-1997. Monthly Minimum Temperature; April. Raster digital data, 1 km resolution. http://www.daymet.org</p> <p>Daymet Monthly Minimum Temperature in January for Colorado. Units are degrees Celsius. Daymet represents an average from 1980 - 1997, at 1 kilometer resolution. Raster was down sampled to 30m, re-projected and snapped to be compatible with other environmental inputs.</p>

Raster name	Metadata entry
LDI	<p>Environmental Input layer: Landscape Disturbance Index (LDI) Source citation: Colorado Natural Heritage Program. 2020. Landscape Disturbance Index Layer for Colorado. Raster digital data. Colorado Natural Heritage Program, Fort Collins, CO</p> <p>This represents 8 individually modeled anthropogenic impacts that were then combined into a single layer. Impacts represented are:</p> <ul style="list-style-type: none"> * Agriculture * Urban Development * Oil and Gas Development * Surface Mining * Roads and Trails * Wind turbines * Solar installations <p>Each individual layer has its own relevant weight and decay function type (see Supplemental Information). The individual impact layers are then additively combined to produce an overall disturbance layer. The weights are scaled to produce a final range where scores => 500 are High impact.</p>
marchmintemp	<p>Environmental Input layer: March minimum temperature Source citation: Peter E. Thornton, National Center for Atmospheric Research. 2002. Daymet: Climatological Summaries for the Conterminous United States, 1980-1997. Monthly Minimum Temperature; May. Raster digital data, 1 km resolution. http://www.daymet.org</p> <p>Daymet Monthly Minimum Temperature in March for Colorado. Units are degrees Celsius. Daymet represents an average from 1980 - 1997, at 1 kilometer resolution. Raster was down sampled to 30m, re-projected and snapped to be compatible with other environmental inputs.</p>
max_summertemp	<p>Environmental Input layer: Maximum summer temperature Source citations: Thornton, PE, MM Thornton, BW Mayer, N Wilhelmi, Y Wei, RB Cook. 2012. Daymet: Daily surface weather on a 1km grid for North America,1980-2012. Acquired online (http://daymet.ornl.gov/) on 02/20/2014 from Oak Ridge National Laboratory Distributed Active Archive Center, Oak Ridge, Tennessee, U.S.A. http://dx.doi.org/doi:10.3334/ORNLDAAAC/Daymet_V2 via the USGS Geo Data Portal (http://cida.usgs.gov/gdp/).</p> <p>Daymet Daily surface weather for Colorado. Units are degrees Celsius. Daymet represents an average from 1980 - 2012, at 1 kilometer resolution. The highest temperature during the period including June, July, and August for each year was averaged. Raster was down sampled to 30m, re-projected and snapped to be compatible with other environmental inputs.</p>

Raster name	Metadata entry
maymintemp (may_mintemp)	<p>Environmental Input layer: May minimum temperature Source citation: Peter E. Thornton, National Center for Atmospheric Research. 2002. Daymet: Climatological Summaries for the Conterminous United States, 1980-1997. Monthly Minimum Temperature; May. Raster digital data, 1 km resolution. http://www.daymet.org</p> <p>Daymet Monthly Minimum Temperature in May for Colorado. Units are degrees Celsius. Daymet represents an average from 1980 - 1997, at 1 kilometer resolution. Raster was down sampled to 30m, re-projected and snapped to be compatible with other environmental inputs.</p>
min_wintertemp	<p>Environmental Input layer: Minimum winter temperature Source citations: Thornton, PE, MM Thornton, BW Mayer, N Wilhelmi, Y Wei, RB Cook. 2012. Daymet: Daily surface weather on a 1km grid for North America,1980-2012. Acquired online (http://daymet.ornl.gov/) on 02/20/2014 from Oak Ridge National Laboratory Distributed Active Archive Center, Oak Ridge, Tennessee, U.S.A. http://dx.doi.org/doi:10.3334/ORNLDAAC/Daymet_V2 via the USGS Geo Data Portal (http://cida.usgs.gov/gdp/).</p> <p>Daymet Daily surface weather for Colorado. Units are degrees Celsius. Daymet represents an average from 1980 - 2012, at 1 kilometer resolution. The lowest temperature during the period including December, January, and February for each year was averaged. Raster was down sampled to 30m, re-projected and snapped to be compatible with other environmental inputs.</p>
northness	<p>Environmental Input layer: Northness (aspect) Source citation: Colorado Natural Heritage Program. 2011. Unpublished data using USGS 30m DEM. Raster digital data.</p> <p>The Elevation raster was used to create an Aspect raster, which was then used to create two separate rasters representing northness and eastness. $\text{northness} = \cos(\text{aspect})$ $\text{eastness} = \sin(\text{aspect})$</p> <p>Values range from -1 to +1. Northness will take values close to 1 if the aspect is generally northward, close to -1 if the aspect is southward, and close to 0 if the aspect is either east or west. Eastness behaves similarly, except that values close to 1 represent east-facing slopes. For more information: http://ordination.okstate.edu/envvar.htm</p>

Raster name	Metadata entry
pct_copl_pj	<p>Environmental Input layer: Percent of Colorado Plateau Pinyon-Juniper Woodland vegetation type within 150m Source citation: LANDFIRE, Earth Resources Observation and Science Center (EROS), U.S. Geological Survey. 2020. Existing Vegetation Type (EVT). LANDFIRE Remap 2016, CONUS. Raster digital data. www.landfire.gov</p> <p>Existing Vegetation Type represents complexes of plant communities representing NatureServe's terrestrial Ecological Systems classification. Data for continental US was downloaded, reprojected, clipped to reference extent, and snapped to be compatible with other environmental inputs. Cells attributed as EVT_Name = Colorado Plateau Pinyon-Juniper Woodland were used as the input feature source data for a moving window analysis (focal statistics SUM) to count the number of 30m cells within a circular distance of 150m were of this vegetation type. The sum was divided by the number of cells within the window (81) to obtain a percentage grid.</p>
pct_imb_mont_sage	<p>Environmental Input layer: Percent of Inter-Mountain Basins Montane Sagebrush Steppe vegetation type within 150m Source citation: LANDFIRE, Earth Resources Observation and Science Center (EROS), U.S. Geological Survey. 2020. Existing Vegetation Type (EVT). LANDFIRE Remap 2016, CONUS. Raster digital data. www.landfire.gov</p> <p>Existing Vegetation Type represents complexes of plant communities representing NatureServe's terrestrial Ecological Systems classification. Data for continental US was downloaded, reprojected, clipped to reference extent, and snapped to be compatible with other environmental inputs. Cells attributed as EVT_Name = Inter-Mountain Basins Montane Sagebrush Steppe were used as the input feature source data for a moving window analysis (focal statistics SUM) to count the number of 30m cells within a circular distance of 150m were of this vegetation type. The sum was divided by the number of cells within the window (81) to obtain a percentage grid.</p>
pct_IBM_sage	<p>Environmental Input layer: Percent of Inter-Mountain Basins Big Sagebrush Shrubland vegetation type within 150m Source citation: LANDFIRE, Earth Resources Observation and Science Center (EROS), U.S. Geological Survey. 2020. Existing Vegetation Type (EVT). LANDFIRE Remap 2016, CONUS. Raster digital data. www.landfire.gov</p> <p>Existing Vegetation Type represents complexes of plant communities representing NatureServe's terrestrial Ecological Systems classification. Data for continental US was downloaded, reprojected, clipped to reference extent, and snapped to be compatible with other environmental inputs. Cells attributed as EVT_Name = Inter-Mountain Basins Big Sagebrush Shrubland were used as the input feature source data for a moving window analysis (focal statistics SUM) to count the number of 30m cells within a circular distance of 150m were of this vegetation type. The sum was divided by the number of cells within the window (81) to obtain a percentage grid.</p>

Raster name	Metadata entry
pct_mixed_forest	<p>Environmental Input layer: Percent of mixed conifer forest vegetation type within 150m Source citation: LANDFIRE, Earth Resources Observation and Science Center (EROS), U.S. Geological Survey. 2020. Existing Vegetation Type (EVT). LANDFIRE Remap 2016, CONUS. Raster digital data. www.landfire.gov</p> <p>Existing Vegetation Type represents complexes of plant communities representing NatureServe's terrestrial Ecological Systems classification. Data for continental US was downloaded, reprojected, clipped to reference extent, and snapped to be compatible with other environmental inputs. Cells attributed as EVT_Name = Southern Rocky Mountain Dry-Mesic Montane Mixed Conifer Forest and Woodland, Southern Rocky Mountain Mesic Montane Mixed Conifer Forest and Woodland, or Inter-Mountain Basins Aspen-Mixed Conifer Forest and Woodland were used as the input feature source data for a moving window analysis (focal statistics SUM) to count the number of 30m cells within a circular distance of 150m were of this vegetation type. The sum was divided by the number of cells within the window (81) to obtain a percentage grid.</p>
pct_RM_aspen	<p>Environmental Input layer: Rocky Mountain Aspen Forest and Woodland vegetation type within 150m Source citation: LANDFIRE, Earth Resources Observation and Science Center (EROS), U.S. Geological Survey. 2020. Existing Vegetation Type (EVT). LANDFIRE Remap 2016, CONUS. Raster digital data. www.landfire.gov</p> <p>Existing Vegetation Type represents complexes of plant communities representing NatureServe's terrestrial Ecological Systems classification. Data for continental US was downloaded, reprojected, clipped to reference extent, and snapped to be compatible with other environmental inputs. Cells attributed as EVT_Name = Rocky Mountain Aspen Forest and Woodland were used as the input feature source data for a moving window analysis (focal statistics SUM) to count the number of 30m cells within a circular distance of 150m were of this vegetation type. The sum was divided by the number of cells within the window (81) to obtain a percentage grid.</p>
ppt_s1	<p>Environmental Input layer: Winter Precipitation Source citation: Peter E. Thornton, National Center for Atmospheric Research. 2002. Daymet: Climatological Summaries for the Conterminous United States, 1980-1997. Monthly Total Precipitation. Raster digital data, 1 km resolution. http://www.daymet.org</p> <p>Daymet total precipitation (centimeters) for December, January, & February for Colorado were totaled to represent average winter precipitation. Daymet represents an average from 1980 - 1997, at 1 kilometer resolution. Raster was down sampled to 30m, re-projected and snapped to be compatible with other environmental inputs.</p>
ppt_s2	<p>Environmental Input layer: Spring Precipitation Source citation: Peter E. Thornton, National Center for Atmospheric Research. 2002. Daymet: Climatological Summaries for the Conterminous United States, 1980-1997. Monthly Total Precipitation. Raster digital data, 1 km resolution. http://www.daymet.org</p> <p>Daymet total precipitation (centimeters) for March, April, & May for Colorado were totaled to represent average winter precipitation. Daymet represents an average from 1980 - 1997, at 1 kilometer resolution. Raster was down sampled to 30m, re-projected and snapped to be compatible with other environmental inputs.</p>

Raster name	Metadata entry
ppt_s3	<p>Environmental Input layer: Summer Precipitation Source citation: Peter E. Thornton, National Center for Atmospheric Research. 2002. Daymet: Climatological Summaries for the Conterminous United States, 1980-1997. Monthly Total Precipitation. Raster digital data, 1 km resolution. http://www.daymet.org</p> <p>Daymet total precipitation (centimeters) for June, July, & August for Colorado were totaled to represent average winter precipitation. Daymet represents an average from 1980 - 1997, at 1 kilometer resolution. Raster was down sampled to 30m, re-projected and snapped to be compatible with other environmental inputs.</p>
ppt_s4	<p>Environmental Input layer: Fall Precipitation Source citation: Peter E. Thornton, National Center for Atmospheric Research. 2002. Daymet: Climatological Summaries for the Conterminous United States, 1980-1997. Monthly Total Precipitation. Raster digital data, 1 km resolution. http://www.daymet.org</p> <p>Daymet total precipitation (centimeters) for September, October, & November for Colorado were totaled to represent average winter precipitation. Daymet represents an average from 1980 - 1997, at 1 kilometer resolution. Raster was down sampled to 30m, re-projected and snapped to be compatible with other environmental inputs.</p>
ppt_yrly	<p>Environmental Input layer: Total Annual Precipitation Source citation: Peter E. Thornton, National Center for Atmospheric Research. 2002. Daymet: Climatological Summaries for the Conterminous United States, 1980-1997. Monthly Total Precipitation. Raster digital data, 1 km resolution. http://www.daymet.org</p> <p>Daymet total precipitation (centimeters) for all months for Colorado were totaled to represent average annual precipitation. Daymet represents an average from 1980 - 1997, at 1 kilometer resolution. Raster was down sampled to 30m, re-projected and snapped to be compatible with other environmental inputs.</p>
relief	<p>Environmental Input Layer: Local Relief Source citation: Derived from U.S. Geological Survey. 2006. 30m Digital Elevation Model for Colorado. Raster digital data. A measure of surface roughness. Created from 30m DEM for Colorado by using FocalRange command: FOCALRANGE(coelev30, Circle, 16, DATA)</p>

Raster name	Metadata entry
riparian_dist	<p>Environmental Input Layer: Distance to wetland/riparian area Source citations: United States Forest Service. 2006. LANDFIRE Current Vegetation for Colorado. Raster digital data, 30m. http://landfire.cr.usgs.gov/viewer/viewer.html U.S. Geological Survey. 2010. High Resolution National Hydrography Dataset. File-based geodatabase, vector digital data 12,000 - 24,000. http://nhd.usgs.gov/index.html</p> <p>There is not a complete statewide dataset for wetland or riparian areas. Using available partial datasets (NWI, CDOW riparian) may just bias to mapped areas. Decided to try using NHD & LandFire as described below, but this is known to be an imperfect solution. USGS High Resolution National Hydrography Dataset (NHD) for Colorado and USFS LandFire Current Vegetation were queried for wetland and riparian areas. Results were converted to 30m raster, and a distance raster calculated. NHDWaterbody: "FType" = 361 OR "FType" = 466 OR "FCode" = 39001 OR "FCode" = 39005 OR "FCode" = 39006 LandFire Current Veg: "SYSTMGRPNA" LIKE '%Riparian%' OR "SYSTMGRPNA" LIKE '%Wet%</p>
shortgrass_mod2	<p>Environmental Input layer: Boosted Regression Tree model of Shortgrass Prairie in Colorado. Source citation: Fink, Michelle. 2014. Final model of Shortgrass Prairie for use in Colorado Wildlife Action Plan Enhancement: Climate Change Vulnerability Assessment. Colorado Natural Heritage Program, unpublished 30m raster digital data.</p> <p>This a distribution model produced using the Boosted Regression Tree method, with values representing an approximate probability of shortgrass prairie occurrence at each cell. Original raster was resampled and snapped to the reference extent.</p>
slope_deg	<p>Environmental Input layer: Slope (degrees) Source citation: Colorado Natural Heritage Program. 2011. Unpublished data using USGS 30m DEM. Raster digital data.</p>
snow_persistence	<p>Environmental Input layer: Snow Persistence Index Source citation: Hammond, J. C., F. A. Saavedra, S. K. Kampf (2017). MODIS MOD10A2 derived snow persistence and no data index for the western U.S., HydroShare, https://doi.org/10.4211/hs.1c62269aa802467688d25540caf2467e Raster digital data.</p> <p>Images from each year were reprojected, clipped to the reference extent, and averaged. Values are the percent of year where snow is on the ground.</p>

Raster name	Metadata entry
soil_pct_org	<p>Environmental Input layer: Average % organic matter in soil</p> <p>Source citations: Miller, D.A. and R.A. White. 1998. A Conterminous United States Multi-Layer Soil Characteristics Data Set for Regional Climate and Hydrology Modeling. Data derived from Natural Resources Conservation Service (NRCS) State Soil Geographic database (STATSGO). Tabular digital data. http://www.essc.psu.edu/soil_info/index.cgi?soil_data&conus</p> <p>NRCS. 1994. State Soil Geographic (STATSGO) data base for Colorado. United States Department of Agriculture, Natural Resources Conservation Service. Available at http://datagateway.nrcs.usda.gov</p> <p>NRCS. 2012. Gridded Soil Survey Geographic (gSSURGO) Database for Colorado. United States Department of Agriculture, Natural Resources Conservation Service. Available at http://datagateway.nrcs.usda.gov. November 20, 2012 (FY2013 official release).</p> <p>Values are supplied for each of 11 standard soil levels, down to 2.5m. Values of 0 are really NoData. Non-zero values were averaged from layers 1 - 6 as a proxy for percent clay composition down to 60cm soil depth. Due to the coarse scale of STATSGO (NRCS 1994) and the incomplete nature of SSURGO (NRCS 2012) in Colorado, all soil inputs used in CODEX PGCN models were based on the combined STATSGO-SSURGO version. Tabular data was joined to the combined STATSGO-SSURGO vector digital dataset for Colorado and exported as a 30m raster.</p>
soil_ph	<p>Environmental Input layer: Soil pH</p> <p>Source citations: Miller, D.A. and R.A. White. 1998. A Conterminous United States Multi-Layer Soil Characteristics Data Set for Regional Climate and Hydrology Modeling. Data derived from Natural Resources Conservation Service (NRCS) State Soil Geographic database (STATSGO). Tabular digital data. http://www.essc.psu.edu/soil_info/index.cgi?soil_data&conus</p> <p>NRCS. 1994. State Soil Geographic (STATSGO) data base for Colorado. United States Department of Agriculture, Natural Resources Conservation Service. Available at http://datagateway.nrcs.usda.gov</p> <p>NRCS. 2012. Gridded Soil Survey Geographic (gSSURGO) Database for Colorado. United States Department of Agriculture, Natural Resources Conservation Service. Available at http://datagateway.nrcs.usda.gov. November 20, 2012 (FY2013 official release).</p> <p>Soil pH values are supplied for each of 11 standard soil levels, down to 2.5m. Values of 0 are really NoData. Non-zero pH values were averaged from layers 1 - 6 for this project. Note - a mathematical mean is not technically the appropriate way to lump multiple pH values, but we are restricted by how the data were originally recorded. Surface pH alone was not seen as sufficient information, so we averaged the values of the first 6 layers as a proxy for actual total pH down to 60cm soil depth. Due to the coarse scale of STATSGO (NRCS 1994) and the incomplete nature of SSURGO (NRCS 2012) in Colorado, all soil inputs used in CODEX PGCN models were based on the combined STATSGO-SSURGO version. Tabular data was joined to the combined STATSGO-SSURGO vector digital dataset for Colorado and exported as a 30m raster.</p>

Raster name	Metadata entry
soil_type	<p>Environmental Input layer: Soil type (Map Unit) Source citations: NRCS. 2012. Gridded Soil Survey Geographic (gSSURGO) Database for Colorado. United States Department of Agriculture, Natural Resources Conservation Service. Available at http://datagateway.nrcs.usda.gov. November 20, 2012 (FY2013 official release).</p> <p>In deductive models, map units were selected as appropriate, merged, and converted to 30m raster as required.</p>
ssurgo_depth_cm	<p>Environmental Input layer: Soil depth Source citations: Miller, D.A. and R.A. White. 1998. A Conterminous United States Multi-Layer Soil Characteristics Data Set for Regional Climate and Hydrology Modeling. Data derived from Natural Resources Conservation Service (NRCS) State Soil Geographic database (STATSGO). Tabular digital data. http://www.essc.psu.edu/soil_info/index.cgi?soil_data&conus NRCS. 1994. State Soil Geographic (STATSGO) data base for Colorado. United States Department of Agriculture, Natural Resources Conservation Service. Available at http://datagateway.nrcs.usda.gov NRCS. 2012. Gridded Soil Survey Geographic (gSSURGO) Database for Colorado. United States Department of Agriculture, Natural Resources Conservation Service. Available at http://datagateway.nrcs.usda.gov. November 20, 2012 (FY2013 official release).</p> <p>Depth to bedrock (field ROCKDEPM) is a single value per soil polygon. Units are centimeters. Note that a value of 152 really means >= 152 cm and a value of 0 is really NoData (occurs on Water polygons only). Due to the coarse scale of STATSGO (NRCS 1994) and the incomplete nature of SSURGO (NRCS 2012) in Colorado, all soil inputs used in CODEX PGCN models were based on the combined STATSGO-SSURGO version. Tabular data was joined to the combined STATSGO-SSURGO vector digital dataset for Colorado and exported as a 30m raster.</p>
ter_rough_index	<p>Environmental Input layer: Terrain Ruggedness Index Source citation: Colorado Natural Heritage Program. 2021. Unpublished data using USGS 30m DEM. Raster digital data.</p> <p>The Elevation raster was used to create an index of terrain ruggedness reflecting the difference in elevation between neighboring cells. R script provided by Michelle Fink, CNHP.</p>

Appendix B: Basic modeling methods

Element Occurrence or location processing

- For each species use a single-species shapefile output from BIOTICS (or other source)
- Use the Multipart to Singlepart tool to separate all polygons (this interim step not kept)
- Use the Feature to Point tool to convert the polygons to centroid points (check the "Inside" box), and name the shapefile something like: sppname_pts.shp
- Use the Add XY Coordinates to generate location data for each point (don't use X Y fields already there from the polygon shapefile, since they belong to a single EO).
- Open the sppname_pts.dbf file in Excel
- Check for low-precision/very old EO records – these may need to be left out of the modeling dataset
- Copy SNAME and POINT_X, POINT_Y info into a three-column multispecies .csv file, omitting old, low-precision points, although you might keep these if consistent with range of other points. Generally anything older than 1970 could be left out. The csv spreadsheet should look something like the example below, with as many rows for each species as there are good locations. Species names can be as they would normally appear, including ssp. or var. Make sure your XY coordinates are all in the same spatial reference as your environmental grids (e.g., NAD83 Zone 13)

SNAME	X	Y
Species1 Name	277935.1488	4184812.83
Species1 Name	282537.3615	4191508.725
Species2 Name	228874.062	4168829.166

Environmental data processing

- All grids must share a common projection, extent, cell size, and alignment, and be in the same folder
- I typically use the 30m elevation grid as my reference grid, but just be sure to always use the same reference grid, and make sure it covers the full extent of the study area.
- The Maxent software requires ASCII grid files, but other modeling methods can use ESRI grids or geotiff (geotiff preferred, since it is smaller and easier to work with)
- Use the settings under Environments (either under Geoprocessing menu or use the button at the bottom of the raster to ASCII conversion tool) to set processing extent, snap to grid, output cell size to the reference grid. Be sure environments are correct each time an input processing step is run.
- For categorical variables such as soil type, geology type, vegetation type, it is much better to use a "distance to a particular type" instead of the categories themselves, so try to narrow down one or a few types. Use the Euclidian Distance tool with a shapefile as the input feature to generate the distance to X grid.

Maxent modeling

- Unzip the Maxent files in location on your computer (not on the network, unless using modeling server)
- Make sure you have java installed

- To open Maxent, double click the maxent.bat file
- Browse to the location of samples .csv file, and folder of environmental layers files (ascii rasters)
- Select one species in the left-hand window and select appropriate environmental layers in the right window. Be sure to specify if a layer is categorical instead of continuous (default).
- If you always use the same location for environmental layers, model runs will be faster after the first time because Maxent makes a cache of layer info. Every time a new layer is used, it will have to write that into the cache.
- Check boxes on right side (create response curves, make pictures..., do jackknife), OK to use auto features - left side box. If there are plenty of points (like more than 100 or so, you could, under settings at the bottom, put in a number like 10 to 25 in the Random test percentage box. This gives a better estimate of model fit. For these rare species models, it isn't really necessary.
- **Use the default asc output, the other ones don't work in ArcMap**
- Note that all climate variables (precipitation and temperature) are highly correlated with elevation, so I sometimes omit the elevation grid
- Specify the output directory, leave projection layers field blank (this is for projecting under future conditions, e.g., climate change)
- See this pdf for help with Maxent settings P:\C NAP_2021_2025\CODEX PGCN Models\references\modeling\a_maxent_model_v7.pdf
- When ready, click the run button in lower left. Program will let you know if there is a problem with any inputs. One or two layers i.e., soil_pH have "no data" in some areas such as the reservoirs, so OK to just say ignore and suppress additional warnings
- A run will probably take an hour or two, depending on if data is all cached.
- If you want to do another run, make a new Maxent folder (e.g., maxent2) so as not to overwrite the previous run.

Model results

- It is best to set ArcMap to not automatically turn on added layers (Customize > ArcMap Options > General tab, uncheck box "Make newly added layers visible by default")
- Add the .asc file that is in your Maxent folder for that species to an ArcMap mxd
- Convert the .asc to a .tif raster using Conversion tools > To Raster > ASCII to Raster. Specify .tif for the output raster, and use FLOAT for output data type
- Classify the tif raster under symbology tab - yes to calculate statistics first
- Potentially keep everything above 0.50; botanists to review, determine final extent
- Initial cutoffs used were orange= Equal training sensitivity and specificity, red= 0.50+

Post review processing

- To "erase" an area (i.e., reservoir), make a polygon of the feature to be erased, and convert it to a raster, using the complete model raster as processing extent, snap raster, and raster analysis cell size in environment settings. In raster calculator, use a statement like this to set reservoir cells to "NoData": SetNull(~(IsNull("EchoCynRes_PolygonToRaster")), "Townsendia_glabella.tif"). This is now the new full model. Now reclassify according to desired cutoff and export for CODEX model, use additional range clip if needed.
- For CODEX binary version, classify the full model into two display classes, then reclassify this so that everything not kept (e.g., cells <0.50, and areas of NoData) becomes NoData, and cells above cutoff = 1. Then clip this binary if needed.
- For clipping with non-rectangular shape, check the "Use input features for clipping geometry" box

- To make a shape incorporating an elevation contour, use Contour (spatial analyst tools / surface) on a classified elevation grid, then draw a boundary polygon and use this in feature to polygon to get the contour portion. Then select appropriate part and export as clip shape.

Deductive models

- Maxent can be used to investigate the contribution of selected variables to predictive ability in a model, but if results are unsatisfactory, a deductive model may be needed.
- Deductive models are constructed by combining grids of the various factors, using the "envelope" of applicable conditions (e.g., elevation between 5000 and 9000 ft).
- To identify the envelope or range for each factor, use the Extract Values to Points tool to intersect species location points with each environmental raster input. Values will be output in a new column in the new output shapefile
- When you have the values for each factor, use the Raster Calculator with CON statements, or the Reclassify tool (after setting the display classes) to pull out the ranges (values outside the range of interest should become NoData, values in range 1). Add the binary rasters together in Raster Calculator, then Reclassify again. For a binary result, use the highest value as 1, everything else becomes NoData. Or you may want to keep areas where all but one factor agrees (next highest value) as moderate probability, etc.

Appendix C: Expert review of SDM for rare plant species

Introduction

In the past two decades, the popularity and accessibility of species distribution modeling has increased dramatically (Franklin 2013, Guisan et al. 2013, Zurell et al. 2020). Accessible computing equipment, a proliferation of predictive algorithms, and user-friendly interfaces have made it possible for nearly anyone with access to spatial data to produce a predictive distribution model for a species of interest, regardless of their knowledge of the species and its habitat. This is in stark contrast to previous times when species distribution maps were produced by one or more specialists, using extensive field knowledge of “their” species, a paper map, and an indelible marker. These early “models” were easily converted to digital format, and often remain the most useful representation of where a particular species can be expected to occur.

The growth of species distribution modeling has resulted in countless digital models, covering various areas, but often only occasionally tested in the field. A primary validation of any distribution model is that it permits the discovery of previously unknown occurrences. Standard statistics used to assess the quality of a model (for example, the widely reported Area under the Receiver Operating Curve or AUC), do not necessarily tell us much about whether the model is good in the field. Field validation is expensive, time consuming, and can even be dangerous when predicted habitat is in difficult terrain. A somewhat more accessible evaluation technique is the expert review performed by those who have observed the modeled species in the field.

In the course of a project to produce models for use in conservation planning, we solicited expert review from regional botanists/ecologists in a uniform format. Models were produced for 80 of the Plants of Greatest Conservation Need as specified in the 2015 Colorado State Wildlife Action Plan. These distribution models will be included in the conservation data sharing platform, the Colorado Conservation Data Explorer (CODEX), to help conserve and protect these species through environmental review and conservation planning. As such, the modeling process was focused on producing products that kept a balance between predicting too much suitable habitat to be reasonable in field survey, and models that too closely fit already documented locations.

Our objectives are to report on the results of the review process, identify sources of reviewer dissatisfaction with models, recognize corrective actions that can be taken to address reviewer comments, and recommend best practices for future reviews.

Methods

Reviews were solicited from botanists active currently or in the past in Colorado and adjacent states, on the basis of having some familiarity with a particular species. Reviewers were affiliated with a variety of local, state, and federal agencies, universities, regional herbaria, and/or the Colorado Native Plant Society. An online survey google form with a combination of radio button choices and fields for text comment was used in combination with a binary model draft displayed on ArcGIS Online, both viewable in an internet browser (supplementary material). CNHP botanists made an effort to remind reviewers to fill out the survey, and to clarify unclear survey responses. In a few instances, reviewers chose to respond by email, and their answers were entered into appropriate columns to the extent that could be

determined from the email content. Not all reviewers completed all questions. The questions asked in the google form are shown in Table C-1.

Table C-1. Questions included in the Google Form for expert model review, Round I. Questions with an * were included in the google form for Round II review.

Question	Response Type	Choices
What is your name?	Short answer	
What is your email address?	Short answer	
Which species did you review?	Short answer	
Briefly describe your familiarity with the species and areas where you have observed it.	Long answer	
Overall correctness	Multiple Choice	Un-usable, Poor, Reasonable, Good, Excellent
Please add any comments to justify the overall score which you chose.	Long answer	
Fit	Multiple Choice	Underfit (model covers too broad of an area), Overfit (model covers too narrow of an area), Suitable
Please add any comments to justify the fit which you chose.	Long answer	
Distribution	Multiple Choice	Should include additional areas, Should exclude certain areas, Suitable
Please add any comments to justify the distribution which you chose.	Long answer	
Do you have a shapefile or circled map image of the area you believe should be included or excluded? If so, please email to jp.smith@colostate.edu	Multiple Choice	Yes, No
Additional Comments	Long answer	
*May we identify you in the final report as an expert reviewer of this species?	Multiple Choice	Yes, No

Responses were tabulated in a spreadsheet that preserved all online entries. Categorical responses were tallied, and comments were summarized into five general categories of dissatisfaction with the model.

Comment summary category definitions

Too broad / widespread: The model area was perceived as including too much area outside the current documented range of the species

Has incorrect inclusions: The model area included unsuitable habitat types to a degree that prompted comment (e.g., riparian areas included for an upland species, reservoirs as habitat)

Missing known occurrences: The area within modeled suitable habitat did not cover one or more documented locations for the species

Data refinement desired: The reviewer suggested an environmental factor that appeared to be omitted from the model, or a more fine-scaled representation of included data

Reviewer unclear on model purpose: The reviewer expected the model to show detailed microhabitat at a scale much finer than that of the model resolution, or expected the modeled area to conform closely to only known occurrence locations

Results

Thirty-nine individual botanists reviewed a total of 46 models created in Round I of this project, for a total of 75 reviews. Individuals reviewed from one to eight species models, depending on the extent of their expertise. As expected, model review detail and useful feedback was variable both within and between reviewers. Over 80% of reviews concluded that the model in question was in the Good or higher level of Overall Correctness. Ten models were assigned Poor, and two were labeled Un-useable. The majority of rankings fell within the three middle Overall Correctness categories from Poor to Good (Table C-2). As reviews were submitted, we realized that it would have been advisable to better define evaluation parameters, and to adequately describe the purpose and use of the models. Reviewer comments indicated that model characteristics making a model un-useable to one reviewer might be acceptable to a different reviewer. In some instances, the disinclination of a reviewer to make use of the standard review form (replying instead by email) detracted from the utility of their response.

The overall goal of the review process was for each reviewer to compare a model of predicted suitable habitat against their personal knowledge of a species and its habitat. Although reviewers had all observed their reviewed species, there was considerable variation in the level of experience needed to make useful comparisons between an external GIS model and the internal “model that I carry in my head with regard to this species.” There was naturally variation in amount of reviewer experience in locating and identifying the species in the field; some reviewers discovered and described the species, others had seen it only a single time, but most reviewers had at least moderate experience with the species. Reviewer experiences were highly variable in how recently the species or its habitat had been observed. Some reviewers had surveyed during the most recent field season, others had not seen the species for one or more decades. Although their most recent observation date was not usually reported by reviewers as part of this survey, it appeared from responses that agency personnel and those with a particular investment (i.e., discoverers/publishers) are those most likely to be making frequent and recent surveys. Finally, individual biases and beliefs about what a model “should” be were apparent in many responses. This information is difficult to quantify, but should be useful in guiding future review efforts, and illuminating potential areas where communication can be improved.

Categorized reviewer comments revealed that even models ranked Good or Excellent might have room for improvement. In general, reviewers were most likely to indicate that a model was too broad or included too much unsuitable habitat (Table C-2). The inclusion of unsuitable habitat in predicted areas was the second most frequent comment type.

Table C-2. Tally of reviewer comments by comment summary category and overall model correctness score.

	Excellent	Good	Reasonable	Poor	Un-usable	% of total reviews
Too broad / widespread		3	11	5	2	31%
Has incorrect inclusions	1	5	6	4	1	25%
Missing known occurrences		2	4	1	1	12%
Data refinement desired	1	3	6	3		19%
Reviewer unclear on model purpose			1	5	2	12%

In spite of the variable responses from reviewers, the exercise did produce important feedback that was used to revise some models. Three models were revised back on feedback; these were for *Aletes humilis*, *Draba smithii*, and *Pediocactus knowltonii*. The most easily implemented revisions were removing portions of predicted habitat too far outside the known range when a botanist indicated that they had surveyed unsuccessfully for the species in that area (e.g., *Draba smithii*), or removing obviously incorrect habitat areas (e.g., reservoirs) from modeled habitat (e.g., *Aletes humilis*). Revisions that would be too complex to implement under the scope of the project, or feedback suggesting other data types were noted and those species models flagged for revision priority in future work. All reviewer input, even if no solutions were suggested, will be considered in future revisions.

Discussion

The five Overall Correctness categories were presented without definitions in the first round of review. For some reviewers, a misunderstanding about what would make a model poor or unusable required a follow-up between CNHP botanists and the reviewer to clarify. This experience shows that those soliciting model review may need to educate botanical experts on how/why models are produced, and work to clarify review expectations. For instance, the ranking categories should be well defined (e.g., Table C-3 below), and/or perhaps condensed into the three middle categories of Good, Reasonable, and Poor.

Table C-3. Suggested definitions for use in model review.

Overall Correctness Category	Definition
Five-category method	
Excellent	Model covers occurrences, appears to narrow down suitable habitat area well, given coarseness of available data, indicates convincing areas for survey.
Good	Model covers occurrences, contains some areas of unsuitable/unlikely habitat, but these do not dominate and could be omitted, indicates plausible areas for survey.
Reasonable	Model covers most good occurrences, may include significant unsuitable area, but could be trimmed, indicates some likely areas for survey.
Poor	Model misses some important occurrences, includes large unsuitable areas, generally does not represent species habitat, but may indicate additional likely areas for survey.

Overall Correctness Category	Definition
Un-useable	Model misses many occurrences, is no better than random survey, does not agree with reviewer beliefs about the species, indicates additional areas that are thought to be highly improbable.
Potential three-category method	
Good	Model covers occurrences, narrows to suitable habitat in a generally satisfactory way, may contain some areas of unsuitable/unlikely habitat, but these do not dominate and could be omitted, indicates plausible areas for survey.
Reasonable	Model covers most good occurrences, may include significant unsuitable area, but could be trimmed, indicates some likely areas for survey.
Poor	Model misses many occurrences, includes large unsuitable areas, generally does not represent species habitat well, may include highly improbable area, but may also indicate additional likely areas for survey

In the Round II of the project, the model review form included the following project explanation at the beginning: These models were created for CODEX (Colorado's Conservation Data Explorer), a free web-based mapping and environmental review tool hosted by CNHP (<https://codex.cnhp.colostate.edu/>). The intent of these models is to facilitate conservation and protection of these species through environmental review and conservation planning. Models are a broad, inclusive representation of locations recommended for survey where suitable habitat may exist. Due to modeling limitations, some unsuitable habitat may be included. The definitions in table C-3 were also included on the form with the question for overall model correctness.

Through more detailed evaluation of reviewer comments, we were able to identify and assign sources of model dissatisfaction to several entities, namely, the review process, the modeling process, or the data (Table C-4). We also identified potential corrective actions that can be implemented to improve the review process. Although some sources of model concern arising from data cannot be corrected, improved communication with regard to model process and objective together with the realities of geo-spatial data, are likely to improve the utility of the review processes for all involved.

Table C-4. Source of model dissatisfaction

Concern	Potential corrective actions
Review process	
Reviewer expects model to show microhabitat	More detailed explanation of purpose and intended use of model
Reviewer expects model to be tightly constrained to vicinity of known locations (i.e., to “show the range”)	More detailed explanation of purpose and intended use of model; work to implement a cost-effective method of either selecting small portions of statewide environmental datasets, or else constraining background samples to be nearer known occurrences. The goal of this effort would be to eliminate distant and dissimilar areas from model construction.

Concern	Potential corrective actions
Reviewer expects wants finer scale model	Detailed explanation of realities of data availability; or work with modeler to develop better inputs
Modeling process	
Lack of experience with modeling technique and/or data preparation	Work with, hire/retain experienced modelers; train additional staff
Insufficient bio-eco-geo knowledge to permit interpretation of inputs and draft results	Modeler works closely with botanist, ecologist, etc. to determine appropriate inputs
Data - Occurrences	
Missing occurrences	Locate and add additional occurrences; request data from reviewer; consult with field botanists who may have unsubmitted records
Occurrences poorly mapped	Improve mapping if possible; remove or down-weight poorly mapped occurrences
Too few occurrences	Deductive model based on specimen information, re-do model if additional occurrences found
Some clusters of occurrences in well surveyed areas dominate model results	Subset and/or weight occurrences
Data - Environmental covariates	
Desired input not available	Explain realities of data availability, try to identify surrogate
Desired input not available across species range	Detailed explanation of realities of data availability, try to identify surrogate data, or constrict model area
Scale too coarse or shows too widespread predicted distribution	Constrict model area, down sample data
Predicted area includes too much unsuitable habitat	Find data that better represents key factors, down sample data, cut unsuitable areas from final model
Many highly correlated inputs	Correlation analysis considered in selection of inputs
Errors or unexpected patterns in data layer produce unlikely results	Check validity of original data and preparation process; omit this covariate
Distribution not driven by any known covariate	More research on species requirements needed

Tradeoffs between model production costs and validation efforts are a crucial consideration in the often under-funded sphere of rare species conservation. Most species distribution models have a statistical evaluation of some sort, ideally with a separate, independent set of test points that have not been used to create (train) the model. In our work modeling the predicted distribution of rare plant species, we

typically do not have enough location points to justify withholding a useful percentage from model construction. While it is clearly ineffective to hold out one or two of ten points for testing, there are no recognized standards for when and how many points to hold out as a test percentage. With ample time and funding, this question could be researched, but support is generally wanted for more essential tasks such as field survey and conservation efforts. Consequently, we take comparative note of statistics such as AUC that compare the performance of the model against the combined set of training and background points, but do not regard them as high-quality model evaluation tools in this situation.

Field validation can be statistically rigorous, with spatially balanced randomly generated sample points enabling statistical evaluation. This type of exercise is often prohibitively expensive and time consuming, unless the species is a well-funded conservation target. Less rigorous, informal field validation is more realistic, and should be employed whenever possible. In this scenario field biologists have access to the model (either in the field or in the office after collecting accurate location data) and can report success or failure of model at a visited location. Success and failure can take several meanings here. A clear validation is if the target species is in fact found at a predicted location. However, the species absence at a location that otherwise appears highly suitable is not always a failure. Possible reasons for the absence include:

- the species is mobile or only seasonally apparent and was missed during the field visit
- the species could become established in the location, but has not dispersed to the area (for various reasons), or was formerly present but extirpated by anthropogenic activity
- the habitat is in fact unsuitable, and could not support the species if it has ever dispersed to the area

Our goal is to test whether the model is useful for our objective, not whether it is correct, however that is defined (Pearson 2010). Expert review of species distribution models can act as an informal post hoc field validation, similar to the in-office post field-visit scenario, but with more variable results, depending on the experience and model evaluation skill of the expert. Our experience with expert review indicates that our rare plant models are acceptable tools and representative of on-the-ground conditions. In addition, the cost-benefit balance, together with the network-building effect and iterative nature of the model production-review-revision process can lead to a synergistic effect that benefits both the species of greatest conservation need and those who study them.

A quick search turned up several hundred to a few thousand publications on the topics of boundary organizations and knowledge co-production in conservation, land management, and climate change response. A special issue of *Frontiers in Ecology and Evolution* (Enquist et al. 2017) featured a series of articles about the emerging field of translational ecology (TE) wherein boundary-spanning organizations are depicted as key players in the translational process between scientists (especially academia) and stakeholders (land management agencies, etc.). Clearly there is widespread feeling that researchers and practitioners could benefit from stronger ties. This has led to a proliferation of discussion about boundary spanners at the science/policy divide, boundary objects (physical objects or tools providing a common point of reference among stakeholders), evidence bridges (facilitating knowledge exchange between researchers and practitioners), and related concepts (Morissette et al. 2017, Gustafsson and Lidskog 2018, Salafsky et al. 2019, Kadykalo et al. 2021).

There is, however, little recognition of the value of the vertical communication and long-term network building that greatly reduce boundary effects. TE is a delightful concept, but the authors seem unaware of a number of networked organizations that have traditionally (albeit quietly) filled this role in conservation and ecological decision making, particularly in the US. The members of the NatureServe network (state Natural Heritage Programs, Conservation Data Centers, etc.) have a decades-long history of acting as evidence bridge organizations between university or agency scientists and a variety of stakeholders, including local, state, tribal, and federal land managers, non-governmental environmental advocacy, conservation organizations, political entities, local educators, “amateur” field observers, and more.

Furthermore, Natural Heritage Program (NHP) staff typically embody a pool of both extensive field-based ecological knowledge as well as trusted long-term stakeholder relationships in their state or region. Program staff often possess biological knowledge and field experience that spans decades, providing the ability to discern trends in the biodiversity of an area. Staff may serve as recovery team or technical committee members for a particular taxa or group of taxa, as well as being members and leaders of local or regional stakeholder groups. NHPs serve as training grounds for qualified ecologists (botanists, zoologists, etc.) who may later take agency positions, and often provide training for agency personnel and other stakeholders. NHPs are well positioned to get all parties to the table, being connected with researchers and practitioners, as well as other boundary-spanning groups. NHPs preserve the tradition of natural history but have also adapted to current conservation trends and issues. The fact that we were able to elicit expert reviews from so many individual botanists is a remarkable demonstration of the value of our program’s partnership-building expertise.

In the case of rare species, especially those of less glamorous taxa, it can be a challenge to find experts who are able to evaluate a distribution model. This loss of expert knowledge is ongoing and difficult to mitigate against in a quickly changing world but provides an important source of useful (although often denigrated) evidence in conservation. The same holds true for indigenous community expertise as well.

There are, however, opportunities for all involved in the practice or funding of botany and plant ecology to facilitate the preservation and ongoing maintenance of expert knowledge. Staff of agencies and conservation organizations can work to instill the enthusiasm for field work and support upcoming botanists and ecologists on the track to subject-matter expertise while experts in senior or career positions can mentor younger colleagues to pass on both their field knowledge and other observations about a species. Academic modelers can and should move beyond highly technical explanations and learn to communicate the basic modelling process and concepts in plain English, following the advice of E.B. White that “No one can write decently who is distrustful of the reader's intelligence, or whose attitude is patronizing.” In the end, any single species distribution model cannot be all things to all users but can play a role in facilitating and sustaining networks of conservation practice.

References

Enquist, C.A.F, Jackson, S.T., Garfin, G.M et al. 2017. Foundations of translational ecology. *Frontiers in Ecology and the Environment* 15:541-550 doi: 10.1002/fee.1733

Franklin, J. 2013. Species distribution models in conservation biogeography: developments and challenges. *Diversity and Distributions* 19: 1217–1223.

Guisan, A., Tingley, R., Baumgartner, J.B. et al. 2013. Predicting species distributions for conservation decisions. *Ecology Letters* 16:1424–1435.

Gustafsson, K.M. and R. Lidskog. 2018. Boundary organizations and environmental governance: Performance, institutional design, and conceptual development. *Climate Risk Management* 19:1-11.

Kadykalo, A.N., R.T. Buxton, P. Morrison, C.M. Anderson, H. Bickerton, C.M. Francis, A.C. Smith, L. Fahrig. 2021. Bridging research and practice in conservation. *Conservation Biology* 35:1725–1737.

Morisette, J.T., A.E. Cravens, B.W. Miller, M. Talbert, C. Talbert, C. Jarnevich, M. Fink, K. Decker, E.A. Odell 2017. Crossing Boundaries in a Collaborative Modeling Workspace, *Society & Natural Resources*, 30:9, 1158-1167, DOI: 10.1080/08941920.2017.1290178

Pearson, R.G. 2010. Species' Distribution Modeling for Conservation Educators and Practitioners. *Network of Conservation Educators and Practitioners, Center for Biodiversity and Conservation, American Museum of Natural History Lessons in Conservation, Vol. 3, pp. 54-89*
<https://www.amnh.org/content/download/141368/2285424/file/ncep.amnh.org/linc/>

Salafsky, N., Boshoven, J., Burivalova, Z., Dubois, N. S., Gomez, A., Johnson, A., Lee, A., Margoluis, R., Morrison, J., Muir, M., Pratt, S. C., Pullin, A. S., Salzer, D., Stewart, A., Sutherland, W. J., & Wordley, C. F. R. 2019. Defining and using evidence in conservation practice. *Conservation Science and Practice*, 1, e27.
<https://doi.org/10.1111/csp2.27>

Zurell, D., Franklin, J., König, C., Bouchet, P. J., Dormann, C. F., Elith, J., Gusman, G. F., Feng, X., Guillera-Arroita, G., Guisan, A., Lahoz-Monfort, J. J., Leitão, P. J., Park, D. S., Peterson, T., Rapacciuolo, G., Schmatz, D. R., Schröder, B., Serra-Diaz, J. M., Thuiller, W., Yates, K. L., Zimmermann, N. E. and Merow, C. 2020. A standard protocol for reporting species distribution models. – *Ecography* doi: 10.1111/ecog.04960

Appendix D: Projected suitable habitat under potential future climate conditions

Climate model basics

General circulation models (global climate models) or GCM are computer models that simulate how various physical processes interact in the atmosphere, oceans, and landmasses to produce world-wide climate patterns. GCMs are used for all types of investigation into climate behavior, both short and long term. These models are tested to see how well they predict past conditions. Global scale models use a three-dimensional grid of large cells (on the order of 1 x 2 degrees – about 16 of which cover Colorado).

Dozens of modelling groups (centers) around the world use GCMs under various scenarios to predict what climate conditions might be like in the future. Scenarios represent the complex relationship between the socioeconomic forces driving greenhouse gas and aerosol emissions and the levels to which those emissions would climb during the 21st century. In more recent model efforts, scenarios are called Representative Concentration Pathways (RCP). Models are set up with known historic conditions in 1950, and run without additional correction to 2100, using input specifications (scenario or RCP values) about how greenhouse gases will change under global circulation patterns.

Each model run produces complex multidimensional output (a global 3-dimensional climate grid over time). The data we use is typically available in NetCDF format. Extensive data manipulation is required to convert the NetCDF output into the various monthly, seasonal, and annual rasters of precipitation or temperature that we use in species distribution models. Following standard practice in weather data, climate data is typically averaged over a 30-year “normal” period for comparison with new observations or future projections.

Because all models have their particular biases, it is important to predict future species distribution by using two separate “slices” of the same 150-year model dataset (model space), one representing the recent past (historic normal), and the other representing the projected future normal for a particular period. This controls for model bias and allows us to have confidence that the observed change is due to scenario conditions, not a difference resulting from using two models.

Test model of predicted future distribution for *Draba smithii*

From prior climate change related project work, CNHP had available a selection of processed seasonal precipitation and temperature statewide datasets. Unfortunately, a complete set of rasters equivalent to climate inputs used in the PGCN distribution modeling was not available. We were able to match seasonal precipitation, and to substitute average high for summer and fall for maximum summer temperature, and average low temperature for winter and spring to approximate winter minimum temperature. Frost date and other climate datasets were not available without significant additional computations. This climate data was based on the hadgem2.es.1.rcp85 climate model for the period 1980-2012 to represent current climate normal.

This is a model that shows a hot and dry future climate for most of Colorado, with an average annual reduction in precipitation of 4% and increase average temperature of nearly 7°F (Figure 1.). Projected future normal values were for the 30-year period centered around 2050 (i.e., 2035-2065 from the complete 150-year model dataset).




Figure 1. Graphs of the % change in precipitation and temperature change from baseline conditions between 1971-2000 in the “hot and dry” scenario predicted in the hadgem2.es.1.rcp85 climate model for the 30-year period centered around 2050.

Table 1. Comparisons of the relative contributions of model inputs for the two “current” distribution models. The model on the left used Daymet data. Daymet data is interpolated from historic weather observations (weather station locations) and represents an average from 1980 - 1997, at 1 kilometer resolution; the raster was down sampled to 30m. The model on the right used the hadgem2.es.1.rcp85 climate model, taking data from 1980-2012 to represent current climate normal.

Daymet current model

Variable	Percent contribution	Permutation importance
dist_tvolc_sel	23.1	8.5
ter_rough_index	19.1	0.6
ppt_s1	14.6	5.7
ppt_s3	12.7	40.8
max_summertemp	8.4	1.2
co_ned30m	5.4	0.1
ppt_s4	3.3	24.8
slope_deg	2.8	0
dist_ignmet_xy	2.6	1.6
northness	1.9	0.3
may_mintemp	1.4	2.7
ppt_s2	1.4	11.4
eastness	1.1	0.3

hadgem2.es.1.rcp85 current model

Variable	Percent contribution	Permutation importance
dist_tvolc_sel	24.1	39.6
ter_rough_index	22.1	0.5
ppt3_hd	13.8	7.8
co_ned30m	11.2	0
ppt1_hd	8.2	10.4
ppt4_hd	4.7	24.4
dist_ignmet_xy	3.6	6
eastness	2.7	0.8
northness	2.3	0.4
tmax4_hd	1.9	3.2
ssurgo_depth_cm	1.4	1
slope_deg	1.4	0
tmin1_hd	1.1	0.3

Variable	Percent contribution	Permutation importance	Variable	Percent contribution	Permutation importance
avg_lastfrost	0.9	0.1	ppt2_hd	1	2.5
average_clay	0.8	1.2	average_clay	0.3	0.6
min_wintertemp	0.1	0.3	average_silt	0.2	2.3
average_sand	0.1	0.2	tmax3_hd	0.1	0.1
ssurgo_depth_cm	0.1	0.1	average_sand	0	0
apr_mintemp	0	0	tmin2_hd	0	0
average_silt	0	0			
avg_firstfrost	0	0			

The figure below (Figure 2) illustrates the predicted model habitat produced during this project using Daymet climate data (top row) and predicted habitat using the hot and dry model, the hadgem2.es.1.rcp85 climate model (bottom row).

The top left shows the full Maxent model from 0-1. Climate data used in this model included seasonal precipitation, April and May mintemp, avg first and last frost, max summertemp and min wintertemp. This climate data was based on Daymet monthly data, which is interpolated from historic weather observations (weather station locations). The data is based on the average values over the period 1980-2012. For reference, the top right is the clipped binary model of high probability habitat (cut off was 0.5) which will be used in CODEX.

The bottom left is the full Maxent model for the current (recent past) distribution but using the “model space” period of the hot and dry model, the hadgem2.es.1.rcp85 climate model. Overall statewide patterns of predicted suitable habitat are similar between the two “current” full Maxent models. Differences between the top and bottom left maps are due primarily to the different temperature datasets used to represent climate. Non-climate inputs were the same as in the top left full model.


Bottom right shows the full Maxent model using the hadgem2.es.1.rcp85 climate model for the 30-year period centered around 2050 (i.e., 2035-2065 from the complete 150-year model dataset). Non-climate inputs were unchanged. Although predicted higher probability habitat remains in a few key locations (vicinity of Pikes Peak, Sangre de Cristos and Wet Mountains, Raton pass), future predicted habitat is severely diminished under hot and dry future conditions.

Conclusion


Most distribution models for rare species, including those produced for this project, are based on recent historic climate conditions, and focused on identifying survey areas, in hopes of increasing our knowledge of the species range and population levels. Species distribution models intended to facilitate the development of adaptation strategies in response to rapidly changing environmental conditions must instead focus on changes in patterns of precipitation and temperature within a species range. The delineation of a species’ climate envelope and evaluation of current population stressors, together with predicted degree of change in the near future, can provide direction for a variety of adaptive management strategies.

This test run made clear that comparable climate change focused models of future habitat for PGCN would require a significant climate data preparation effort. As expected for most rare and restricted-range species, Colorado's Plants of Greatest Conservation Need are nearly all highly vulnerable to changing environmental conditions, since they are apparently already adapted to narrow environmental niches. The pace of environmental change is fueling pressure to develop faster methods for identifying climate niche specialization and adaptive capacity, instead of relying on lengthy common garden studies and decades-long demographic monitoring. Adaptive management strategies for these species could be guided by species-specific, model-based investigation of climate constraints and tolerances.


Full model using historic climate normals



Clipped binary model for CODEX



Full model using modeled current climate normals



Full model using future modeled climate normals




Figure 2. *Draba smithii* species distribution models. Top row: predicted model habitat using Daymet climate data, with the model on the left showing the full distribution probability (0-1) and the model on the right showing the CODEX version with probability greater than or equal to 0.5 probability. Bottom row: predicted modeled habitat using climate data from hadgem2.es.1.rcp85 climate model, with the model on the left showing conditions under the current time frame, i.e., 1980-2012, and the one on the right showing a future time frame, the 30-year period centered around 2050.