STUDIES ON THE OVERWINTERING OF THE UREDOSPORE OF THE BLACK STEM RUST OF WHEAT IN THE VICINITY OF LARIMER AND WELD COUNTIES, COLORADO. WINTERS OF 1921-1922-1923

bу

E. A. Lungren

A THESIS SUBMITTED FOR PARTIAL FULFILLMENT
OF THE REQUIREMENTS FOR THE DEGREE
OF
MASTER OF SCIENCE
IN BOTANY

COLORADO AGRICULTURAL COLLEGE
FORT COLLINS, COLORADO
1923

378.788 A O 1923

Approved by

Chairman and Professor of Entomology

e paulo re

Lo Dewson

Graduate Committee

This thesis has been read and approved and recommended for credit.

Head of Department of Botany.

Colorado Agricultural College, Fort Collins, Colorado. May 22, 1923

ACKNOWLEDGEMENTS

The writer takes the opportunity to express his sincere thanks and gratitude to those who have aided him during the present study.

Sincere thanks are extended to Dr. A. K.

Peitersen and Professor C. D. Learn under whose instruction this study was carried to completion, for kindly suggestions and criticisms.

Thanks are extended to Dr. E. C. Stakman and Dr. F. E. Kempton for suggestions and materials used in parts of this study.

Appreciations are extended to the field men for their co-operation and material received.

INTRODUCTION

There is a problem presented to the people in Colorado as to whether the uredo stage of the black stem rust (<u>Puccinia graminis</u> Pers.) can overwinter and cause spring infection without the common barberry (<u>Berberis vulgaris</u>) as an alternate host.

In order to meet this problem the author
has carried on experiments at the Colorado
Agricultural College during the winters of 1921,
1922 and 1923. Field observations have also
been made during the early spring and summer months
of 1921 and 1922. A review of literature
on black stem rust is given and also the results
obtained on the overwintering of the urediniospores.

OVERWINTERING OF THE UREDINIOSPORE

CONTENTS

Part I

- I. Introduction
 - A. History
 - B. Damage caused by Rust
- II. Kinds of Grain Rust in the United States
 - A. Black stem rust
 - 1. Host
 - B. Orange leaf rust
 - C. Yellow stripe rust
 - D. Crown rust on oats
- III. The Causal Organism
 - A. Morphology
 - B. Physiology
 - C. Biologic Forms
 - D. Life History
 - IV. Epidemiology studies
 - A. Relation of barberry rust to the propagation of black stem rust
 - B. Observations on uredinial overwintering at selected stations in the field

 - C. Immediate topography
 D. Method of Germinating spores
 - E. Tabulated Data
 - F. Inoculation experiments
 - G. Weather report
 - H. Spring observations
 - I. Resume of 1922, 1923 work.

Part II

- I. Report on Barberry Eradication from 1918-1923
- II. Summary
- III. Literature Reviewed.

HISTORY OF DISEASE

The cereal rusts have been known for several thousand years. In the bible it is spoken of as "mildew" which is the term applied to rust in England. In those biblical times it was thought to be a pestilence as a judgement of God for the peoples' sins. About 1000 B. C., Solomon prayed to God for clemency, "If there be in the land famine, if there be pestilence mildews, etc." I Kings 8:37.

The Greeks and Romans also feared the rusts. Aristotle described the disease about 350 B. C. Pliny, a Roman, said "wheat was more susceptible to mildew attacks than barley." He also stated that it was the most destructive disease of cereal crops and recommended that the grain be sown early so as to ripen before the attack came.

There were also many ficticious beliefs as to the source of rust. Many believing that certain stars exerted an especially injurious influence from which the so-called secret evil arose, among which are the rusts and mildews.

The first law toward destruction of barberries was passed in Ouren France, in 1660. This was done as they were supposed to have held some mysterious relation to epidemics of wheat rust.

In 1726 Connecticut passed a law restricting cultivation of barberry bushes as it was observed the rust was heavy where the barberries were located.

In 1797 Person found the fungus causing the disease and gave it the name of <u>Puccinia graminis</u>.

The first definite observations on rusts as diseases were published by DeBary in 1865. He proved that the rust was heteroecious and the barberry was the aecial host of <u>Puccinia graminis</u>. Teliospores of <u>Puccinia graminis</u> were germinated and the sporidia placed on the barberry leaf producing aecia and pycnia. With the aecidiospores, wheat was inoculated which produced the red pustule containing the uredinia.

De Bary opened the scientific field on rust investigations and a more complete study of this organism has been made. Pathologists of today are continuing the research on other disease as well as the rusts trying to find a way of combatting them.

DAMAGE DONE BY BLACK STEM RUST.

According to Dr. E. C. Stakman (1) the black stem rust is the limiting factor in grain production in many sections of the United States. It is very destructive on spring wheat and will sometimes cause great loss to winter wheat.

Through a period of years it has been observed that the rust destruction has occurred at irregular intervals. In 1904 the loss caused by rust in Minnesota and the two Dakotas was estimated at \$20,000,000.00. In 1916 it was estimated that South Dakota had prospects of 150 million bushels, but in this same year there was an epidemic of rust and the crop run less than 40 million bushels. A conservative estimate of loss due to black stem rust in 1916 was 180 million bushels. These points bring out the importance of the disease in this country.

KINDS OF GRAIN RUSTS IN THE UNITED STATES.

There are four important grain rusts in the United States, the black stem rust (Puccinia graminis) which will attack our grains and several of the wild grasses. This is the most important rust.

The orange leaf rust (<u>P. triticina</u>), attacking the seed leaves of wheat. Yellow stripe rust (<u>P. glumarium</u>) attacking the stem and leaves, and crown rust of oats (<u>P. coronata</u>).

The black stem rust develops on the common barberry. The uredinial stage occurs on the stems and leaves of the cereals in reddish sori which are long or ovate in shape. It occurs mostly on the stems but can

also be found on the leaves. It can readily be distinguished by the naked eye. It may also occur on other hosts.

Stakman (1):

Hordeum jubatum
Agropyron repens
Agropyron smithii
Agropyron caninum
Hystrix patula

Flymus sp.
Festuca sp.
Koleria custata
Anthoxanthum puelli
Bromus sp.

with <u>Puccinia graminis</u>, although there are forms of this rust which act somewhat differently. For examples, one form will grow on wheat and also on barley but normally will not attack oats or rye. There is also a form which attacks rye and barley but will not attack wheat or oats. All the races or forms can live on several of the wild grasses but not all of them can attack the same grasses. There are thirty—eight different forms of black stem rust and each race has a particular host which it attacks.

The early orange leaf rust, <u>Puccinia triticina</u> attacks the leaves of the plant when they first appear above the ground. The attack takes place in the spring sown grain and in the fall on winter wheat. It can be readily distinguished from the stem rust by the rounded erange colored pustules. Microscopically the spores are

round or circular while those of graminis are ovid or egg shape. (See Plate I).

This rust seems to be continually producing uredospores or summer spores, and live over the winter as uredinial mycelium. Some pathologists consider the damage caused by this rust very slight, only slight delay in maturity, while others say that there are no injurious effects.

The uellow stripe rust, <u>Puccinia glumarium</u>,
was found in Arizona in 1915 on wheat and in the western
United States on barley, rye and many wild grasses.
The pustules can be determined from the graminis
species by being very long and yellow in color. The spores
are much longer than those of graminis. Recently
Dr. Stakman has found considerable stripe in Mexico.
Aside from Arizona it was found in Wyoming. It has never
been found in Colorado.

Crown rust of oats, <u>Puccinia coronata</u>, has a similar life history to that of graminis. The spore stages are the same but the shape of the teliospore can readily be distinguished by the crown shaped apex.

The sporidia go to the buckthorn or Rhamnus sp. and develop the aecia stage. Epidemics of crown rust have occurred in areas where the Rhamnus is grown. The

spores germinate spreading very rapidly from wheat to wheat. Berberis spp. and Mahonia spp. shrubs susceptible to rust.

DESCRIPTION OF PUCCINIA TRITICINA (ORANGE LEAF RUST)

Uredinia, generally on under side of leaf, spores somewhat small, oblong, black and covered by the epidermis. Spores are oblong, obtusely and somewhat pointed above. According to Jackson and Mains (3) the aecial stage has been found. The report is as follows: In the spring of 1919, ten specimens of wintered telia of Puccinia triticina were germinated and sown on 48 species included in 19 genera of the families Ranunculaceae, Boraginaceae and Hydropyllaceae. results were negative, except in the case of two species of Thalictrum repens where pyonia were produced. the spring of 1920, 37 collections of wintered telia were germinated and sown upon 14 species of Thalictrum. Infection resulting in the development of pycnia or aecia was obtained in 84 instances upon 12 of these species. The most vigorous development of aecia was obtained on four exotic species. Infection was only occasionally obtained on the native species used with but a weak development of aecia. The collection of telia giving infection were from Pennsylvania, West Virginia, North Carolina, South Carolina, Alabama, Georgia,

Mississippi, Tennessee, Kentucky, Indiana, Michicgan, Iowa, Minnesota, Iowa, Missouri, Louisiana, Texas, Arizona, California, Washington, and Idaho. Aeciodio-spores obtained from 20 of these cultures were sown back upon seedling wheat producing in every case uredinia typical of <u>Puccinia triticina</u>.

DESCRIPTION OF PUCCINIA GLUMARIUM (YELLOW STRIPE RUST)

Abundance on both surface of leaves and also occurring on the stems. The pustules are arranged in long lines being yellow in color, frequently occupying half of the leaf blade. Uredospores are globose to broadly illipsoid.

Teliospores occur on underside of leaf or on the stem arranged in long lines with a few scattered on the inflorescence. They are oblong and rather club shaped being dark brown or black in color.

Beside the cereals they are found on several of the wild grasses, such as Agropyron spp., Bromus, Elymus and Hordeum.

Puccinia glumarium has no known aecial. It is called spring rust on account of its early appearance, and yellow stripe on account of the sulphur yellow color. The sori are as long as 7 cm. and found on the upper side of the leaf. The uredinio are abundant as early as the beginning of May and it is known to

overwinter easily.

Morphology of Puccinia graminis.

The species <u>Puccinia graminis</u>, has been divided into a number of different forms or races by Ericksson and Henning. Morphologically the races or forms appear nearly identical but differ physiologically in the manner in which they infect hosts of different genera or even of different species. The names of different forms are:

<u>Puccinia graminis tritici</u>, <u>Puccinia graminis secales</u>, <u>Puccinia graminis tritici</u> compacti

Puccinia graminis phlei-pratensis

They may be classified as the following:

Class -- Basidiomycetes
Sub-class --Protobasidiomycetes
Order -- Uredinales
Genus - Puccinia
Species -- Graminis
Variety -- tritici
tritici compacti
secalis
avenae
agrostis
phlei-pratensis

Stakman (2), states that size and shape of urediniospores are similar; however, if a large number of spores are measured and the arithmatical mean or biometrical mode determined, it becomes apparent that there is some difference.

Puccinia graminis tritici, spores are nearly constant in shape, size and color. They are long, but in width are about the same as those of <u>Puccinia graminis</u>

avenae. Shape is eliptical to ovate, color bright
yellow. Mode 32.64 u x 19.71 u.

Puccinia graminis tritici compacti. Quite similar to Puccinia graminis tritici. Color practically the same. Spores slightly longer, shape ovate to ellipsoid, Mode 31.76 u x 19.52 u.

Puccinia graminis secales, uniform in size, shape and color. Shape elongate - elliptic, color dull yellow to grayish, mode 27.23 u x 16.93 u.

Puccinia graminis avenae, color bright cadmium yellow, size and shape very variable, shape ellipsoid or globose. Mode 28.64 x 19.49 u.

Puccinia graminis agrostis spores similar to form above. Somewhat smaller. Mode 22.72 x 16.16 u.

Bio	ological forms	Size Limits	-	Modes
P.	graminis tritici graminis avenae graminis phlei-	23.04-40.96 x 19.20-37.12 x		32.64x19.71 38.64x19.49
P.	pratensis graminis secalis graminis tritici	16.00-29.76 x 17.92-38.72 x	13.44-21.44 2	33.40x17.31 27.23x16.93
	compacti	24.96 -38.08 x	14.40-25.28 3	31.76x19.52

Stakman and Piemeisel worked with the various strains or races to determine the hosts for various biologic forms of Puccinia graminis. The work was carried on in the greenhouse. Inoculations were made on different grasses, especially cereals and the results tabulated.

Agropyron caninum is the host for both P. graminis tritici and P. graminis secales.

Agropyron smithii both P. graminis tritici and secales were present.

Agrostis fatua can easily be infected with P. graminis avenae, also P. graminis coronata.

Avena sativa is infected with P. graminis avenae.

<u>Dactylis glomerata</u> may be infected with either <u>P</u>. graminis avenae or <u>P</u>. graminis phlei-pratensis.

Elymus canadensis can be attacked with ease by the following forms: P. graminis tritici, P. graminis tritici, P. graminis tritici compacti, P. graminis secales.

Cultivated rye (secales) is infected by P. graminis tritici, P. graminis secales, the latter can not infect wheat.

Festuca elatior contained P. graminis phleipratensis, oats, barley and rye may be weakly infected.

Deer successfully infected oats with P. graminis tritici, taken directly from wheat.

Freman and Johnson also succeeded in infecting rye more easily with barley rust than with wheat rust.

On field observations the author has noted several cases where the above could be used for explanation. For example, a rusted rye field may be near barberries heavily rusted. The strain could be P. graminis secales. The next year wheat was planted

on the same field, very little rust appeared, while the bushes were heavily rusted. The P. graminis secales form produced the aecidia on the bushes but this form will not attack wheat.

PHYSIOLOGY

The incubation period of the various forms of Puccinia graminis varies considerable on the same host according to Stakman and Peimeisel (2). The age of the plant, the temperature, the light intensity may effect the rate of development of the rust materially. The cereals are usually susceptible at any age up to the ripening time. The susceptibility of some grasses under greenhouse conditions seems to be greater. Species of Agropyron and Elymus seem to to be extremely susceptible when young and much less when older. Although under natural conditions the plants may be older and become infected. Phleum pratensis and Agrostis alba seem to increase their susceptibility with age. They found other grasses such as Dactylis glomerata to be equally susceptible at all ages. He believes that if such observations were carried on an extensive scale under various conditions and with a large number of species of plants, and different forms of rust, it would seem quite possible to explain any conflicting results obtained by investigators.

Temperature is important in determining the rate

of development of the various biologic forms. distinct differences were noted by Stakman between the different forms. He says that contrary to a fairly general belief, stem rust seems to develop best at fairly high temperatures, provided conditions have been favoragle for infection. The best infection usually results when the temperature is low enough to cause the condensation of moisture in a very fine film on the This usually occurs at night. Then infection leaves. having taken place, the urediospores seem to develop more rapidly at higher temperatures up to a certain point, probably about 75 degrees F. During extremely hot weather, it is difficult to secure heavy infection, probably on account of the difficulty of inducing a film of water to form. Low temperatures below 65 degrees F. inhibit the development of the rust because of the effect of heat on the vigor of the plant. Considerable amount of sunlight is also necessary for development of rust. Cloudy weather may lengthen the incubation period a week or more.

Stakman and Piemeisel (2) inoculated plants, placing them under bell jars for equal lengths of time, keeping them side by side. Some being partially shaded and others kept in direct sunlight. The shade plants

were weakly infected. Plants where the light was completely excluded, developed no rust.

LIFE HISTORY

Puccinia graminis has two stages on barberry, the spermogonial and the aecidial, and two on the grasses; the uredinial and telia. The first appearance on the barberry is the latter part of April or the first of May. In the summer of 1922, the first infected bushes found in Colorado were found on May 8, on sprouts east of Denver. This was the pycnial stage, occurring on the upper surface of the leaves. spots are round and convex above the surface ranging from 3-4 mm. in diameter. On ripening they turn orange in color due to the development of the spermogenes which are the small bodies walled in by a large number of orange-like cells, the spermatia. These small oblong to roundish bodies do not germinate freely and their exact function is not known. They are supposedly gametes which have lost their function.

On the lower surface of the leaf just below the epidermis a large mass of hyphae arise, causing a rounded dome appearance. The hyphae within are shorter giving rise to parallel chains of spores known as acciospores. They are gradullay pushed up each new spore formed below the older ones. The epidermis is soon ruptured, the edges of which revolute and there is formed an orange cup full of chains of aecidiospores.

The older spores at the top are blown away and are replenished by the new produced. The spores germinate readily with long germ tubes. The spore has one well defined germ pore. It is not resistant to drouth.

The aecidiospore is carried by the wind to the grain and here lodges on the stem or leaves. On germinating, the germ tube grows over the surface of the plant and soon finds a stomatal pore. On entering it works its way through the intercellular spaces occasionally sending haustorium into the mesophyll cells.

After several hours this mycelium begins to form a third spore, the urediniospore. Aknot-like hypha is formed just beneath the epidermis some of the branches turning upward forming layers or rows running parallel to the surface. The hyphae then divides into two daughter cells, the lower of which develops a stalk and the upper becomes a urediniospore.

The spore is oval in shape containing a double cell wall the outer being cutinized, containing spines. Two nuclei are present and four germ pores. Anumber of spores forming bursts the epidermis forming a pustule. Several pustules on the stem constitutes a sorus.

It is very easy for the urediniospore to break from the pedicel and be carried by the wind to adjoining plants or other susceptible grains. Here they will

germinate producing mycelium which will penetrate
the epidermis and in turn form more fresh uredo material.
This can be repeated indefinitely as long as conditions
remain favorable. It is also very easy for the mycelium
to go up and down the leaf producing fresh sori.
The linear pustule can be accounted for by the movement
of this mycelium.

Urediniospores may retain their viability for several months depending on the environmental conditions. Drying or freezing conditions they are able to live for about three months. Alternate freezing and thawing kills them in a short time.

Due to the former drying conditions which occur, the mycelium which had produced the urediniospores now begins to produce (at first in the same, afterwards in separate sori) the fourth kind of spore, the teliospore. In P. graminis there are two cells formed, each being a spore and capable of independent germination.

This spore is also formed on a pedicel being club-shaped. The walls are heavy, the contents appearing brown, granular and at first oily. One large nucleus is the center.

The function of the teliospore is to act as a resting spore, tiding the rust over the winter. Due to the heavy cutinized wall it is able to withstand severe 'conditions.

Each cell has one germ pore and when conditions are favorable in the spring, there protrudes through it a rather long tube which divides twice forming four separate cells called the basidium. Each cell under ordinary conditions produces a small stalk on the end of which a basidiospore or sporidia is formed. This spore can germinate indefinitely and form a conidium resembling basidiospore. If one of the detached basidiospores or conidium are carried to the surface of the common barberry leaf its germ tube bores through the cuticle entering the tissues forming mycelium which in turn gives rise to the spermogenes and aecidia.

In connection with the aecial stage it is interesting to know of the number of spores produced which has been worked out by Christman at St. Paul. One aecidium contains 8000 aeciospores. There are on an average 36 aecidiums to one cluster cup. An average of 8 cups on a leaf. Eighty per cent of the leaves were infected on a bush containing 35,000 leaves. From the former there would be 23 million aecidiospores on one leaf and for a bush 7 feet tall, $2\frac{1}{2}$ feet in diameter there would be 64 billion, 512 million aeciospores. At this rate it would not take many bushes to saturate the air with spores in the near vicinity.

STEM RUST EPIDEMIOLOGY

The relation of Barberry rust to the propagation of Black Stem Rust.

Stem rust epidemiology takes up the work of the study of overwintering of the urediniospore and observations of the rust in the field. In finding rust one should be able to account for its presence and obtain the evidence for proof. It is often done by tracing the rust to infected barberries or by finding the uredinia to overwinter, or a study by the use of slides exposed at different elevations to see it the spores are blowing into the area from adjoining territories.

We have been able to trace the rust to infected barberries in several cases. The spread is illustrated by the maps (See Plate 3), one of the property of D.E. Bliss, Greeley, Colorado, the other of P. Dines at Berthoud, Colorado. The rust in the surrounding territory must be accounted for by the presence of these bushes.

Another striking example is the one which was found at Burlington, Colorado, in June, 1922. In the town there were three properties on which 12 heavily infected barberries were growing. It was found that the rust from these bushes had spread for a distance of ten miles or more. It was also noticed that the rust was heavier near the bushes than it was further away. There were several wild grasses such as Agropyron

sp., Elymus spp., and Hordeum spp. heavily infected with the rust which indicated that it can pass from the bushes to wild grasses and in turn to the wheat. This was a local epidemic, rust only existed in the area near the bushes.

In Minnesota, when tracing the spread from infected bushes they find that they can only trace them a short distance before running into a spread from other bushes. This is due to the large number of barberries growing in this State. A number of small local epidemics within a short time leads to a large epidemic which may cover a great territory.

OBSERVATIONS ON UREDINIAL OVERWINTERING AT SELECTED STATIONS IN THE FIELD.

If rust is present and no barberries can be found for several miles, the problem is then a case of overwintering or blowing in from infected areas.

The author has made a series of observations on overwintering by selecting various stations in the northern part of the State where rust occurred in 1921.

The substratum of the rust was selected on the volunteer grains, fall sown grains, wild grasses and straw or sutbble. The varieties and locations are tabulated on the result sheet. Each plot contained an abundance of fresh viable uredinia in the fall.

After locating the plots it is necessary to make notes on the conditions of the host. The fall sown and volunteer grains were vigorous and contained plenty of rust. Some plots were seedlings while otherswere grain which had been planted late in the summer. The stubble was selected from rusted fields after the grain had been cut. An abundance of material was available so as to have plenty for tests throughout the season.

IMMEDIATE TOPOGRAPHY

The plots were located in protected places from animals to avoid any destruction. The stations listed from 1 to 6 were exposed to weather without protection. The material on the stubble was exposed in one plot, another covered with leaves and another placed in a hole and covered. Rusted material was also set out on a ditch bank, one on a south slope and one on a north slope. It is necessary to see if the slope affected the life of the spores.

METHODS OF GERMINATING SPORES.

After the material was located and all notations made, it was then necessary to collect material from each plot at different intervals throughout the year and make germination tests. Two methods were tried the

latter being the better. Take a glass slide and fasten a glass ring to it by the use of vaseline. Place in a few drops of distilled water, sow the spores over this and invert it over the ring. Place the slide in a moist chamber and allow to stand for 48 hours, then make counts. The second method is by taking a slide and placing a small circle of vaseline in the middle. Place on a drop of distilled water and sow the spores on this. Place the slide in a petri dish and the petri dish in the germinator. The germinators used were those in the Colorado Seed Laboratory which are kept at a constant temperature of 20 degrees C., which is the optimum for the best germination of the spores. Counts were made at 48 and 72 hour intervals, before being taken out. The percentage of germination was made by marking off half or fourth of the field and counting the number of spores present; also the number that germinated, and in this way a percentage was obtained.

TABLE I

LIST OF STATIONS WHERE THE PLOTS WERE
LOCATED GIVING HOST, STAGE
AND DATE COLLECTED

				Date
No.	Location	Host	Stage	Located
1	3 mi. S. of Fort			
2	Collins South of City Park	Oats "	Volunteer	Sept.5,1921 Oct.1, 1921
2 3 3	n n n n	Barley	ti	1 1 1 II
3	Experiment Station	Wheat	Heading	
			out	Sept.15,1921
4 4 ₇	2 mi. N.of Greeley	Wheat Oats	Volunteer	Oct.3, 1921
4 ₁ 5 5 ₁	3 mi. N. of Loveland	_	Mature	Oct.8, 1921
ρĪ	11 11 11 11	Barley (wild)	11	11 11 11
6	l Mi. E.ofBerthoud	Oats		u u u
61	и и и и	Barley	·II	tt tt tt

The following gives the data concerning the collection of spores and the percentages of germination. The preparation date is given, that is, the date the spores were collected and placed in the germinators. They were immediately brought in from the field and placed in the germinator by the method already explained. The stations are indicated, the percentage of germination, the germination test and the temperature at date of collection. The date of germination test is the date on which the counts were made. A thermometer was carried to the station when spores were collected. The collections and tests were made at the given intervals from September, 1921 when first collections were made until

the spores failed to germinate, several check
tests were made after no germination was obtained.
The tests were then dropped.

Table II.

PERCENTAGE GERMINATION OF UREDINIOSPORE OF
PUGGINIA GRAMINIS DURING THE WINTER
1921-1922

					
Prepara-	1.	Station	1% of	Condi-	Temperature
tion	Germi-	Number	Germi-	tion	at date of
Date	nation		nation		Collection
0 /2 0 /02	Test	 	Test		
9/16/21	9/20/21	 1	94	Fair	Fair
9/16/21 9/16/21	9/19/21	2	84	11]
9/16/21	9/19/21	3	89	11 -	# #
9/23/21	9/28/21	1	84	11	11
9/16/21 9/23/21 9/24/21	9/28/21	2	50	n	tt
9/24/21	9/28/21				J
	8 A. M.	3	94	ff	11
10/1/21	10/3/21				
70/7/07	8 A. M.	1	45	! !! !	11
10/1/21	10/3/21		7 5 7 7		12
10/1/21	8 A·M· 10/3/21	2	75 Barley	. 11	! 1
10/1/21	8 A.M.	3	45 Oats	II	11
10/5/21		4Wheat	40	11	73°F
10/5/21	10/6/21	4 Oats	85	Good	73 F.
10/7/21	10/10/21				, 5 1 9
	A. M.	2 Oats	no	II	65 F.
10/7/21	10/10/21				
10/0/21	A • M •	2 Barley	87	ti	80 F.
10/8/21	10/11/21 A• M•	6 Barley	07.	11	00 1
10/8/21	10/11/21	o parrey	ופ		80 F.
20/0/21	A.M.	67 Oats	50	Dried	80 F.
10/11/21	10/13/21	-1000			
	A. M.	3 Wheat	96	Good	63 F.
10/15/21	10/17/21	_			
10/15/01	A. M.	1	98	II	62 F.
10/15/21	A. M.	3	99	11	CO TO
10/15/21		J	98	••	62 F.
·	A. M.	2 Oats	95	11	62.F
10/15/21	10/17/21				ON TE
	A. M. 10/17/21 10/20/21 A. M.	2 Barley	95	11	62 F.
10/15/21	10/17/21	6 Oats	93	11	62.F.
10/18/21	10/20/21				
Ì	A • M •	4	75	ii.	Warm
	•	•			
			,	i i	

Table II conintued

				1	
10/18/21	10/20/21	41	Low	Dried	Warm
P. M.	A. M.	Two Tibes	The same	-	
	10/24/21	·			04 15
P. M.	A. M. 10/24/21	1	Low	Dried	84 F.
$A \cdot M \cdot$	A. M.	2 Hordei	50	Fair	79 F.
10/21/21	10/24/21	2 Avenae	50	Fair	79 F.
A . M .			· ·		
10/21/21	10/24/21	2 avenae			
70/07/07	7.0/04/07	telia	Act appropriate		79 F.
10/31/31	10/24/21	2 philei-		Good	79 F.
10/91/91	10/24/21	pratensis	3		
10/21/21	10/24/21	tritici	93	Good	79 F.
10/26/21	10/28/21	6 Avenea		Good	79 F.Clear
10/28/21	10/31/21	8 phlei-	and the state of t		
		pratensis	25	Good	67 F. "
P • M •	A • M •	aveneae			
10/20/21	10/31/21	Z D ~	(low)	G ood	67 F. "
P M.		tritici	74	Good	64 F. "
10/31/21		41	Low	Good	no ·
	A • M •				
10/31/21				Į.	
17 /5 /57	A. M.	4	Low	Good	no
11/5/21	11/7/21	5	20	Good	73.5 F.
P• M• 11/5/21	A. M. 11/7/21				
P. M.	A. M.	5 1	40	Good	73.5 F.
11/5/2 1	11/7/21		}	9000	10.00 P.
A • M •	A • M •	1	75	Good	73.5 F.
11/5/21	11/9/21				
$A \cdot M \cdot 11/7/21$	A. M. 11/9/21	3	90	Good	73.5 F.
$A \cdot M \cdot$	A • M •	6	70	Good	71.5 F.
11/7/21	11/9/21			4000	(TOO LO
A . M .	A. M.	2	50	Good	
11/7/21	11/8/21		21.384	T in	
A. M.	A. M.	Timothy			
11/10/01	j j	Lake	25	Good	73.5 F.
11/12/21 11/12/21	11/14/21	1	75	Good	73.5 F.
P. M.	A. M.	6	50	Good	64. F.
11/12/21	11/14/21)	J	u u u	OT• I•
, , ,	A . M .	Brigh-	Blanchur		
	o page of the same	ton	50	Good	65. F.
	7	į	1	1	

Table II continued

3 3 /3 6 /63	33 /38 /03	· · · · · · · · · · · · · · · · · · ·			
	11/17/21		i de la composición della comp		, de acceptant de la constant de la
A • M •		Greeley			
11/01/01	11/05/01	41			
	11/25/21		120	0000	
A • M •	17/05/07	3]19	30.0.K	.63. F.
	11/25/21	_		00 7 7	0.5
A. M.	77 /50 /07	1	6		. 63. F.
	11/30/21		22	SO G•K	• 63. F.
A. M.	77 /00 /07	Barley			
	11/30/31		70	20 2 7	F0 F
A. M.	70/5/07	Berth	30		. 58. F.
12/2/21 12/2/21 12/2/21	12/5/21	5	No	30 C.K	
10/0/01	12/5/21 12/5/21	- 5 1	No	20 C.K	43. F.
10/0/01 10/0/01	10/5/61	3	∮No .	20 0 · K	43. F.
12/2/21	12/5/21	3 leaf	37.0	20 a 12	47 1
19/9/91	79/5/97	Rust	No		43. F.
12/2/21 12/9/21	12/5/21	5.	No	20 C.K	43. F.
TN/3/NT	12/12/21 A.M.	3	Low 10	20	14.77 TO
10/9/21	12/12/21	1	Low 10 No	20 20	43. F.
10/9/21	12/12/21	1:	100	۵0 '	43. F.
10/0/21	10/10/01	tritici	30%	20	43. F.
1/3/22	1/5/22	3	10%	20	58 F
1/3/22	1/5/22		10/2	٠,	00 F
- 7 9 7 ~ ~	A · M ·	3 "	30%	20	58. F.
1/3/22	1/5/22	8 Barn	no	20	58 F
1/3/22	1/5/22	9 Hole	no	20	58 F
1/13/22	1 /16/22			~ ~ ~	
P.M.		5 !	25%	20	48. F.
1/13/22	1/16/22	5'Elmis	no .	20	48. F.
P • M •					
1/13/22	1 /16/22	5'Agro-			T- 1
P.M.		pyron 5'	1% (20	48. F.
1/13/22	1/16/22	5¹.	30%	20 _	48. F.
$P \cdot M \cdot$					
1/13/22	1/16/22				
P.M.	A - M -	5'H.j.	10%	20	48. F.
1/13/22	1/16/22	5'triti-			
		cina	10%	20	48. F.
1/13/22	1/16/22	5'h.j.		ļ	
- 1 1	artista -	graminis	10%	20	48. F.
	1/16/22	5'wheat			•
A • M •	ļ	grass	no	20	48. F.
	1/16/22	3	no	20	48. F.
A • M •		_	,		
1/13/22	1/16/22	3	5%	20	48. F.
	,	,	ŀ		
				•	

Table II continued

		{		ţ	
	2/13/22	5'hordei	J. no	20	42. F.
2/11/22	1/13/22	5'h.	no	20	42. F.
2/11/22		jubatum 5' " 5'phelum 5'h.	no 20	20 20	42. F.
0.420.400	0/07/00	jubatum	no	20	36. F.
2/19/22	2/21/22	5 H. jubatum	no	20	36. F.
2/19/22	2/21/22	5 h.			
2/28/22	3/2/22	jubatum 5 h.	ño	20	36. F.
, ,	, ,	jubatum	5 %	20	40. F.
A • M •	3/12/22	5 h. jubatum	no	20	34. F.
3/10/22	3.12/22	5 h. jubatum	no	* 20	34. F.
4/20/22	4/23/22	5 h.			
		jubatum	no	20	40. F.

Table III indicates the date of first germination and the percentage of viable spores; also the date when no spore germination could be obtained. One can compare the time elapsed from high germination to zero for the different hosts.

It can be noted by this data that the longest viable urenidiospore was on Hordeum jubatum.

TABLE III.

		 			
Sta. No.	Host	Date of First Germi- nation Test	% via- bility of Spores	Final Date of Germi- nation Test	1
1	Oats	September		December	
2	Oats	15, 1921 October	94 84	12, 1921: January	0
		1, 1921		12, 1922	0
21	Barley	October 1, 1921	75	Janúary 13, 1922	
3	Wheat	September		January	Ŭ
		15, 1921	89	16, 1922	0
4	Wheat	October		J	
		3, 1921	40		
4_{7}	Oats	October 3,		November	•
		1921	85	2, 1921	0
5	Timothy	October 8,		December	
		1921	20	2, 1921	0
51	Barley	2		March 12,	
_	(wild)	8, 1921	40	1922	0
6	Oats	October		January	
		8, 1921	50	<u>16, 1922</u>	0

The graph indicates the highest germination of urediniospore from any one plot by months from September 1921 to April, 1922.

The following chart indicates the test made on the uredo material collected on stubble. Ten small bunches were made containing several stems to a bundle. These were placed in an erect position out in an open field exposed to all weather conditions, protected by an open screen.

The plot was established September 15, 1921 and tests made at intervals until no germination was obtained.

The results are tabulated as to date of collection, and placing the spores in the germinator, date on which the counts were made, percentage germination, remarks and temperatures at time of collection.

TABLE IV.
STATION NO. 4 NEAR BOTANY
ANNEX STUBBLE TEST III.

	Date	, Date	100	T	
No.	Placed	of	Germi-		Temperature
	in	Germi-	nation	Remarks	on
	!	nation			Collection
604	9/27/21	9/30/21	No	Dried	
538	tt.	9/29/21			
		A. M.	15		: (
537	11	" A.M.	65		
614	11	" A. M.	45		
431		9/30/21	n e	TET	
237	11	A.M. 9/29/21	75	Warm	!
201		A.M.	25		
416	tt	9/30/21	۵٥		
		A.M.	10		
245	II ·	9/30/21			
•		A.M.	3 5	Cool	
435	1.11	9/30/21			
	-	A • M •	60		
434	11	9/30/21			
en a	10/7/07	A. M.	no	Dried	
574	10/3/21	10/5/21			
604	P.M. 10/3/21	A•M• 10/5/21	no	roor sild	e Cool in 8:45A.
001	P.M.	A-M-	Fair	Dry	n
538	10/3/21	A·M· 10/5/21	1.00.1.1	Diy	
	P.M.	A.M.	75	Good	11
537	10/3/21	10/5/21	- and annual		
	P.M.	A.M.	Fair	On Plate	11
615	10/3/81	10/5/21			
=	$P \cdot M$.	A.M.	3 5	Fair germ	• 11
431	10/3/21	10/5/31		~	
O IZ IZ	P.M.	A.M.	85	Good	tı
2 37	10/3/21	10/5/21	-	_	
416	P.M.	A • M •	25	Dry	11
#TO	10/3/21	10/5/21	Pois	D3-3-1	2.77
245	10/3/21	A. M. 10/5/21	Fair	Dried out	II
₩.	+0/0/81	A.M.	Low	11 11	n
		t Zr ● INI ●	1 TOW	11	48

-28-

Table Inv continued

435	10/3/21	10/5/21			
434 574 604 538 537 615	10/3/21 10/13/21	A. M. 10/5/21 10/7/21	no 87 no Low "	Very Good Dried Good " Dried Conditions	Clear 0 64 11 11
431 237 416 245 435 434 435	" " " " " " " " "	11 11 11 11 11/6/21	Low 10 20 Low 45	Good Good " " Dried Good	11 11 11 11 11
416 604 615 574 431 434 235 538 574	3 P.M. 11/3/21 11/3/21 11/3/21 11/3/21 11/3/21 11/3/21 11/3/21 11/3/21 11/3/21 11/3/21 9/27/21	A· M· 11/6/21 11/6/21 11/6/21 11/6/21 11/6/21 11/6/21 11/6/21 11/6/21 11/6/21 11/6/21	No 10 15 no n Low No	Good " " " " Dried Dvied Good Poor culture	68 n n n n n n n n
537 604 431 237	12/20/21 12/20/21 12/20/21 12/20/21	12/22/21 12/22/21 12/22/21 12/22/21	n n Low	Dry 32.5 32.5 32.5	32.5 32.5 32.5
538 435	12/20/21	12/22/21	1% No Low		32.5 32.5 32.5
537 574 416 434 245 574 416 537 604	12/25/22 12/25/22 12/25/22	12/22/21 12/22/21 12/22/21 12/22/21 12/22/21 12/27/22 12/27/22 12/27/22	NO n n n n n n n		32.5 32.5 32.5 32.5 32.5 28 28 28

Table IV. continued

435	12/25/22	3 12/2	7/221	No	28
431 '	12/25/22	3 12 2	7/22	п	28
343	12/25/22				28
5 38	12/25/22	3 12/2'	7/22	ti .	28
237	12/25/22	3/12/2	7/22	[‡	28

A bundle of stubble was placed in a hole and covered with straw. It was protected from winds and extreme cold weather. It was found to contain fresh uredo material January 27, and no spores were found viable.

A similar test was made on fresh material placed on the ground and covered with leaves. The results were the same as the above. A test was made also of spores on stubble which was placed in a small shed protected from all moisture and wind. No germination was to be had.

Plots containing rusted material were placed on the side of a ditch, one on the north side and the other on the south. The plot on the south side of the ditch retained viable spores longer than did the one on the north. The tabulated results are as follows:

TABLE V. STUBBLE PLACED IN HOLE
Date placed in, November 8, 1921

	Date of Germination	% Germination	Temperature on date of Collection
	Test		
1/25/22	11/28/22	No	28

TABLE VI. STUBBLE WHICH HAD BEEN IN BARN Date placed in, November 8, 1921

Date of Collection	Date of Germination test	Percentage Germination	Temperature on date of Collection
1/25/22	1/27/22	No	280

TABLE VII. PLOTS EXPOSED ON DITCH BANK Date set out, November 7, 1921

A STATE OF THE STA	Date Collected	nation	% Germi- nation	Temperature on date of Collection
	<u> </u>	Test	<u> </u>	
South Pot	11/28/21	11/30/21	45	58°F•
a n	11/28/21	11/30/21	no	58 F.
North pot	12/20/21	12/22/21	11	32 F.
South pot	12/20/21	12/22/21	11	32 F.

INOCULATION EXPERIMENTS

A series of pots were planted in the greenhouse with different varieties of grasses and inoculations made to establish rusted material for field purposes, and to observe rust development.

The material received for inoculum was of the following nature.

Host	From		<u>Kind</u>	Collector	Date	_
					Collec	
Wheat	Lincoln, Nebr.		P.g.	${ t Peltier}$	9/26	/21
Ħ	St.Paul, Minn.	,	P.g.	O.S. Ammodt	10/7	/21
Erickson	Pullman, Wash.		P.g.	Humphrey	10/7	/21
Wheat	Manhattan, Kan,		P•g•	C.O.Johnstor		

TABLE VII.

GREENHOUSE PLANTINGS

Pot No.	Variet	ty	Date Planted
1	Marquis		9/16/21
2	Turkey I	Red	u
3	Kanred		
	Marquis	R.R.	Ħ
4 5	Calif. I		
b	(Barley)		H
Cl		(10922)	10/7/21
C2	11	` #	n , ,
C3	11	11	11
C4	11	II.	tt .
Dĺ	11	II	ft
D2	tf	11	:
D3	11	n	11
D4	f t	ff	11
El	Oats		II.
ES	ti .		If
Fl	Timothy		н
F2	11		Ħ

TABLE IX.
INOCULATIONS

Pot No.	Variety	Inoculum	From	Date
D1	Defiance	P.g.	Nebraska	10/6/21
Cl	Defiance	n _	u	††
3	Kanred	tt	P. Davies	11
5	Calif. Pearl			
	(Barley)	TI .	Sta. No. 21	tt
4	Marquis R.R.	H	Sta. No. 4	11

TABLE X.

RESULTS OF INOCUATIONS.

		773 3		Date	
Date		Kind	_		Doman
Inoculated		of	From		Degree
	Varieties	Rust		Earliest	of
	2.			Infection	Infection
10/6/21	Seedling		Lincoln,		
	(Defiance)	P.g	Nebraska	no	no
11	Seedling				{ }
	(Marquis)	11	Greeley	11	tt
11	Seedling		•		÷
-	(Calif.		Fort		1
	Pearl		Collins	11	n
	Barley)	11	OOTITIE		
t t			Fort		-
,,	Seedling	ST.		t	u
	(Defiance)	,,	Collins	· · ·	**
11	Seedling				
	(Kanred)	11	P.A.Davi	es "	t1

The results were negative due to a blight which attacked the plants after they had been in the germinator. The results of the 1923 inoculation experiments were very successful as you will see later.

METHOD OF MAKING AN INOCULATION TEST.

After the seedlings have grown to 4 or 5 inches it is then time for inoculation. Obtain fresh uredo material and place several spores on a flat needle. Moisten the leaves with distilled water, run the needle with the spores down over theleaf and in this way establish a thin film of moisture with the spores suspended. Then place plants in incubator. Another successful method is to apply, or push spores on the moistened leaf and with the thumb and forefinger run

over the leaf. This is all right when one is working with but one strain. If several strains are used the other method is best as one can take care and not mix the forms.

The pots are then incubated for 48-72 hours in a glass covered incubator or moist chamber. After this place them out on a bench until the pustules begin to form. Then place out of doors during the daytime in good weather to harden. It takes about 6 days for rust to form after incubation.

TABLE XI.

WEATHER REPORT

Date of		
Collection	Station 1	Station 2
		- Con Annual Control
9/15/21	Fair	Fair
9/16/21	Fair	Fair
9/17/21	Fair	Fair
9/18/21	Fair rain in P.M.	Fair
	Fair	Fair - rain
9/20/21	Windy - frost	Windy - frost
	(night)	
	Fair	Fair
	Fair	Fair
9/23/21	Fair	Fair
	Fair - windy	Fair - windy
	Fair	Fair
9/26/21	Fair	· Fai r
9/27/21	Fair	Fair-
9/28/21	Fair _	Fair
9/29/2 1	Cool	Cool
9/30/21	Fair	Fair
10/1/21	Cloudy	Eloudy
10/2/21	Fair	Fair
10/3/21	Fair	Fair
10/4/21	Warm	Warm
10/5/21	Warm	Warm
10/6/21	Cloudy all day	Cloudy - rain
	rain in A.M.	in A.M.
	<u>.</u>	

TABLE XI continued

```
10/7/21
                 Cool in A.m. clear Cool in A. M. clear
10/8/21
                 Warm
                                      Warm
10/9/21
                                      11
                 Clear - windy in
10/10/21
                 A. M.
10/11/21
                 Warm
                                      Cloudy wind 64° F.P.M.
10/13/21
                 Clear P.M. cloudy
                  wind P.M.
                 Cloudy 52 degreesF. 65 F. P.M.
10/14/21
10/15/21
                 Cloudy 63
                                     73
                                         F. P.M.
                 Clear 70
10/16/21
                                     63 F. P.M.
10/17/21
                 Clear 65
10/18/21
                 Clear 60
                                     75 F. cloudy
10/19/21
                 Clear 52
                                     70 F. clear
                                     73 F. "
10/20/21
                 Clear 70
10/21/21
                 Clear 56
                                     71 F.
10/22/21
                 Clear 58
                                     75 F.
                                     68 F. "
10/23/21
                 Clear 55
                 Cloudy 52 F.
10/24/21
                                     70 windy - rain in P.M.
10/25/21
                                     54 F. windy - clear
                 Cloudy - windy
                  49 F.
10/26/21
                 Clear 48 F.
                                     63 F. clear
10/27/21
                 " 55 F.
                                     66 F. cloudy
                 " 49. F.
10/28/21
                                     62 F. clear
10/29/21
                 " 64 F.
                                     65.8 F. "
                 " 62 F.
10/30/21
                                     65 F.
10/31/21
                 " 46.3 F.
                                     67 F.
11/1/21
                 " 53 F.
                                     64 F.
11/2/21
                 Clear 58.5 F.
                                     70 F. clear
11/3/21
                 " 10 A.M. 68 F.
                                     68 F. " 3 P.M.
11/4/21
                    9:30 A.M. 55 F.
                                     68 F.
                                           11
                                              2 P.M.
11/5/21
                 55 F.
                                     58 F.
                                            " 3 P.M.
11/6/21
                 57 F.
                                     58 F.
                                              3 P.M.
11/7/21
                 53.5 F.
                                     51.5 F. cloudy 4 P.M.
11/8/21
                 Cloudy rain 38.5 F. 48. F.
11/9/21
                 " 34 F.
                                     53 F. clear
11/10/21
                 Clear 42 F.
                                     63 F. "
11/11/21
                 " 64 F.
                                     64 F. Cloudy
12/1/21
                 Clear windy 42.5F
                                     44 F. windy
12/2/21
                 " 45.5 F.
                                     43 F. windy
12/3/21
                 Snow 4 in. 45 F.
                                     43 F.
12/4/21
                 Snow .70pp. 45 F.
                                     43 F.
12/5/21
12/7/21
                 Clear - 39 F.
                                     42.5 F. clear
12/8/21
                    38 F.
                                     Clear
```

TABLE XI. continued

```
12/10/21
12/11/21
12/12/21
                 Clear 65 F.
                                    | Clear 73 F.
                                    # 54 F.
12/13/21
                 " 53.5 F.
                                     " 60 F.
                 " 48.5F.
12/14/21
12/15/21
                 " 51 F.
                                     # 67.5 F.
12/16/21
                 " - windy 22.5 F.
                                     "23.5 F. windy
                 # 23 F.
12/17/21
                                     30 F.
12/18/21
                 " 25 F.
                                     28.5 F.
                                    29.5 F.
                 "-30 F.
12/19/21
                 Cloudy 32.5 F.
12/20/21
                                     Cloudy - dew 32.5 F.
12/21/21
                 " heavy dew 30 E. Cloudy - 32.5 F.
12/22/21
                 30 F.
                                     32.5 F.
12/23/21
                 Snow - cloudy 20 F. Snow - cloudy 17 F.
                                   Ħ
12/24/21
                         " 21 F.
                                              # 17 F.
                         " 25 F.
                                              " 18 F.
                 11
                                    11
12/25/21
                 11
                                     11
                         " 20 F.
                                              " 25 F.
12/26/21
12/27/21
                 Clear 23 F.
                                    Clear 29 F.
12/28/21
                 " 22 F.
                                     " 30 F.
                 " 28 F.
                                    " 35 F.
12/29/21
                                    # 30 F.
12/30/21
                 " 27 F.
                                     " 40 F.
12/31/21
                 " 30 F.
                                     #-55 F.
1/1/22
                 " 30 F.
1/2/22
                 Cloudy, snow 2 in.
                 pp. .24, 31.5 F.
                                     Snow 34 F.
1/4/22
                 Clear - snow 12 F.
                                     Clear 33 F.
1/5/22
                 " 15 F.
                                     " 32 F.
                " 17 F.
1/6/22
                                     " 30.5 F.
1/7/22
                 " 20. F.
                                     "33 F.
                 " 25 F.
1/8/22
                                     " 45 F.
                 " 29 F.
1/9/22
                                    " 38 F.
1/10/22
                 " 40 F.
                                     C 37.5 F.
                 " - windy 32 F.
1/11/22
                                     " 38.5 F.
1/12/22
                 32 F.
                                     38.5 F.
1/13/22
                 Clear 48.5 F.
                                     38.5 F.
1/14/22
                 " 30 F•
                                     56 F.
1/14/22
                 Clear 30 F.
                                     45 F.
1/15/22
                 _____
                                     _____
1/16/22
                 Clear 32.5 F.
                                     38.5 F.
1/17/22
                 Snow 20 F.
                                     18.5 F.
1/18/22
1/19/22
                 " 14 F.
                                     10 F. snow
                25 F.
                                    14 F. "
1/20/22
                Clear 12 F.
                                    26 F. clear
                " 20 F.
                                   38 F. "
1/21/22
                " 18 F.
1/22/22
                                    40 F. "
                " 30 F.
                                    44 F. "
1/23/22
```

TABLE XI. continued

	 	
1/24/22 1/25/23 1/26/22 1/27/22 1/28/22 1/29/22	Clear 28 F. " 30 F. " " " "	36 F. clear 40 F. "
1/31/22 2/1/22 2/2/22 2/3/22	11 11 11	
2/4/22 2/5/22 2/6/22	II Snow II	,

SPRING OBSERVATIONS

Observations were made inthe early spring of 1922 to find the first appearance of rust on grasses. No stem rust was found until June 5, 1921, south of Wray, Colorado. Leaf rust was found May 28, 1922 and June 3, 1921 at Sterling, Colorado. Other reports of leaf rust were sent in later by county agents.

The map indicates the locations of stem rust in the State, observations were made during the summer of 1922. The largest epidemics were traced to the barberry.

OVERWINTERING OF UREDINIOSPORE 1922 - 1923

No field plots could be located in this section due to the scarcity of rust. The work for the year was devoted largely to inoculation tests as well as overwintering.

Germination of rust \underline{P}_{\bullet} graminis tritici collected the summer of 1922.

TABLE XII December 8, 1922 set in germinator December 10, 1922 set in germinator December 11, 1922 set in germinator December 15, 1922 set in germinator " 20 F. " " "

No germination could be obtained from this material.

A TEST OF STUBBLE COLLECTED IN 1922 GAVE THE FOLLOWING RESULTS.

Light petri dishes each containing red spores on distilled water medium; 20 degrees and 30 degrees. Set in germinator December 15, 1922

TABLE XIII.

No.	December 18 A.M. % germination	December 20, 1922 % germination	Temperatu re
1 2 3 4 5 6 7 8	no n n n n	no n n n n	200 11 11 11 11

TABLE XIV
WHEAT PLANTED IN SIX INCH POTS IN GREENHOUSE
NOVEMBER 2, 1922

2	Pots	13274	Certified	Marquis
11	#1	13408	tt .	Kanred
11	.it	13407 13409	tt	
11	11	13409	11	u
11	ff	1310	11	!!

These pots were inoculated December 28, with P. graminis from St. Paul. Set in incubator 48 hours and then set out on bench. January 3, 1923 flecks appearing, and rust starting to develop on the two pots containing Marquis. Plants were set out in daytime to harden.

January 12, 1923, the rusted pots were exposed to out of door conditions. The spores of inoculated material lived only a short time. The results are given below:

TABLE XV.

Date set in	Date of		Temperature
Germinator	Germination	<u>Germination</u>	
January 19, 1923	January 22,1923	85	20
January 26,1923	January 29,1923	30	11
January 26,-1923			. tt
February 21,1923			11
February 24,1923			11

January 30, 1933, Marquis wheat was planted in one large plot in the greenhouse for further inoculations.

February 18, 1923, the wheat was ready for inoculation. The inoculation was by the thumb and finger method, already described.

My March 5, every leaf of the plots was covered with rust. It was set out on this date one half being exposed and the other part covered. Material was collected on March 9, 1923, and thespores were found to

germinate 30 per cent. On a revisit to the plot it was found to have been destroyed so no other tests were made.

SUMMARY OF WORK DONE

- (1) From the enormous losses caused by black stem rust as shown in the past, compared to the rate of growth of our present population, it is necessary to apply some problem for the destruction of this disease, no matter what the sacrifice may be.
- (2) The black stem rust is most common and the most destructive of all the rusts.

Orange leaf rust attacks the plants when they are very young and does very little damage.

Yellow stripe rust is characterized by the long pustules and sulphur color. It does not occur in Colorado but where it does occur there is some loss from it.

Corwn rust on oats. The telia spore names this rust by the crown shaped apex. Where you find this rust you find the Rhamnus spp. of shrub which is the alternate host. Severe attacks in the east have caused much damage.

(3) We find several rust spores which appear to be alike but act differently to various hosts. This can be accounted for by having rust on barberries and no rust

The various races or strains have been analyzed by Dr. E. C. Stakman as to grains and wild grasses they attack.

The rust is heteroecious thus having an alternate host. In the case of graminis it is the Berberis spp.

The aecial or cluster cup develops here producing aecia which attack the wheat and here starts the epidemic. In the fall the mycelium changes to the winter telia which rests over and the next spring germinates, making the new attack to the shrub; thus completing the cycle.

(4) The epidimiology studies brings out the importance of overwintering. There is convincing evidence of the spread of rust from barberries, but it has also been observed that some rust exists where no barberries are present. This leads to two problems; overwintering or blowing in from infected areas.

Field plots were selected for the overwintering experiment. Various grains and grasses were selected at different locations in the northern part of the state. Notes were taken as to the topography and condition of the host. Plots were exposed and some covered.

In germinating spores the best method was found by using the slides and Petri dishes. Spores germinated best at 20 degrees C.

From tests made at intervals throughout the season on all 9 plots there were no spores that retained their viability. The list below gives the number of months the spores lived in each plot.

TABLE XVI

Plot	Length of germination
No.	Period
1	Four months
2	Three months
21 3 4	Three months
3	Four months
4	One month
	One month
4 1	Two months
5า	Five months
5 ₁ 6	Three months

It was also noticed that the percentage of germination decreased from one month to the next.

Hordeum jubatum contained P. graminis spores which reatined life the longest. They germinated 5 per cent March 2. Several check tests were taken later but no germination was to be had.

Red spores will not overwinter on stubble either exposed or protected.

Alternate freezing and thawing kills the urediniospores. Favorable conditions in the fall causes them to germinate and then die.

A south slope is more favorable for germination than a north slope.

Barberries become infected in this State as early as May 5.

Leaf rust appears around the first of May. Stem rust around the first of June.

Plants were successfully inoculated in 1923 and plenty of rust material was to be had. Inoculated material when exposed, contains viable spores for only a short time. No spores were found to germinate after March 19, 1923.

At present there has been 24,001 barberry bushes found on 1642 properties, out of which 23,822 bushes have been removed from 1639 properties. On the resurvey of 1922, 18 per cent of the bushes were found sprouting. Several of these have been treated with chemicals. Our recent spring resurvey shows that no sprouts are coming up so chemical eradication is very successful. The removal of the bushes has eliminated a great deal of rust and it is the author's opinion that complete eradication will aid to solve the rust problem.

If conditions were more favorable for the uredinial to live thru the month of March, the author believes that overwintering could be obtained. To answer this question exactly it would be necessary to run the experiment


thru a series of years. Stations should be located in each part of the state and tests made from each. It may be possible that there are certain sections where overwintering may be successful but as to the northern part of Colorado we conclude that the spores do not live through the winter of 1921 and 1922.

REFERENCES


- (1) Stakman, E. C., A study in cereal rusts. Minn. Exp. Sta., Bulletin No. 138 (1914).
- (2) Stakman, E. C. and Piemeisel, F. J., Biologic forms of Puccinia graminis on cereals and grasses. Jr. Agr. Res. 10:a; pp. 429-496, 1917.
- (3) Jackson, H. A. and Mains, E. B., Abstract
 Phytopathology 2: 1: 40.
- (4) Christman, A. H., Sexual reproduction in the rusts. Bot. Gaz. V. 39, 1908.
- (5) Olive, E. W., Rusts of cereals and other plants.
 Agr. Exp. Sta., Bulletin 109, 1908.
- (6) Hungerford, C. W., Rust in seed wheat and its relation to seedling infection. Jr. Agr. Res. 6/5/20
- (7) South Dakota Bulletin 109, 1908 Rusts on cereals and other plants.
- (8) Melchers, Leo E. and Parker, J. G., Rust resistance in winter wheat varieties. U.S.D.A. Bulletin No. 1046.
- (9) Kempton, F. E., The origin and development of the pycnidium. Got. Gaz. Vol.LXVIII, Oct. 1919.
- (10) Stakman, E. C., Destroy the Common Barberry. Farmers Bulletin No. 1058.
- (11) Stakman, E. C., The black stem rust and the barberry. Farmers' Bulletin No. 796.
- (12) Learn, C. D., Save the Wheat. Colorado Agr. Exp. Sta, ExtensionBulletin No. Series 1, No. 176-A 1921.
- (13) Robbins, W. W., Eradicate the common barberry, Colorado Agricultural Experiment Station, Extension Bulletin No. 147, Series 1. Sept., 1918.
- (14) Thompson, Noel T., Kill the common barberry with chemicals. Department Bulletin 268, 1923.

- (15) Hungerford, C. W., Ruse in seed wheat and its relation to seedling infection Jr. Agr. Res. 19, No. 6., 1920.
- (16) Duggar, Book on Plant Pathology
- (17) Strasburg, Textbook in Botany.
- (18) Martin, Text book in Botany.


Plate No. I



Leaf Rust Uredinia L.P.




Stem Rust Uredinia H.P.



Stem Rust Uredinia in Lactic Acid H.P.

Plate II. Cluster Gups on Barberry Leaves








Plate IX
Germinators Used for Incubating Inoculated Material







Plate V

Rust on Inoculated Grain Set out to Harden







BARBERRY REPORT FROM BEGINNING OF CAMPAIGN TO DECEMBER 31, 1922

TABLE OF CONTENTS

	rage
Title Page	
Table of Contents	
Resume of Barberry Eradication in Colorado from beginning of Campaign to December 31, 1921.	59
Introduction	5 9
State Law	65
State Financial Aid	7 5
Publicity	76
Cities and Towns Completed in Survey from beginning of campaign to December 31, 1931	- 83
Counties covered in farm-to-farm survey from beginning of campaign to December 31, 1921	88
Legend	90
Progress Map, Resurvey from Beginning of Campaign to December 31, 1921	
State Totals, Form A	941
Original Survey, Form B	95
Resurvey, Form C	96
Legend	. 97
Map, Location of escaped bushes	98
Properties where escaped bushes were found	99
Average Cost by years	100
General Discussion	. 101

TABLE OF CONTENTS (Continued)

Progress of Eradication Campaign, January 1 to December 31, 1922.

Introduction	105
Personell	107
State Financial Aid	108
Publicity	109
Illustrations	110-115
Legend	116
Progress Map, Farm-to-farm Survey from beginning of campaign to December 31, 1922	117
Legend	118
Progress Map, Resurvey, 1922	119
Progress Map, Resurvey from beginning of Campign to December 31, 1922	120
State Totals, Form A	121
Original Survey, Form B	122
Resurvey, Form C	123
Results of Barberry Eradication, Colorado 1918 to December 31, 1922. Counties Totals	124
Cities and Towns covered, 1922, original Survey	125
Legand	126
Map, Location of Escaped Areas	127
List of Properties where escaped bushes were found	128

TABLE OF CONTENTS (Continued).

Resurvey Results	129-130
Summary of Barberry Survey by Federal Department	13 1
Summary of Costs, Original Survey	132
Summary of Costs per Acre	133
Resuvey Costs	134
Discussion of Resurvey	135
Epidemiology Studies	139
Megend	146
Chemical Eradication	147
History of Plantings	148
Discussion of year's Work	150

RESUME OF BARBERRY ERADICATION IN COLORADO FROM BEGINNING OF CAMPAIGN TO DECEMBER 31, 1921

Introduction

The first stepr were taken toward the eradication of the Common Barberry in the state of Colorado in the year 1918 thru the co-operation of the U. S. Bureau of Plant Industry, the Division of Extension of the Colorado Agricultural College, and the Department of Botany of the Colorado Agricultural College. this time a general and rather hasty survey was made for the purpose of determining the number and general location of the bushes within the boundaries of the state and the magnitude of the task of permanently removing them. At this time, Dr. Gillette, of the Extension Department, declared the Common Barberry bush to be a pest and as such to come under the laws of the state concerning such pests. In conjunction with this act a quarantine was placed upon the shrub on January 20, 1918.

Thru the efforts of the Federal Office a man was placed in the field this first year. His duties consited of making inspections of properties and urging the removal of Barberry Bushes were found. He

succeeded in locating many bushes and also in having a great many of them removed. However, his greatest accomplishment was the paving of the way for his successors by bringing before the public the true character of Berberis vulgaris and educating the people as to the necessity of its complete removal from the small grain growing states. During this first season work was done in the counties of Boulder, Larimer and Weld.

Since the beginning of the campaign has been carried on in Colorado under three main heads: publicity, inspection and eradication.

early in the campaign as it was felt that the success of the work depended upon the co-operation of the people as gained thru a thoro understanding of the nature and object of the movement. Less has been done along this line with each succeeding year, the people becoming more familiar with the work as the campaign advanced. During the last two years publicity has been limited to the districution of bulletins and life cycle cards and an occasional news article in the town paper for the purpose of freshening the people's memory.

"Inspection" and "eradication" have gone hand in hand, especially during the later years of the campaign. At first the bushes were merely tagged and

left for owners to remove. After the majority of the bushes were removed in this way it was found to be more practical, except in the instances of hedges or other large findings, to remove the bush when found as this saved the expense and time involved in making a return trip, sometimes of many miles, perhaps to find the bush still in.

The greatest number of bushes were removed during the year 1919 when six men were placed in the field. They walked and used bicycles according to the nature of the territory in which they were working and during the latter part of the year they were honored with an automobile of rather doubtful vintage and still more doubtful characteristics. The area covered was comparatively small as most of the work was done in the cities and towns, some rural scouting was attempted in the counties of Bent, Crowly, Kiowa, and Prowers and also a small amount in Larimer and Weld.

The scouts wore arm bands to designate their calling and took care to have the people understand the nature of their visit. Scouts neglecting to do this in the early days of the campaign were known to have been taken as chicken thieves and otherwise placed in embarassing possitions.

During the year 1919, 17,919 bushes were found on 1330 properties. During the year 1919, 1,747

bushes were found on 158 properties. Thus a total of 19,666 bushes were found from the beginning of the campaign to December 31, 1919 and these on 1,488 properties. Up to this time the scouts had been successful in having 16,909 bushes removed from 1,239 properties.

Since July 1, 1919, the work has come under two sub-heads: original survey and resurvey. Resurvey is very important because of the fact that, regardless of the care exerted in removing the bushes, there are great possibilities of sprouts shooting up the following spring. This in fact presents a very real problem. In addition, resurvey often leads to the discovery of bushes missed the first time over.

The following summer of 1920 was marked by a scarcity of dependable men due to the high wages being paid in all fields of endeavor. As a result much difficulty was experienced in securing reliable scouts and for the greater part of the season the State Leader worked alone, covering the field in a car. An assistant was put on for a month working chiefly in the cities and towsn of Boulder and Larimer counties. During this season several large findings were made, 1976 bushes being located on 25 properties.

By June of the following year, there were four men in the field, forming two teams of two scouts each, each time having a car. Several rust epidemics

broke out during the season and prompt investigation was made in each instance. None of them were as serious as reported and only in a few places was the damage done of any great consequence and this on spring grain. Most of these out breaks seemed to be the result of wind blown urediniospores coming in from outside sources as no barberries could be found in the immediate vicinity. In a few instances the outbreaks were traced to barberry and the cause removed. Much of the work during 1921 was in regions where the farms were well scattered and considerable distances apart most of the ground being covered in cars. Also considerable resurvey was done. At the end of the season, 22548 bushes were reported found on 1596 properties and of these 22,457 were removed from all but three properties. Also 3226 sprouts were found and removed up to December 31, 1921.

By the end of the year 1921, the most important sections in Colorado had been given an original survey and a considerable part of this area resurveyed. The persistency of the bush was by this time well realized and very seldom did the scout feel sure that he would not be greeted by a sprout the following season. Especailly in rocky and wooded areas did the bush seem unconquerable for no amount of digging could reach all the roots. This problem will no doubt be successfully met in the future years thru chemical eradication a small amount.

of which was attempted in the state this season and with results. Next seasons' work will involve the use of chemicals to a considerable degree since it is much cheaper in the end than continued digging and much more effective.

STATE LAW IN COLORADO

Regarding the Common Berry.

Colorado's Amended Pest Law:

An Act to Provide for the Protection of Trees,
Fruits, Crops and Pasturage against the Ravages of
insects, Rodents, Weeds and Plant Diseases; to
Provide for the Formation of Pest Inspection Districts,
to be under the General Authority of the State Entomologist; and to Make an Appropriation for Carrying out
the Provisions of this Act.

Be it Enacted by the General Assembly of the State of Colorado:

Section 6. Whenever the State Entomologist, his deputy, or county inspector, has reason to believe, or has been credibly informed, that within the State there exist lands infested by pests which are liable to spread to the injury of others, it shall be his duty to make investigation of the suspected premises, and if they are found so infested, shall notify the owner or persons in charge of control of such premises, in writing, of the nature, extent and location of the infection and demand that within a specified time certain specified work shall be done on the infested premises for the extermination.

of such pests, and if the occupant of the infested property refuses or fails to de effective work in the premises in accordance with the instructions of the officer in charge, such officer shall take possession of the infested premises, and do the work necessary for the extermination of said pests, as provided in this Act. The reasonable and necessary expenses for doing such shall be paid for by the county without unnecessary delay, upon filing with the county commissioners of the certificate of the State Entomologist showing the necessity for such work, the reasonable cost and expense thereof, and giving a description of the land or property upon which the work was done. And, upon filing such certificate, such charges shall become alien of the property treated, collected as real estate, or, in the event of personal property, it shall be effected by levy and salt after ten days advertising, as required by law upon the delivery of such certificate to any officer authorized by law to make levy and sale under execution. PROVIDED However, that either the State Entomologist or the County inspector may, where cause exists outside of any pest inspection district as above provided, pursue the same remedies in all respects as to such lands and owners of

lands outside of any such district the same as though embraced within a pest inspection district."

SECTION 7, Each and every violation of any of the provisions of this Act, or non-compliance with any motice or direction given by the inspector in charge under the provisions of this Act, shall render the offender liable in an action of debt for not less than Twenty-five dollars (\$25.00) and not more than Three hundred dollars (\$300.00) and each and every day that any person, firm or corporation shall fail to comply with any notice in writing received from the inspector in charge under the provisions of this Act, shall be deemed a separate offense. If the land in question be within or without a pest inspection district and the defendent be found guilty, the Court shall also render judgement against the defendent for the cost and expense."

An Act to Prevent the Introduction, Importation, or Spread of Injurious Insects and Plant Diseases in Golorado; to Provide for their Extermination when found in the State; to Require the Inspection of Nurseries, Nursery stock, and Orchards; to Make an appropriation for the Carrying out of this Act; To Regulate the sale of Insecticides, and to Repeal Chapter Fifty-five (55) for the Session Laws of 1897, and providing a Quarantine

for Nursery Stock, Potatoes, and Other Objects
Infected with Insect pests or Plant Disease. Be it
enacted by the General Assembly of the State of
Colorado.

"SECTION 1. For the purpose of this Act, the Entomologist of the State Agricultural College is hereby declared to be State Entomologist, whose duty shall be
directly or thru his deputies or Horticultural Inspectors, to carry out the provisions of this Act. The
compenstaion of the State Entomologist shall be ----- etc."

"SECTION 4. It shall be the duty of the State
Entomologist, directly or thru his deputies or Horticultural
Inspectors, to inspect all the nurseries, orchards, and
shade trees of the State each year, and when it seems
necessary, to inspect potatoes, and other farm or garden
crops or other articles whatsoever, for the purpose
of determining whether or not any of them are infested with
injurious insects of fungous or other plant diseases that
are likely to be disseminated to the injury of others, and to
determine the best methods of control for the pests and
diseases with which he has to deal.

When a nursery, farm or otherpremises that has been inspected seems to be free from insect pests and plant diseases, the Entomologist shall, upon request,

issue to the owner, renter or occupant, a certificate stating these facts. But if any nursery stoc, orchard or shade trees, potatoes, or other farm or garden crops or any objects whatsoever, be found to be infested with any serious insect pest or fungous or other plant disease that is likely to be disseminated to the injury of others, the owner or renter or person in charge of the infested or diseased property or premises must disinfect or destroy the same under the direction and to the satisfaction of the State Entomologist, or his deputy or Horticultural Inspector in Charge, and at the owner's expense, unless the State Entomologist or Inspector in charge sees fit to remove, disinfect or destroy the infested or diseased articles himself:"

"SECTION 6	
ورن ورن من جن المن المن ورن ورن ورن المن المن المن المن المن المن المن الم	

Each and every violation of any of the provisions of this Act, and each and every non-compliance with any notice or direction given by the Horticultural Inspector in charge under the provisions of this Act, shall be punishable by a fine in the sum of not less than five dollars nor more than five hundred dollars, and each and every day that any person, form or corporation shall

fail to comply with any notice in writing received from the State Entomologist, or Horticultural Inspector, in charge under the provisions of this Act, shall be deemed a separate offense."

"SECTION	8	

Whenever the State Entomologist, his deputy or the Horticultural Inspector in charge, in their investigations shall find a section or any portion of the State to be infected with a serious pest or plant disease, and in their judgement, the shipment of potatoes or other farm seeds, products or other articles from that district or section should be liable to spread the insect or disease into other sections, they, or any any of them, may declare a quarantine against such section to prevent the shipment of potatoes or other farm products or seeds or other articles, from the quarantine sections; also, whenever it is ascertained that infested potatoes of other farm seeds, products or other articles are being imported from another state or states, the State Entomologist may declare a quarantine against the importation of such infested potatoes or other farm seeds, products or other articles from such state or states."

STATE ENTOMOLOGIST QUARANTINE NO. 8.
On Account yf Black Stem Rust.

The fact has been determined and notice is hereby given, that the common barberry (Berberis vulgaris) and its horticultural varieties, as well as other species of berberis and Mahonia, are capable of harboring the Black stem rust of wheat, oats, barley, rye and many wild and cultivated grasses. Thru co-operation with the Department of Agricultura, Colorado Agricultural College, local organizations, and individuals, susceptible species of barberry have recently been largely eradicated from the State of Colorado.

NOW, THEREFORE, I, C. P. Gillette, State Entomologist, of the State of Colorado, by virtue of the authority vested in me by the Amended Horticultural Inspection Act, Chapter 131 of the Session Laws of 1917, and for the protection of the great food producing cereal crops of the State do hereby establish a quarantine forbidding of all plants of the following species, into the State of Colorado, or outside the State of Colorado.

Berberis aethenis, B. altiaca, B. amurensis, B. aristata, B. asiatica, B. atropurpurea, B. brachybotrys, B. brevipaniculata, B. buxifolia, B. canadensis,

B. caroliniana (carolina), B. coriaria, B. cretica, B. declinatum, B. fendleri, B. fischeri, B. fremontii, B. heterapoda, B. ilicifolia. B. integerrima, B. laciflora, B. lycium, B. macrophylla, B. neapalensis, B. neubertii, B. umbellata, B. vulgaris, including its subspecies and horticultural varieties, Mahonia aquifolia, M. diversifolia, M. glauca and M. repens.

Therefore, by authority vested in me by the Amended Horticultural Inspection Act, Chapter 131, of the Session Laws of Colorado for 1917, I also declare all the names species to be pests, harboring a serious disease to grains and grasses, and order their extermination wherever they occur in the State of Colorado.

This order supersedes Quarantine Order No. 6, issued July 20, 1918.

This order shall be in effect on and after July 25, 1919; and until further notice.

C. P. Gillette,

State Entomologist,
Fort Collins, Colorado.

Fort Collins, Colorado, July 19, 1919.

RESOLUTION ON BARBERRY BY COLORADO COUNCIL OF DEFENSE.

"Whereas, it has been shown by the most accurate scientific work that the black stem rust of wheat and other cereals and grasses develops on both the green and purple varieties of common barberry (Berberis vulagris) bushes in the spring, producing countless numbers of minute spores which may be carried long distances by the wind to wheat, other cereals and grasses, and whereas the chief wheat producing states, under the encouragement of the U.S. Department of Agriculture, are engaged in the eradication of common barberry in order to increase food production; and whereas, it has been shown in this and certain European countries that black stem rust has disappeared gradually and contemporaneously with the barberry bush; and, whereas, we have in the Japanese barberry (Berberis thunbergii) a variety which is immune to rust and may be planted with safety; therefore, be it

RESOLVED, That we, the State Council of Defense, request that there be no further planting of any barberry (berberis spp.) bushes except the species and variety known as Japanese barberry (Berberis thunbergii), and that all barberry bushes except the immune variety above mentioned be destroyed on all premises within the

bounds of the State of Colorado. Further, we declare the barberry varieties which support black stem rust a pest, and we support the State Entomologist in his efforts to enforce the provisions of the Colorado Pest Inspection ACT ."

STATE FINANCIAL AID

No direct financial aid has been given by the State to the Barberry Eradication Campaign in Colorado. However, the Department of Botany of the Colorado Agricultural College has furnished office space for those carrying on the work and has permitted them the use of the Department's furniture, sotrage space, general laboratory and office equipment and conveniences and considerable amounts of various office supplies.

The use of the above mentioned things has saved the Campaign approximately three hundred dollars per year.

In addition to the financial aid those connected with the Department have given their hearty support to the Eradication owrk and have given much assistance in the form of advice and suggestions. Their co-operation has been much appreciated.

PUBLICITY

As stated in the introduction to this report the greatest efforts in the way of publicity were put forth during the early years of the campaign. Less was done each succeeding year in this connection as the continued advance of the work furnished its own advertising. Bulletins and descriptive circulars and eards are always carried by the scouts and a large number distributed in this manner. These and the articles printed in the different papers constitute the publicity agents being used in the state at this stage of the campaign.

The following list furnishes more definite date upon the nature and extent of the work done in getting the campaign before the people.

BULLETINS AND CIRCULARS.

Three extension bulletins have been published sinde the beginning of the campaign. They are:

"Eradicate the Common Barberry." Sept. 1918, Extension Bulletin Series 1, No. 147, 6000

"Destroy the Common Barberry" Feb. 1919, Extension Bulletin Series 1, No. 155-A 10,000

"Save the Wheat." June 1931, Extension Bulletin Series 1, No. 176-A, 8000.

The following bulletins and advertising matter were also distributed.


Farmers Bulletin No. 1058	6,000
Reprint (Year Book 1918) No. 796	5,000
Sheets of Story: "Banish the Barberry	3,000
Colored plate showing life cycle spores, etc.,	1,000
Life cycle cards	3,000
large colored posters	9,000
Copies of State Entomologist Quarantine No. 8	1,000
Form letters: A. mailed to every property owner on whose property common barberr had been found B. Mailed to all schools, rural news papers, and farmers	y 20,000
Arbor day poem, describing common barberry	1,000

NEWS PAPER PUBLICITY.

papers, they for the most part being eager to assist the campaign by the donation of whatever space was desired. About 500 articles have been published many with descriptibe illustrations or cartoons. These have appeared in every paper in the state and thus have reached all the news paper reading inhabitants.

Some of the leading stories and their authors were:
"A Fungous Parasite and its Host."

E.L. Sargent.



etc.

EDUCATIONAL EXHIBITS.

Many window displays have been placed before the curious eyes of the public in the course of their education. These have varied somewhat in design

but all have been planned with the same purpose in view; namely that of conveying to the people the nature of the common barberry, its characteristics, the object of the campaign, and thus thru their acquired knowledge to secure their co-operation in eradicating the bush. The usual exhibit consisted of bushes of both common and Japanese barberry bushes, samples of rusted grains, and different articles to attract attention, such as sacks of flour, bundles of grain, loaves of bread, and so on. Numerous placards were used to designate the different articles and also show in detail the essential facts concerning the rust and part played by the barberry. impressive statistics were printed on attractive cards and also worked into the display. Thes e exhibits were placed in every town of consequence and in the larger cities several were shown at the same time.

Similar exhibits were placed in the Post Offices of the larger cities where they were subjected to the inspection of many different classes of people.

Likewise the City Parks were used to advantage, large common barberry bushes being placed at the entrances with this sign in close proximity:

DESTROY THIS BUSH
THE COMMON BARBERRY SPREADS
STEM RUST OF WHEAT
DESTROY YOUR BUSH
U. S. DEPARTMENT OF AGRICULTURE.

The City Libraries were utilized in the effort to place before the public the object and necessity of the campgaing. These displays usually consisted of a placarded bush or possibly two, one of each variety with plenty of descriptive literature near by for free distribution.

The following warning was also placed in the street cars of Denver.

"Have you destroyed your Common Barberry Bushes, Government Inspectors here again. "

Complete and striking demonstrations were maintained at all the leading fairs over the state and at the National Stock Show, hold in Denver. Special effort was made to make this last exhibit complete in every detail, because of the great number of people who would be attracted to it.

Bublicity Thru the Theater.

The motion picture houses over the state provided an effective means of advertising the campaign and getting the facts before the people. All the important play houses inDenver and the other cities of the state ran an educational slide for as much as a week at a time. The return of the scouts was also often heralded in this manner and at the same time the people's minds freshened concerning the campaign.

Illustrated lectures were given in the schools and at

farmers meetings over the state. In order to cover the field, several sets were put in use at the same time.

In this connection the film "Barbarous Barberry" was used to advantage and was enthusiastically received wherever shown.

The following slides were used:

"Do you know that a fungus parasite, stem rust, destroyed more American wheat in 1916 than 200 Coloradoans could eat in a year, that common barberry, an ornamental shrub spreads this disease?"

Pictures of common barberry with the words: "the common barberry. Triple spines, sawthothed leaf edges, berberis in clusters. Spreads wheat rust, destroy this bush."

Picture of Japanese barberry and these words: "the Japanese barberry. Single spines, single berries, smooth leaf edges. Does not spread rust, save this bush."

Picture of red rust of wheat and the words: "Stem rust of wheat. Red rust, summer stage. The barberry spreads this stage to the wheat in the spring."

Picture of black stem rust of wheat and the words: "Stem rust of wheat. Black rust, autumn stage. This stage can only infect the barberry. It can not infect wheat."

Life cycle of stem rust. Picture of red rust with label. Picture of black rust with label. Picture of barberry rust with label. Wording "Life cycle of stem rust. Remove the Barberry bush and destroy this cycle."

Colored slide made from Barberry poster.

Slide made from back of stationary.

"Picture of girls with skirts and hats blowing and words: "Colorado winds blow dust across the state. Rust spores are lighter than dust particles. They travel hundreds of miles."

Picture of common barberry leaves, stems and spines and words: "Common barberry. Destroy this bush. Spines mostly 3 in a group. At tip of stem sometimes single. At base of stem sometimes in fours.

Picture of Japanese barberry leaves, stems and spines and words: "Japanese Barberry, save this bush. Spines mostly single. Seldom in two's or three's. Leaves smooth.

Picture of Japanese barberry berries and words: "Japanese barberry, save this bush. Berries born singly or in groups of Two's or three's."

Pictures of common barberries and the words: "The common barberry, destroy this bush. Berries in clusters dull red, shriveled.

"Have you destroyed your common barberry bush?"
Government Inspectors are here again to save the wheat
from rust."

George's cartoon: "The Hun of the Grain World."

Cities and Towns Completed in Survey
From Beginning of Campaign to December 31, 1921
(Very small places omitted.)

Adams County
Adams City, Brighton, Bennett, Darlow, Derby, Eastlake,
Hazeltine, Henderson, Sand Creek, Junction, Scranton,
Strasburg, Westminister.

Alamaso County
Alamosa, Estrella, Hays, Hooper, Henry, McGinty,
Mosca, Willis.

Arapahoe County
Deertrail, Englewood, Littleton, Petersburg, Sullivan

Archuleta County
Arboles, Blanco, Chromo, Pagosa Junction, Pagosa Spgs.

Bacca County
Springfield, Stonington, Two Buttes.

Boulder County
Boulder, Broomfield, Erie, Eversman, Hygiene, La-Gayette, Longmont, Louisville, Lyons, Niwot.

Ohaffee County
Buena Vista, Granite, Poncha, Salida.

Cheyenne County.

Cheyenne Wells, Kit Carson, Wild Horse.

Clear Cræek County Dumont, Georgetown, Idaho Springs, Silver Plume

Conejos County
Antonito, Conejos, La Jara, Manassa, Ortiz, Romeo, Sanford,

Costillo County
Blanca, Costillo, San Luis, San Pablo.

Crowley County,
Crowley, King Center, Numa, Olney Springs, Ordway,
Sugar City.

Custer County, None

Delta County
Austin, Decaredge, Delta, Eckert, Hotchkiss, La
Zear, Paonia.

Denver County
Denver

Dolores County Rico

Douglas County
Castle Rock, Larkspur, Sedalia.

Eagle County
Eagle, Gypsum, Red Cliff

Elbert County
Elbert, Elizabeth, Kiowa.

El Paso County Colorado Springs, Fountain, Manitou, Monument, Palmer Lake

Fremont County
Brwester, Canon City, Concrete, Florence, Howard,
Penrose, Portland.

Garfield County
Carbondale, Glenwood Springs, Grand Valley, New
Castle, Rifle, Silt.

Gilpin County
Blackhawk, Central City, Gilpin.

Grand County Granby, Kremmling, Sulphur Springs.

Gunnison County Crested Butte, Gunnison, Irwin.

Hinsdale County None

Huerfano County La Veta, Ravenwood, Rouse, Strong, Walsenburg

Jackson County None Jefferson County
Arvada, Edgewater, Colden, Morrison, Wheatridge,
Willow Springs.

Kiowa County
Aroington, Brandon, Chivington, Eads, Galatea,
Haswell, Sheridan Lake, Towner.

Kit Carson County
Burlington, Flagler, Seibert, Stratton.

Lake County
Leadville, Malta, Birdseye

La Plata County Durango

Larimer County
Berthoud, Ft. Collins, Loveland, Timnath, Wellington.

Las Animas County
Borwind, Earl, El Moro, Forbes, Hastings, Hoene,
Ludlow.

Lincol n County
Arriba, Bovina, Boyero, Genoa, Hugo, Limon.

Logan County
Atwood, Crook, Flemming, Padroni, Peetz, Rrocter,
Palisades, Whitewater.

Mineral County Creede, Wagonwheele Gap.

Moffat County Craig, Juniper, Great Divide.

Montezuma County
Cortez, Dolores, Mancos

Montrose County
Cimmaron, Montrose, Olathe

Morgan County
Brush, Fort Morgan, Orchard, Snyder, Wiggens.

Otero County
Fowler, La Junta, Manzanola, Rocky Ford, Swink.

Ouray County Colona, Ouray, Ridgeway.

Park County None

Philips County
Amherst, Haxtun, Holyoke, Paoli

Pitkin County
Aspen, Norris.

Prowers, County
Briston, Granada, Hartman, Holly, Lamar, Wiley

Pueblo, County
Avondale, Baxter, Boone, Lima, Pueblo, Rye, Siloam.

Rio Blanco County None

Rio Grand County
Del Norte, Monte Vista, South Fork.

Routt County
Hayden, Phippsburg, Steamboat Springs, Yampa.

Saguache, County Grestone, La Garita, Mirage, Saguache, Villa Grove.

San Juan County
Howardsville, Silverton

San Miguel County
Placerville, Telluride

Sedgewick County
Julesburg, Ovid, Sedgwick.

Summitt, County
None

Teller County
Cripple Creek, Divide, Florissant, Goldfield,
Independence, Woodland Park, Victor.

Washington County Arkon, Otis

Weld County

Ault, Briggsdale, Eaton, Erie, Evans, Fort Lupton, Greeley, Hudson, Johnston, Kersey, Lucern, Mead, Miliken, Nunn, Platteville, Pierce, Severence.

Yuma County Eckley, Laird, Wray, Yuma

Counties Covered in Farm-to-Farm Survey From Beginning of Campaign to December 31, 1921

Name of *	1 5	1	No or	No. mi.	No. of	No.
Name Or	Area of		No. sq.	travel-		Bushes.
County	County	Survey-	miles	1	ings	Dasies.
`	7.47.0	ed	Surveyed 789	ed 710	10	31
Adams	1410	5 6		616	2	7
Alamosa	700	80	560)	34	549
Arapahoe	890	77	685	612	34	04 5
Archuleta	1415	30	437	488		
Baca	2530	15	380	364		
Bent	1345	.30	. 469	605 730	163	1911
Boulder	785	58	455 97	110	4	5
Chaffee	97.5	10 66	1235	1164	4	J
Cheyenne	1850 410	50	205	194	4	42
Clear Creek	1215	15	183	300	6	29
Conejos Costillo	1260	16	198	210		20
Crowley	800	65	524	351	2	10
Custer	750	00	J254	001		
Delta	1215	25	303	366	8 :	27
Denver	65	100	65	1400	669	9261
Dolores	1095	20	219	195		
Douglas	660	20	602	410		
Eagle	1725	1 5	258	214	2	2
Elbert	1625	40	730	700		
El Paso	2140	69	1486	1648	230	3912
Fremont	1600	25	400	523	67	2150
Garfield	2700	20	540	481	3	6
Gilpin	175	25	44	76		
Grand	1925	20	385	295		
Gunnison	3275	10	328	368	-	
Hinsdale	1075				·	
Huerfano	1530	15	225	240	3	3
Jackson	1660					
Jefferson	770	55	423	900	13	115
Kiowa	1690	40	676	586		
Kit Carson	2210	40	884	751	·	
Lake	400	60	240	312		
La Plata	1680	25	586	625	5	43
Larimer	2675	55	1471	1974	11 6	1195
Las Animas	4790	12	572	673	4	6
Lincoln	2640	45	1188	972	1 1	3
Logan	1865	90	1678	1782	4	12
Mesa	3420	30	1026	1263	10	169
Mineral	900	25	225	200	ļ	
Moffat Montezuma	4810 2215	3 25	144	112 594	2	3
MOII PASaula	DALU	80	004	054	1 5	၁
		•				
			1	*		

Montrose Morgan Otero Ouray Park Philips Pitkin Prowers Pueblo Rio Blanco Rio Grand Moutt Saguache San Juan San Miguel Sedgwick Summit Teller Washington Weld	2310 1335 1260 565 2220 690 980 1360 2435 3200 865 2415 3255 435 1350 549 650 549 590 24185	20. 100 35 27 100 18 37 60 50 5 20 100 5 60 50 60 50 60 50 60 60 60 60 60 60 60 60 60 60 60	462 1335 441 153 690 176 422 1455 432 121 423 152 270 549 29 1560 3347	511 1721 594 195 570 195 555 1895 304 211 485 285 470 55 1368 3990	7 12 11 6 1 3 63 9 1 115	50 151 99 10 1 1433 24 2 2 1361
Yuma Totals 1	2420 04834 q• mi	30	726 34194 sq• mi	586 38307 miles	6 1596 Props	22548 bushes

(No data on cost per county. Therefore could not give cost per mile per county).

LEGEND

Areas Covered in Farm-to-Farm Survey.

Beginning of campaign to June 30, 1919

July 1 to December 31, 1919


January 1 to June 30, 1920

July 1 to December 31, 1920

January 1 to June 30, 1921

July 1 to December 31, 1921

All work done of Federal Funds.



LEGEND

Areas Resurveyed


July 1 to December 31, 1919

January 1 to June 30, 1920

July 1 to December 31, 1920

January 1 to June 30, 1921

July 1 to December 31, 1921



		Campaign	Entire			Since Jul. 1				Jan. 1	Since					Month	This		terroc	
:Total : 59	:Removals:		: :Original: 59	:Total : 16	Removals:	: Uriginal: 10		Total 37	:Removals:	••	Original: 27	:Total :		:Removals:	:Resurvey:		:Original: (:Surveyed:	: Survey : No. or :	· ··
				218:	323	100		43 5	73:		362:			,,	••		STATE	days	mem .	
: 1499			1499	7	.,		J	31 :			3 - -						tota	1	H: prof. T.	Nu
19		,	19 :	H				છ			ω		••				2/55. *	l ro	:Having :To	Number of
97 :			97 :	24			ีย 4	<u>ဌာ</u> လ			58			٠.		;	RealMy		tal:	roper
1596:			1596: 1	31				83;			83:	.	**			· ;	MA C	Found: Removed:	bushes	d T
1593 ::	69	· ••	1524 ::	28	80		 S	85	16	•••	69	::	::	:;	•••	.:	1 (a.s.) 10	moved::	were::	: :
19172:		• ••	19178:	37		0	1 1 1	607 :		••	607 :		••		`);	halin.	,	and:	C4#46.
1411:			1411:	<u>ا</u>			ر.	14 :			14 :						7 6 6		:Escaped: T	Number
3376:		• •,	3376:	213			2 .	299 :		••	299			••	••	1/2 /	EC 3/		otal	หารทย
22548: 22457		*	22548:	250			ប ភ	906 :		•-	906			• •		-	1921	Found:	Country	in nt
28457 :	1889 ;		20568	159	35:	F 5	7 7 1	1212	583	••	629		• •					Found : Removed : Survey	GI JU	22
3226	3226			289	289			630	630									Survey	on Re-	e mords:


and the Contract of the Contra

Part of the of			The second secon	,									
Fig.											-		
Fig. of Chils. Statisty Data Da					-								
### CHANGE OF CHANGE CHANGE						C				5			Yuma
Fig.	948	1861	345 :	45	910	0	- -	26	2	89 -	300	con :	Weld
	7	7									G 1		Teller
The control of Properties Hother or election of the North of Properties Hother or election Hot	2	S					٠				100 :	K .	Sedgyic
Column C											25 	ie]	San Mig
Sec. Cont.			•								13)e	Saguaci
### Control State Control State Control Contro											1 (SOUT	Rio Bla
### 16 of Colors Co	34	24 :			44	0 03	o 53			63	50 0	2	Pueblo
	0	9				3	3 :			3	37 :		Prower
### Character Country Character Char				**************************************	1 C	- 0:	- 0.		**************************************	- 0:	100:	6	Philli Patkin
)			•									Park
	72				98					10	275		Ouray
Notice N	151	: 191			151		123 :			18:	100:	000	Morgan
Number of Power Number of Power Number of Register Number of Register Number of Register	50	50		•••	υ υ υ υ	3 00				3 03	25	uma.	Montez
											300		Moffat
Number of Properties In	1.69	169			1.69	10	10			10	30:		Mesa
Control Country Coun	12	12			12		4			4	90:	ŭ	rogan
Co of Sites Number of Fungations Number of Husbes in the country Section Sites Section Sites Section Sites Section	30	.a 02			3 6	4	4			4.	ري دا د	Da Ta	Las Ani
Interest Interest Country Co	838	1195	776	122	: 419	109 :		28	7	& C	 ວາ ວາ	5 00	Larima
No. of Sho. of Short Strikes in Country Every Short Strikes Every Short Strike	43	43			A	n	n	, ,			. 60		Lake
Number of Bucker in											400	7000	K1 Cwa
Number of Bushes in Number of Subshes in Number of S	1.62	165	5 8	٦	113	12	: 13 :	20			55	BOD	Jefferson
Number of Properties in Number of Bushes in Number of Bushes in Number of Bushes in Number of Bushes in Number of States Number of Bushes in Number of Bushes in Number of States Numbe				ada a gira da		•	. 0			3	נו נו		uerfan Huerfan
Number of Properties in	3	3			3	7	1					P	Hinsdale
Number of Properties in Number of Bushes in Country Each of its in Country Each of its in Country Each of its in Each of its											: 20 ::00		Grand
Number of Properties in		The second secon			A contract of the state of the		•				: 25 :		
Number of Properties in	σ	0	- H	1001		300		24	7	33	 ວັນ	الم	Fremont
Number of Preparties In Number of Bushes in Hothers Man Starting Total Starting Total Starting Total Starting Total Total Found Total Total Found Total	3872 1841	3912	. 1087	ייני מייני	$ \omega $	229				230	: 69 :	erin universitä paragotti etti mattaatika	El Paso
Number of Properties in	8	50			20	2	03			N	: 15		Hag le
Number of Proporties in											: 20 : 70		D olore s Douglas
Number of Properties in	8711	9261	•		: 9261	658	688:			: 669	: 100		Denver
Number of Properties in Number of Bushes	27	: 27			: 27	77	: 7		And the second s	. 7	 25	Territorio del Primero de servir de selecto del se	Delta
Number of Properties in						ເນ	<i>∞</i>	•	*	<i>δ</i> υ	တ	7	Crowle
Number of Proporties in Number of Bushes in the country South of thes Country South of the country South of					20		c			C	9 F	e	Costil
Number of Properties in	C4 C2 C3 C3 C3 C3 C4	44 C				40	46			40	50	iro i	Clear Cre
Number of Properties in Number of Bushes in	D	0		-		4	1	• • •	,	4	00 :		Chaffe
Number of Properties in Number of Bushes in Country Both, on which Cities Country Both cities Country	1496	1911	103		80	S	0	: 13		150	: 58		Boulder
Number of Properties in Number of Bushes in to of :No. of :Country : Both, on which :Cities : Country : Both, on which :Cities : Country : Both cities : Country : Count											255 215		Ba.ca
Number of Properties in	000	D#8		2	: 437	: 33	: 34	 လ	20	32	: 77	i do	Arapabo
Number of Properties in Number of Bushes in Country Both, on which Cities Country Both, on which Cities Country Both cities Country Both cities Country Both cities Country Both cities and Escaped Total Found Found Regiments Found Foun	7				7	20 0	100 C			23	80		Alamosa
of :No. of:Cities: Country :Both, on which :Cities: Country :Both cities : Country : Both cities : Country : Coun	251	3	4		27	»	0	. · ·		0	veyed		Adama
of :No. of:Cities : Country : Both, on which : Cities : Country : Both, on which : Cities : Country : Both cities : Found : Removed : Towns : Found :	1	Found	foral		s s	shes were Removed:	Found:	rtry : Total	@ 05 (C)	Cities and Towns	~	cally	Countie: phabetic
Number of Properties in Number of Bushes in No. of Cities : Country :Both, on which : Cities : Country :Both cities : Country :Both cities : Country :Both cities : Country :Both cities : Country : Both cities : Country : Country : Both cities : Country : Cou	. i	bushes in	mber of	Nu	Y COUNT	LY REPORT	ur set	propert	Number cr		et	>	
: Number of Properties in : Number of Bushes in : No. of:Cities : Country : Both, on which : Cities : Country : Both cities : S. Man : and : Having : Total : bushes were : and : Escaped : Total : Found : Removed : Towns : Escaped : Total : Escaped : Total : Escaped : Total : Escaped : Towns : Escaped : Total : Escaped : To	ŀ	22548	į	141	19172	48	:1596		1	1499			Camp'gn
Number of Properties in Number of Bushes in Stand of Cities: Country Both, on which Cities: Country Both cities Country Both cities: Country Both cities Stand Having Fotal Bushes were and Escaped Fotal Found deays: Towns: Escaped Found Regimning of Amipaign to Dec. 31/92/ Original Survey From Acgimning of Amipaign to Dec. 31/92/ 368 31 9 58 83 69 607 14 299 906	1		D + C				: 31	24	, · ·		86		Jul. 1
Number of Properties in Number of Bushes in No. of:Cities: Country :Both, on which: Cities: Country :Both cities: No. of:Cities: Country :Both cities: Country :Both cities: Man: and :Having: Total: bushes were: and :Escaped: Total: Found: di days: Towns: Escapes: Found: Removed: Towns: Criginal Survey From Acquaning of Carupaign to Dec. 31/92/	7 0 L	277	7 77	ا ر	3			00	ď	C	: 004		Since
: Number of Properties in : Number of Bushes in : No. of:Cities : Country :Both, on which :Cities : Country :Both cities : S: Man : and :Having : Total : bushes were : and : Escaped : Total : Found :d: days : Towns : : : : : : : : : : : : : : : : : : :	<u>ෙ</u>		299	14	607	So ,	7.8	o į	4	7	, J		Since
: Number of Properties in : Number of Bushes in :No. of:Cities : Country :Both, on which :Cities : Country :Both cities : Having : Total : bushes were : and :Escaped : Total : Found :d: days : Towns :Escapes : Found :Removed :: Towns : :			31.19	to De	10/1/0	0066	:	OM Ro		12/184	2.		This Month
: Number of Properties in :: Number of Bushes in :No. of:Cities : Country :Both cities : Country :Both cities : Found :				ì	Towns	Removed	Found	. c	Escape	Towns	days	Surveyed	
: Number of Properties in :: Number of Bushes in		12	T _o		Cities	on which:	:Both,	गुत्र <u>भ</u>	Count	Cities	:No. of	'Counties	Period
			25	Numbe			s in	pertie	ber of Pi	Nun			

										υπ co		Weld Yum
3											Summit Summit Teller Washington	Summ Tell
								••	0	រាបា្	Juan Miguel	San San
											Grand t:	Rio
: <u>194</u>												Pitk Pueb
ω.											1 00 · · ·	Park Phil
27 : 1					4						ro ro v:	Mor Ote
: 17										.	tezuma trose	TOM
											+ 12	Mes
											Animas	Las
357 : 202		• • • • •			7	.)lata mer	
											Garson	7 K K K K K K K K K K K K K K K K K K K
											reon	Jeffe
										 B	iale	Hing
										 I טון	ion ion	taurin Sasan Sasan Sasan
										 වා හ	#1d	Frenc Carfi
250 :288										: 35 5	Tr Da	Elber Egiper
										5 O 8	1. 10 B	Douglas Earle:
509 : 889					10					100	Fs	Denve
υπ (C						-) I O	P. O.V	Custe
										 5	0:3 0:3	Conej Costi
							••			: 19	nne Creek	Cheye:
388 : 450 :					26				. . .	 68 0	31°	Bould Chaff
										30	l:eta !	Archu Baca
10 : 411										: 70 40	15 6	Alamosa Arapahoe
Removed : on Re- : survey	Found :	tal	Country Escaped : To	Cities: and: Towns:	Both, on which: bushes were: Found:Removed:	Both, c	Country ng :Total pes:	:Havı	:Citie; y: and :Towns	county: sur- veyed	s Al- cally	Counties Alphabetically
:Sproute		of	IES Nu	BY COUNT		MONTHI	of Prop	na pe		: Anı' t		
.					69				••••			Entire Campign
583 : 630 35 : 289			e procedende er en		8 31	-				73		Jan. 1 : Since :
	3/1921	toDes	gaign	e Carri	\square	n Beyr		miney	Resi	2		Month
lemoved :on he-	Found : h	fotal	Escaped : 7	and :	: Duskes were: :Found:Removed::		g:Total	and :Having :Tota Towns:L'Scapes:	: and : Town	s: man d: days	:Counties: :Sur veyed:	
Both cities and country Found	Bushes in :Both cities	rof	Number Country	Cities		Both,	ountry	S: C		ົດ	No. of	Period:

LEGENDS

- Area of escaped bushes found Jan. 1 to June 30, 1919.
- Area of escaped bushes found July 1, to December 31, 1919.
- Area of escaped bushes found Januarl 1 to June 30, 1920
- Area of escaped bushes found July 1 to December 31, 1920
- Area of escaped bushes found Jan. 1, to June 30, 1921
- Area of escaped bushes found July 1 to Decemmber 31, 1921



LIST OF PROPERTIES WHERE ESCAPED BUSHES WERE FOUND

Name	Town	Ģ ounty	Bushes
.United Oil Co.	Canon City Arvada	Fremont Jefferson	1 1
Greeley-Loveland Irrigation Co.	Loveland	Larimer	2
Big Thompson River	Loveland	Larimer	1
Home Supply Ditch Company	Loveland	Larimer	5
Home Supply Ditch Company	Loveland	Larimer	31
Home Supply Ditch Company School house	Loveland	Larimer	39
(rural) Mrs. M. A. Alps Glen Kester Mr. Buckingham C. C. Richardson J. C. Ruple Loyd Jones Poudre River Poudre River Coy Ranch Poudre River	Loveland Loveland	Larimer	40 15 2 7 7 150 50 15 71 1
Vacant Lot Park Lyons & Johnson	Greeley Greeley Littleton Englewood	Weld Weld Arapahoe Arapahoe	1 4 1 1

(Bushes located near town named)

Average Cost by years.

Year	Miles Travel-	Cost per running mile	Cost per sq. mi.	Cost per Planting	Cost per bush
1918	ed 2702	\$.314	\$ • 457	\$5.38	\$.486
1919	17603	•375	•495	4.80	. 357
1920	11705	.177	.187	83.00	1.05
1921	21556	•226	•21O	58.80	5.386
Campai to	gn	•			
Dec. 3	1, 1921				
	49076	.289	•269°	8.90	•63

General Discussion.

Several methods of scouting have been tried out in Colorado since the beginning of the campaign. Practically all of the work was done on foot the first year, being limited to the towns in the north part of the state. Travel between the towns was by means of the railroads. The cose per square mile amounted to \$.457 . The same system was used the following year of 1919, bicycles, however, being used to advantage and during the latter part of the season a car was pressed into service for a short time. This season found the scouts spreading out into sections where the towns were smaller and further apart. Also some rural work was onde. The cost per square mile amounted to \$.495 1920 witnessed the real advent of the automobile into the field one scout using one car the whole season. In addition another scout was placed in the dities on foot. However, this last scout was used only for a short time. The scout with the car traveled in very favorable sections and proved the value of the automobile in the field the cost per square mmle being \$.187. The following year, two cars were placed in the field but with two men to the car. This proved more costly than where one man was used to the car and would have proven so except for the fact that much of the work was

done in sparsely populated districts where very good time could be made. The cost per square mile for the year came to \$.210.

The cost per bush was of course lowest the first two years of the campaign, since the work was done in the cities and thickly populated regions where most of the bushes of the state were known to be located. The following year saw a material increase in cost per bush as the scouts spread out into the rural sections and a still greater increase the year after that when most of the work was carried on in the dry land sections.

In this report all cost per year is based on the total amount spent that year in all the branches of the work since the records do not indicate how the money was apportioned to the different phases.

Up to date, 446 escaped bushes have been found in the state, most of them being in Larimer county. The rivers and irrigation systems of the regions located close to the mountains seem to have proven themselves the most dangerous agencies for spreading the barberry. Thru this means the seeds would in a comparatively short time be scattered over all the districts by the irrigation waters, while the rivers could carry them for miles. Many escapes were also found in wooded areas, having been carried

there thru the agency of birds. Cattle, horses, and other live stock have also proved themselves to be a means of spreading the bushes since many escaped have been found in pasture which were free from trees.

In making the farm to farm surveys it has been the practice to drive into the farm yard and thoroly inspect all places where barberries might be found. As a usual thing it is advisable to first make known the purpose of the visit. If no bushes are found a bulletin is left with the farmer and no more of his or the scout's time is taken up. Where bushes are located, it is usually necessary to spend some little time in going thru the matter in order that there may be a clear understanding as to the reason for removing the barberries. As a usual thing very little difficulty is experienced in gaining the farmers! interest and co-operation. In the course of the campaign very few people have been encountered who have absolutely refused to have their bushes removed. As the usual thing such parties are city dwellers who think they can go on existing whether the farmer exists or not. In such cases the pressure of the law is called upon to accomplish the desired end. We are thankful that only a few of this type are found over the state. A few of course are essential to normalcy.

Progress of Eradication Campaign

January 1 to December 31, 1922

Introduction.

This report coveres the activities of the barberry eradication campaign in Colorado for the period January 1, 1922 to December 31, 1922.

A slightly greater appropriation was used than during the preceding year, practically the same amount of ground being covered. However, considerable more money was used in administration than in 1921.

Work was done in 45 counties. Original survey work was carried on in 23 of these 45 counties and resurvey in all but two of them.

1453 Bushes were found on 46 properties and
1365 bushes removed from 46 properties, making a
total of 24,001 bushes found during the entire campaign on
1642 properties of which all but 179 have been
removed. One hedge of 175 bushes remains in Jefferson
county, one planting of three bushes is still in
the city of Boulder, and one bush remains on a city
property of Denver, these will be out before spring.

In this report in estimating percent of a county surveyed we consider only that part surveyed which has been actually gone over. Areas too rough and mountainous to be surveyed or of a nature not requiring going over, are not considered as surveyed.

Many counties which appear to be only partially

covered are actually one hundred per cent surveyed so far as the territory requiring inspection is concerned. This explanation is offered beca se of the fact that earlier in the campaign many counties were reported almost entirely covered when the nature of the section would make such practically impossible. Larimer county is a good example of this. Early in the campaign it was reported ninety percent surveyed, which no doubt it was so far as the ground requiring surveying was concerned. However, over half of the county is of a mountainous nature, not requiring going over and offering great obstacles to any attempt at such. This we report as unsurveyed.

PERSONNEL

- Fitzsimmons, J. R. St. Leader Colo. \$150.00 mo. June 15 to September 25.
- Learn, C. D., Collab. Colo. 125 mo. April 1 to Sept. 23.
- Lungren, E. A., Field Asst. Colo. 125 mo. April 1 to Sept. 23
- Thornton, B. J. Field Asst. Colo. 125. mo. July 1 to Nov. 30.
- Miss Downs Stenographer.

State Financial Aid.

1. Cash Appropriations.

No cash appropriations made toward campaign by State of Colorado.

2. Financial aid other than cahs appropriations. Office space, light, heat, stenographic, supplies, etc., estimated at \$350.00

Publicity.

Bulletin and Circulars Distributed.

Name Number of copies

Farmers' Bul. No. 1058 1,000

Circular No. 186 1.000

Reprint (Yr. book 1918) No. 796 .500

Life cycle cards

Circulars

Posters

News Articles

News articles were put in the papers of those towns and regions where barberries had been found or were known to be present. Lengthy articles appearing in Denver Post, Pueblo Chieftan, and other of the larger papers of the Stat.

Educational Features:


Nothing was done along this line beyond the distribution of posters, circulars, bulletins and life cycle cards.


Personal Letters:

Personal letters were sent out to those tenants and land owners where sprouts were expected to give trouble.

Fair demonstrations


No demonstrations of any kind were made during the year. -109-








Heavily Rusted Grain And Grasses, Burlington Colo. 1922



Finding An Abundant of Rust Caused By Infected Barberries
Burlington Colo. 1922.



Barberry Hedge Before Digging; Bentlounty 1822



Escaped Barberry Digging By Hand, Wolhurst Colo. 1922

LEGEND

Areas Covered in farm to farm Survey.

Beginning of Campaign to June 30, 1919

July 1 to December 31, 1919

January 1 to June 30, 1920


July 1 to December 31, 1920

January 1 to June 30, 1921

July 1 to December 31, 1921

January 1 1921 to June 31, 1921
July 1 to December 31, 1922

All work done on Federal Funds.



LEGEND

July 1 to December 31, 1919.

January 1 to June 30, 1920


July 1 to December 31, 1920


January 1 to June 30, 1921

July 1 to December 31, 1921

January 1 to June 30, 1922

July 1 to December 31, 1922





:Original :Resurvey :Resurvey :Resurvey :Resurvey :Resurvey :Resurvey :Resurvey	nal	Surveyed: Surveyed: 33 33 34 14 56	Surveyed: days: Towns: and Surveyed: days: Towns 23 61 27 45 288: 37 14 38 18 14 38 18	Towns: Towns: Tota	HH H		Total 19 19 11 11 11 11 11 11 11 11 11 11 11	Found : Ren Found : Ren 46 4 4 4 4 4 4 4 4 4 4 4 4 4 4 6 6 6 6	bushes were Found : Removed 46 37 46 46 29 24 29 34		ESC	ed :: 44 0	[] [] [] [] [] [] [] [] [] []	Found :Ren Found :Ren 1453 1 1453 1 1379 1	260 260 260 365 30	: Found : on Re- i. Survey : 2458 :2458
Entire : Original Campaign: Resurvey : Removals	nal: vey:	d	Od P	1526		30		25 80 26 44	1561	335	3: 2415		1044:	1379 24001	21828	3167
Total				1586		30 ::	116	1648	1639:	19532	2: 2415		4469	24001	1994 : : 23822:	5684

s and country : Removed		:1260	:1198	31828	mage state out and a state out of the state of the state of the state out	s and country: Removed	. 25	7		: 1547	5	989 30		. 27 8754		83	3850		. The state of the		254	7	43	. 6	086		3 50	: 73			: 29	34			2		933			The state of the s
shes in Both cities a Found	0) 12 / (0	120021/13 1453	1379	34001	\$ 	Both cities: Found:	31	7 28		1919	5	29	10	37 9264		CS	3925 2993	9			429		43	0	ا ا ا	DO I	3 50	151			29 1433	24	e propins de la company de la		23	~	1297		de la companya de la	
r of Bush	2	1901 190	044	4469	i i	Number of oral:	7			106											52			000													345			
Numbe: Country: Escaped		1004	096	8418	S	Count		305			A STATE OF THE STA						2237							Q	energian englis desperage de palemente de la mente englis de la mente englis de la mente englis de la mente en					And the second of the second s							45			
Cities and Towns	,	360	335	19533	BY COUNTIE	Cities and Towns	37	7.438		1813	70	2000	0	27.00		CO	3925	9		12	377	13	43	0 4 0 t	o O O	B O T	3 50	151 98		2	1433	24			03	2	0000 1000			
Both, on which: bushes were: Found:Removed:	77.	29111111 37		1561		Both, on which: buth, on which: characteristics were: Found: Removed:	9	ω α.		140	.	[‡] 0	C3	2		w	234	3		5	Г	3	1 1	9 4	4		200	13		9 [4 8 53	က		**			101 :	***		
Both, o: bus:		77 7 8 8 8 8		: 1648:	MONTHL	:Both, o	. 10			. 168:	7		(Q)	7.48		0	233		••				LO C	.126	7	0	2	2 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7	**	 8 L	: 4 : : 63 :	σ: 					138:	**		
opertie V Total		79	1	-		propert 14ry Total		cc		13		tas este plaking papagan en er est		edit rate, date, dell'editte rate, sant deter		esta establismonte de la calendaria de la c	288	ARRESTOR DE CONTRA LA ARRESTOR LA CONTRA LA CO			CO.	ena aparepapan manaka daga sarah		37		e dem er verb i op 1881 – Alle verdete staat de staat en								e meny and a mentro de minerale in compression in the second seco			30			-
Countributed Count			7 ~	30	G	Number of propert : Country : Having : Total : Escapes :				r-1	The second secon					A season of the	O							12	And the state of t												03			
Cities and Towns		1:11.01.5		153	,	ities and owns	თ		1 1	153		4 CO	Q	7.20		œ	236 43	5		3	7.4	23	വ	© 4	- 4	70	00 2	10		ю Н	63	တ		After any control of the control of			1 00			
No, of Man		05/91	1 7			Am't : of :C :county: :sur- :T	veyed	80.	30	100:	10: 66:		65; 65;	1 85 L	440		0.00 0.00 0.00	,, ,, ,,,	30 .	15:	75	40 :	35	13	000	30	300	35	. 27	100.	100	1 00	00 00 00 00	35 :	100	1 2 2 3	100	•••	*	•
Period : No. of : : Counties:	200	200	5 4			Counties Alphabetically				Bent Boulder		O.K			Dolores	Tagle	Flbert Fl Paso Fremont	Garfield :	Gunni son	Hinsdale :	Jackson Jeffarson	Kiowa Kit Carson	Laka La Olata	Larimer Las Animas	Lingolm:	Mesa Mineral	Montest Montesuma	Morgan Otero				Rio Blanco : Rio Grand :			Sedgwick Strong to	Teller Weahington	Reld Vuma			e de la composition della comp

:Surveyed:	s: man : and :Having d: days : Towns:Escapes	Towns	:Having :Total : :Escapes:	Four:	Found: Removed:	: and : Towns	: Escaped : :	[ota]	: Found	Found : Kemoved :	on Re-
		31100			2			,	10,	7.7	
		120 2 W.	1 / E/ FLO			(1)	Modera	10171	36.01, 192.		
1:48	256				O					105	•• ••
1:40	315				ω				de de des de des des des des des des des	06	
gn:					78					1994	•
				MONT	ILY KEPORT	by COUNTIES					
1. A B A] _	: Am't	C1 + 10	Number of	Properties	in altach	i c	Con	Number of E	Bushes in	U	
phabetically	ζ.	and	Having Total	nd : L	: busnes were:	and:: Towns	יו עט ויי	ta1		Removed	
	veyed										•• •
mosa	06 				4					0	1
Arepanoe Archuleta	100:									10	
±	. O4					,					
lder	. 68				: 27	•				369	• • •
ffee	0			•					And the second s	e, on our activities satisfied debeteen discharge en discharge en entre en entre entre entre entre entre entre	••]••
ar: Oreek	1			*							-
ajos tillo	16			-						and the second s	1
mley ter	0.1								And the state of t	eranden geleichte der eine erweite der mitgen begrechte der gebrechte.	•- ••
	:30			***************************************							
nver lores	300				9					A C	ا٠٠ [٠٠
nglas Gia							•		•••••		•• ••
be rt	1.0			an altra de la companya de la compan						200	•
Paso	. 55 . 25				 					259	• • •
71010	10	•							••	and the state of t	
and	1 1										
เทารอก กลัสลาค											
erfano	13			•				••	.,		
fferson	1 1		Andrews of the second s								
it: Carson							4			œ	
Lake	103							••			
arimer	. 66	•		•	. 7			*		357	507
as. Anımas inclon	0.0									estrupte man integrationalismalismas papelantes est subsential des subsential des subsentials des subsentials	.
ogen ess	:30			•							<u></u>
ineral	o										 .
Morfat Monteruma	30										
ontrose	010	-								Andrews and described the second seco	
tero	: 39				· .					22	.]
ay.	נין ו									- Marian	
1:08	. 44							• ••		egili ingenisian galangan gamaningan sa bapantan gala-ta	.
kin wers	45										
b]'o	: 40 :			••						Tagas et en	
Grand	.45							A CONTRACTOR OF THE CONTRACTOR		et digit i e e estaballis privativas destabas em maran	
; [] ;									Control of the C	ANTONIO CONTRACTOR A SECURITA A S	•
usche Juan	:10										
Mid ह्या हो	01.									Andrews of the party of the continue of the co	.
gariok mi*	c 9										
] ør	2		dan da	der de la company							
norton	0 0 									YOZ	, -
Yuma :	(a)			4	-1					5/4 -	1
	• • • • • • • • • • • • • • • • • • • •			••	•		***************************************				
Andrewsking to the state of the											.
				•	••••		* * **********************************				
				• • • •					A A STATE OF THE PARTY OF THE P		
endometrophete entrement entrement of the control o											
	•			•				Annah Company of the			- **
				The second secon							.
	-	•						***************************************			-
			•••	••	•	•					

-1.83-

on Re-			••••		:Sprouts	ry:Found :on Re- :survey		: 528	the control of the co	: 710			7	2021 :		: 589	080 080 080 080			121		33				(S) (C)	מו							: 703				
Removed	3.3	•		23822	1 1	s and country Removed	31	728	electrical description of the second	1916	42	5 6	0	9263	8	3925	0 3000 0 3000	And the second s	33	254	13	43	9	189		000 L		0	29	 400		co	3	7	0			
Found:	31, 192			34001		Both citie	31	728	Andreas and the state of the st	1919	43	68		9864	8	3935	2993		3	429	12	43 0887	200	169	Ę	טוט וי	00 00 1	0,	29	24		C	3	1397	01			The second secon
Pota]	to Dec.			4469	umber of B	tāl	4	390		106							2876			. 53		0 0 0					F-1							345				
: pedeosa :	; do, 1918	1922		3415	IES	Escaped		: 105									2237					000												45				
and Towns	Colora	Dec.		:19532	BY COUNT	: Citles : and : Towns	227	438		1813	42	55		9264	62	3925	6		23	377	73		300	.: 169 .:		50	00 00 00	0 [29	34			2	952	q			
Found:Removed::	I E	دد	.,	1639	LY REPORT in :	Both, on which: bushes were: Found:Removed:	100	339		167	4	0 0		670 :		336:			8	5		2000	1	40		2 2 0		9	4				-	1.23				
Found:		: : Totals 1918		1648	1 -31	,	000	: 39		168	4,			: 671		: 236	77			18			4	401:		2 2 0			44				4	133	တ			
:Total	Barberry	: County		116	f Prope	mtry :Total		o		72	-						28					42		••	•									36				
:Having :Total : s:E%capes:	of Bar			30	Number o	S: Country :Heving :Total :Escapes:				r-1							О.					61												63				
an : and : ys : Towns:	aults			1526		:Citles : and :Towns	on c	33		153	40	9		671	63	336	43		22	7	23	ın g	4	4 01		25 6	20	Ø,	4 6	တ		.,	4	L 80	co			
3: m 1: day	: Reg			April 4 - 4 - 4 - 4 - 4 - 4 - 4 - 4 - 4 - 4	Am't	Σ₃ +3 τ	>	100		. 58		337	00	100 : 40	. 15	: 45 69	30 :: 25	01:	1 22 1	. 75		10 %		300	3 25	080	35		000		ייי יי		- -	1000	04			
:Counties:	This : Month :	ince :	ince :	ntire:		Counties Al- phacetically	Adams:	Archuleta	Baca : Bent :	Foulder Chaffee	Cheyenne Clear Greek	Costillo	Guster	Denver Dolores	Douglas Eaglei	Elbert El Paso	Fremont Garfield Giapin	Gunison	Huerfano Tackano	Teferson	10ma : 1t Carson	a Plata	Las Animas Lincoln	Logan : Mesa	inerel Offat	Montezuma Montrose	organi oro: Ourav:	Park: Philips	Prowers Prowers	Rio Blanco Rio Grand	Routt: Saguache	San Juan San Miguel	Summit	Teller Weshington Weld:	uma :			


-134**-**

Cities and Towns covered 1922 - Original Survey.

No original survey workdone in towns of Colorado during the year 1922.

LEGEND

- Area of escaped bushes found Jan. 1 to June 30, 1919.
- Area of escaped bushes found July 1, to Dec. 31, 1919.
- Area of escaped bushes found Jan. 1, to June 30, 1920.
- Area of escaped bushes found July 1 to Dec. 31, 1920.
- Area of escaped bushes found Jan. 1 to June 30, 1921
- Area of escaped bushes found July 1 to Dec. 31, 1921
- Area of escaped bushes found January 1 to June 30, 1922
- Area of escaped bushes found July 1 to Dec. 31, 1922



List of Properties where Escaped bushes were found.

Name United Oil Co.	Town Canon City Arvada	County Fremont Jefferson	Bushes 1 1
Greele-Loveland IrrigationCo.	l Loveland	Larimer	2
Big Thompson	HOACTWII	HOT THO I	
River	11	ti	1
Home Supply Ditch Co.	11	It	5
Home Supply	."		J
Ditch Co.	11	II	31
Home Supply Ditch Co.	tf .	11	39
School house			
(rural)	ii.	31	40
Mrs. M.A.Alps	11	it .	15
Glen Kester	11	II .	3
Mr. Buckingham	11	11	7
G. C. Richard-	11	11	7
son	ii	11	150
J. C. Ruple	11	lt	50
Loyd Jones	Ft.Collins	11	15
Poudre River Poudre River	t r • COTTIUS	!!	71
Coy Ranch	11	В	ίŢ
Poudre River	La Porte	11	ì
Vacant lot	100 101 00		
(rural)	Greeley	Weld	4
Park (rural)	II	11	4
	Littleton	Arapahoe	1
Lyosn and			
Johnson	Englewood	Arapaho e	1
Wolhurst			
Farm	Wolhurst	11 .	70
	W.Littleton	ı "	1
M. F. Allen		Boulder	37
G. Osterhout Park	indsor	Weld	11
(rural)	Greeley	Weld	2
McCormack	Greeley	11	2

(Bushes located near towns named)

Resurvey Results 1922

	стт	Υ		1.
County	Props visited	(No. Sprouts		of proper- ties having sprouts
Adams Alamosa Arapahoe Boulder Chaffee Conejos Orowley Delta Denver Eagle El Paso Fremont Garfield Huerfano Jefferson La Plata Larimer Las Animas Logan Mesa Montezuma Montrose Morgan Otero Philips Prowers Pueblo Rio Grand Sedgwick Washington Weld	92304 6276 343315844027206339119	106 148 1 318 300 18 2 86 4 21 1 9 8 1 27 1 27 1 259	148 8 10 3 8 16 33 57 10 18 8 6 16 4 	44 19 50 8 6 9 33 54 40 25 25 30 -14 10 14 11

Resurvey Results

	RU	R AL		l
County		No. Sprouts		% of properties having sprouts
Adams	1			
Alamosa				
Arapahoe	2	12-	10	100
Boulder	13	65	12	54
Chaffee				
Conejos	——			
Orowley				
Delta Denver				
Eagle				
El Paso				
Fremont	24	580	33	26
Garfield				~~~~
Huerfano				
Jefferson	2	34	65	100
La Plata				
Larimer	28	284	24	85 ·
Las Animas				
Logan				
Mesa				
Montezuma				- -
Montrose				·
Morgan				
Otero	1		I	
Philips				
Prowers				
Pueblo			ş	man was base
Rio Grand				tions many
Sedgwiok				
Washington We ld	26	26	58 :	50
14 GTØ	20	ب د	50 .	50
ĺ				

Summary of Barberry Survey By Federal Department, 1922

	100	161100	Mi non	Total C	Cost	Cost	Man
County	Sq.		Mi.per		gost g•mi•		Days:
Adams	1410	run 900	sq.mi.	\$157.85		\$.175	14
Alamosa	490	340	694	62.95		•185	5
Arapahoe	890	1215	1.366	216.15	.243		18
Archuleta	424	325	.766	54.85	.129		5
Bent	1345	1197	•890	168.20	.125		14
	534	1048	1.963		•125 •607		27
Boulder				324.25	•507 •127		
Chaffee	146	105	.719				1
Conejos	486	419	•866	79.79		-190	- 6 5
Costillo	391	318	.813	60.10	•154		_
Crowley	80	53	•662	8.50	.106		. 5
Delta ,	182	130	.714	21.00	.115		2
Denver	65	1076	16.554	360.75	5.550		-30
Dolores	219	183	•336	42.25	.189		4
Douglas	129	183	1.419	19.70	•152		2
Eagle	172	141	•218	24.40	.142		2
El Paso	774	821	1.061	143.25	•185		12
Fremont	352	292	.829	51.90	.147		5
Garfield	270	214	•793	32.10		.150	2
Huerfano	77	65	-844	9.90	.128	3.152	1
Jefferson	577	1573	2.726	228.00	•395	.145	19
Kit Carson	906	758	.837	123.20	•136	.162	11
Lake	100	74	.740	12.95	.129	.175	
La Plata	336	267	.795	38.60	.115	.152	3
Las Animas	575	489	.805	60.10	•105	.123	5
Larimer	1964	3251	1.644	445.40	•226		43
Lincoln	264	212	.821	30.15	.115	.142	3
Logan	373	315	•845	42.30	.1:13	.134	4
Mesa	171	186	1.088	23.05	.135	.124	4
Mineral	45	38	.844	7.05	.156	.201	1 1
Montezuma	317	3.08	.971	40.65	. 128	.132	4
Montrose	115	98	.852	15.50	•135	.158	1.5
Morgan	534	470	•830	58.45	.106	.125	5
Otero	3 1 5	344	1.092	47.35	•150	-138	4
Ouray	85	67	•788	11.35	•133	.169	1
Philips	276	240	.870	33.20	.120		3
Prowers	1360		•952	150.30	.110		15
Pueblo	365	397	1.088	57.45	•158	-145	5
Rio Grand	390		•990	46.15	.119		3
Saguache	488	408	.836	46.20	•095	.113	4
San Miguel	203	168	•828	24.10	•118	144	ĺ
Sedgwick	329	290	.881	34.60	105	.119	3
Teller	89	70	.787	19.40	117	148	
Washington	182	156	. 857	20.30	•111	130	2
Weld	4185	3915	•912	549.25	ъз1	140	46
Yuma	242	190	•785	25.25	.105	.134	
	23222	24990	1.076	4037.25	•103	162	2/10
~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~		~~000	1 = • = • • • • •	1 1001 100	1 • 7 1 4	1.100	349

Summary of Costs - Original Survey

	1	 						1	
	Man	Total		Cost	Cost	Plant	-Cost		Cost
	Days	Cost	Miles	per	per	ings	per	}	per
	:		į	Mile	.Sq•			Bushe	, -
			<u> </u>		mile	3		2	
Adams	5	\$70.10	520	1.133	\$.113	;	<u> </u>		
Alamosa	1 2	12.50	65	.192	.179	i i			
Arapahoe		25.15	200	.122	.123	5	5.03	179	\$.15
Bent	8 .	95.80	742	.129	.109	5	4.25	8	2.65
Chaffee	• 3	6.40	40	.160	.128		- 420	Ü	
Conejos	•5	33.10	267	.124	.109				
Costillo	1.7		172	.124	•113				
Denver	•5		40	.150	•400	2	3.00	3	2.00
Dolores	4	42.25	183	.231	.189	-	3.00	U	2.00
El Paso	•3		23	.152	.140	6	•56	13	•27
Fremont	1.2		80	.187	.87	5	೩.99	893	.02
Jefferson	3	36.50	212	.172	-238		12.17	264	.14
Larimer	3.3	46.10	310	.149	.157	9	5.12	25	1.84
Kit Carson		5.25	43	.122	.240	3	1.75	12	•44
Lincoln	3	30.15	2.2		.115		J. • 1 O	₩	•
Prowers	3 9 1	105.0¢		.128	.112	l \	105.00	20	5.25
Rio Grand		11.50	80	.147	.133				0.50
Saguache	2.5	38.45	269	.149	.119				
San Miguel			121	.133	.119				
Teller	.7		50	.153	.130				
Weld	6.8	94.15	683	.138	.112	7	13.45	36	2.62
<u>Yuma</u>	2 1	25.25	190	.134	.105				2 • 0 2
State									
Totals	61	768.45	5463	.140	.122	46	16.70	1453	•53
							. •		

Summary of Costs Per Acre 1922

Total Costs (Including Administration and Overhead)

Cost per	Cost per	Cost per	Cost per	Cost per bushel small
Acre	A.wheat	A.small	bushel	
\$.000361	\$.00333	grains	whea t \$.000248	grains \$.000149

Field Costs (Including only expenses incurred in field)

Cost per	Cost per	Cost per	Cost per	Cost per
Acre	A.wheat	A. small	bu•	bu. small
		grains	wheat	grains
\$.000272	\$.00249	\$.00193	\$.00018 5	\$.000112

Note: Production statistics per county not yet available

Resurvey Costs.

· · · · · · · · · · · · · · · · · · ·	Man	Total	Total	Total	Cost	Cost
County	12	Props	f	Cost	per	per
					Prop.	Sprout
Adams	9	10		\$ 87.75	\$ 8.78	\$
Alamosa	4	2		50.45	25.22	
Arapahoe	16	34	117	191.00	5.62	1.63
Archuleta	5	ŧ.		54.85		
Bent	6	1	•	72.40		
Boulder	25.5	163	260	303.00	1.86	1.17
Chaffee	7	4		11.75	2.94	
Conejos	3.5	6	İ	46.60	7.77	
Crowley	•5	. 2	1	8.50	4.25	8.50
Delta	2	7		21.00	3.00	
Denver	29.5	669	318	354.75	•53	1.12
Douglas	2	1		19.70		
Eagle	2	2		24.40	12.20	
El Paso	11.7	230		139.75	.61	•45
Fremont	3.8	67	698	36 .9 5	•55	•05
Garfield	2	3 3		32.10	10.70	
Huerfano	1			9.90	3.30	
Jefferson	16	13	120	192.50	14.81	1.60
Kit Carson	10.5	_		117.95	,	
La Plata	3	5	4	38.60	7.72	9.65
Larimer	39.7		305	399.30	3.44	1.31
Las Animas	- 5	4		60.10	15.02	10.70
Logan	4	4	9	42.30	10.57	42.30
Mesa	4 2 1	10	9	23.04	2.30	2.72
Mineral		.0	The state of the s	7.05	00 77	
Montezuma	4	7	8	40.65	20.33	7.04
Montrose	1.5 5	3	0	15.50	2.21	1.94
Morgan		12	4	58.45	4.87	77 04
Otero	4 1	1 11	4	47.35 11.35	4.30	11 .84
Ouray Philips		6		33.20	5.53	Tree trans-
Prowers	5 E	3		45.30	15.10	· ·
Pueblo	3 6 5	63	27	57 • 45 ·	•91	2.13
Rio Grand	2	9	1 1	34.65	3.85	34.65
Saguache	1.5			7.75	0.00	04.00
San Miguel	.8			8.00		or a constant
Sedgwick	3	1		34.60	34.60	
reller	•3			2.75	0.2.00	
Washington	. 3	1		20.30	20.30	
Weld	39.2		285	455.10	3.96	1.60
	<u> </u>		1-3-0		0.00	
STATE						
TOTALS	288	1584	2458	3269 .80	2.06	1.33
	_					

Discussion of Resurvey.

Two methods of resurvey were employed during
the year. The one used depending upon the county
being gone over. In very important sections and where
numerous barberries had previously been found the
resurvey was made with as much care and diligence as
was the original inspections every precaution being
tak n to make sure that no bushes remained. In less important
sections and where the plantings were few and well scattered
the resurvey was made very thoro in the neighborhood of
these plantings and also in such sections as gave the
appearance of furnishing favorable conditions for the
propagation of the rust harboring bushes. The remainder
of such country was of course gone over more less
rapidly, depending upon its general characteristics.

As in the original survey the percent of resurvey in a count refers to that part of the total area of the county. It is evident that earlier reports indicated the percentage of the original survey which has been given resurvey since in many instances the first resurvey was greater than the original. For example counties only partially covered in the original survey were reported as one hundred per cent resurveyed. The short time consumes in making these resurveys as compared to the area of the county in which the survey was made lends further support to the theory

that the original survey was taken as the percentage basis for the resurvey.

In going over the ground a second time it was noted that in the majority of cases where the removal of the bushes was left to the tenant or land owner, due to lack of thoroness in digging out the roots, it being difficult to impress people with the necessity of so doing. In those few instances where the person living on the ground really took the interest and saw to the complete removal of the roots and watched afterwards for sprouts, eradication proved very effective. Their attitude, if it could be instilled into everyone, would prove the greatest means of forever ridding the country of this pest.

It is noticeable that, where bushes are poorly dug in the first place, their co-plete eradication becomes more difficult. Not only are the remaining roots more difficult to remove, but those remaining in the ground seem to have greater powers for sending up shoots, large bushes springing up from small roots in a very short time.

To the most effective means of completely destroying the common barberry bush and preventing the growth of sprouts is to treat the bush with a chemical such as NaCl.

Continued digging of roots whenever sprouts indicate their presence will, after a time, bring about the total destruction of the bush, but in out of the way placed this is a difficult process to carry out. Also, in many instances, we find the barberries growing in wooded and rocky regions where it is impossible toremove all the roots. In such cases and in other instances where digging is impractical chemicals furnish the only means of complete eradication.

In the south western part of the state numerous bushes of Berberis fendleri were found scattered over the hill sides. Cluster cups were present in abundance on the leaves of these bushes but this particular strain or rust did not carry over to the wheat or other grains in the adjacent fields.

There proved to be a great variation in the percentages of bushes having sprouts and in the percentages of properties having sprouting bushes, even in the same and adjacent localities. However, the percentage estimated for the territory resurveyed indicates 15 percent of the bushes to have sprouted on 20.2 percent of the properties having bushes.

A great variation was also observed in the ability of the bushes with regard to their ability to propagate their kind. Many large hedges were found bearing seeds in great quantities, and yet no seedlings were in evidence in the vicinity. In other instances a single

bush was known to produce a great many seedlings, some times reaching a considerable distance with their offspring.

In the State of Colorado we are strong believers in the possibilities of chemical eradication and think it should be employed wherever possible. We are planning a great deal on its use in handling the sprout situation the coming season. Where chemical can not be used complete removal thru digging is the only recourse. In this connection we recommend digging until sure all roots are removed followed by considerable more digging in all directions. If the co-operation of the person living on the property can be secured to the extent that he will conscientiously watch for and remove all sprouts certain death for the plant is assured. Otherwise it is necessary to make frequent visits to the place until sprouting ceases and even after that another wisit should be made after a long period to make sure no sprouts have managed to obtain life from dormant root ends.

Epidemiology Studies.

In the fall of 1921 various plots of rusted grasses were selected near the College campus with a view to making frequent germination tests of the urediniospores. The purpose of this experiment was to determine the longevity or viability of the red summer spores.

No.	The stations or publication		
1 .	3 mi.S. of Ft.Collins		Sept. 15, 1921
2	South of City Park	ff	Oct. 1, 1921
2†	n n n	Barley	tt tt tt
3	Experiment Station	Wheat	Sept. 15, 1921
4	Greeley	Wheat	Oct. 3, 1921
41	u	Oats	tt tt
Б	3 Mi. north of Loveland	Timothy	Oct. 8, 1921
51	n n n	W. Barley	n H
6	l Mi. east of Berthoud	Oats	u u u
7	2 Mi. south of Ft. Collins	W. wheat	Oct. 20, 1921

Germination tests were taken from the various plots at approximately two week intervals.

Two methods were used in making these germination tests. The first practice was to sow the spore s on a cover glass holding distilled water. The class was

inverted ovar a ring placed on a slide so as to held the distilled water. The second method and the one generally used in this experiment consisted of placing a drop of distilled water within a ring of vaseline on a glass slide. The spores were then sown in the water and the whole placed on a petri dish containing a moist blotter. The spores were incubated at a constant temperature of 20 degrees C.

Germination tests were made from each plot at the intervals stated from the date of the first test until zero or no germination was shown. The following data was obtained.

No	• Host	First Made		% spores viable]			spores
1	Oats	Sept. 1	5, 1 9 2 1			12,		viable O
2	11	Oct. 1,	1921	84	Jan.	13,	1921	0
21	Barley	11 11	11	75	11	11	11	0
3	Wheat	Sept. 1	6, 192 1	89	Jan.	16,	1921	0
4	II	Oct. 3,	1921	40	10	st		
41	Oats	Oct. 3,	1921	85	Nov.	2,	192 1	0
5	Timothy	Oct. 8,	1921	20	Dec.	2,	1921	0
51	Wild Barley	u n	ţţ	40	Mch.	12,	1922	Ò
6	Oats	n, n	11	50	Jan.	16,	1922	0

Each germ nation test, from the starting date to the time of zero germination, showed a slight decrease over the preceding test with regard to per cent of viable spores.

The spores remaining viable for the greatest length of time were found on Hordeum jubatum (wild barley) at Station No. 5', the strain on this grass being identified as Puccinia graminis tritici by Dr. E. C. Stakman.

Temperature and precipitation observations were made during the experiment and the temperatures at the Stations were observed when the spores were collected for the germination tests.

Stubble well supplied with red spores was placed out in the open in a pasture, being exposed to full weather conditions. The material was protected by having a small wire fence built around it. Several bundles of the stubble were used in the experiment and each was numbered, the spores being tested for germination at frequent intervals.

The first test was made on September 27, 1921 all proving viable, the highest running 75 per cent. On December 20, 1921, all tests have zero germination both with 48 and 7r hour incubation periods. Another test was made later and with the same results. It was concluded that the red spores could not remain viable over winter when exposed to the elements on stubble.

Asimilar experiment was tried in which the spore laden stubble was placed in a hole in the ground which

was about four feet deep. The results obtained were practically the same as where the spores were exposed idrectly to the elements.

The same results were also obtained when the stubble and spores were placed under several inches of of leaves. In fact the spores remained viable for a shorter length of time in this test than in the one where they were placed out in the open.

IV. Inoculated wheat in pots Decemver 28, with Puccinia graminis, form 3, from St. Paul.

Set in incubator 48 hours and set out on bench.

Jan. 3, 1923, flecks appearing and rust starting to develop on the Marquis.
Placed out of doors to harden.

Jan. 12, 1923, pots exposed to out of door conditions.

Jan. 19, 1921 germination tests taken at 20 degrees C.

Jan. 20, 1921 Marquis wheat placed in greenhouse for further inoculation.

January 22, 1920, spores germinated 85 %.
Note: Inoculation at this time successful on the Marquis. Kanred showing resistance.

Jan. 26, 1923, Spores collected from wheat exposed and germinated at 20 degrees C. Covered with $\frac{1}{2}$ in snow.

Jan. 29, 1923 30% germination

Jan. 30, 1921 40% "

V. Future work when plenty of rust material is available.

Pots kept in dark chamber. Pots exposed to open weather. Pots covered with debris. Pots kept for furture inoculation work.

Germination tests to be made in the usual manner each week.

Inoculations by putting spores in distilled water sprayed on plants, also by thumb and finger method.

VI. Weather reports

VII. Early spring observations.

First indication of rust in field.

First infection of barberry bushes or sprouts.

Overwintering of Urediniospores.

No field plots could be located in this section for the carrying on of overwintering tests because of the absence of rust in the fields. Observations were made by Professor C. D. Learn and E. A. Lungren.

Following are some of the results obtained from the carried on:

I. Rust on leaves, Puccinia graminis tritici, collected summer of 1922.

Set in germinator Dec. 8, 1922.

Three petri dishes containing moist blotters with slides holding distilled water within a ring of vaseline were used. The spores were placed in the distilled water.

II. Red rust from stubble, collected in 1922. Eight petri dishes, each containing red spores in distilled water on glass slide. 20 degree C. and 30 C.

Set in germinator December 15, 1922

$\mathbb{N} \circ ullet$	% germination	% germination	Temp. K
	Dec. 18, 1922	Dec. 20, 1922	-
1	О	0	20 C.
. 2	11	, th	20 C.
. 3	II.	tt	11
4	11	11	r tt
5	11	1!	. 30 C•
6	11	11	tt .
7	rt .	. 11	. 11
8	1 !		11

III. Wheat planted in 6 in. pots in greenhouse Nov. 2, 1922.

2	Plots	13274	Certified	Marquis
11	11	13408	ĮĮ	11
ff	t1	13407.	1f	Kanred
11	11	I3409	. If	11
11	II	1310	Ħ	11

LEGEND

Location of chemically treated bushes July 1 to December 31, 1922

CHEMICAL ERADICATION

Chemical eradication was first practiced in the state of Colorado during the latter part of the season of 1923. The scouts in the field supplied the NaCl, which was the chemical used, at their own expense, there being no cost to the Government. Bushes were treated in the counties of Adams, Arapahoe, Boulder, Jefferson, Larimer and Weld. Varying amounts of chemical were used, but no case did the quantity applied equal that advised by the Departmental circulars which were received later in the year. However, the treatment with these lesser quantities seems to have been effective so far as the bushes in the vicinity of Fort Collins were concerned. None of the others have been revisited and therefore we do not deem it advisable to draw conclusions as to the results of treating the bushes with che icals. Next year definite data on the subject will be obtainable

HISTORY OF PLANTINGS.

One of the most important barberry hedges in Colorado was located on the far m of J. C. Ruple, three miles west and three miles south of Loveland. The hedge was very old and the bushes very large, dynamite and teams with block and tackle being utilized in removing them. These barberries proved to be prolific bearers of fertile seeds and seedlings and bushes were well scattered over the surrounding Thru the farm runs the Home Supply Ditch country. one of the larger irrigation ditches of the community, and this proved an effective means of carrying seeds thruout the rich agricultural region which it waters. Sprouts and escapes still offer a problem in this district. They have, however, for the most part, been removed with the result that for the past two years very little rust has been experienced in the community where formerly the loss from this disease was so high as to almost prohibit grain growing.

The Handy ditch furnishes a similar example of the spread of the barberry over the country. This ditch received its waters from the same stream as does the Home Supply. Near its head was located the farm of Mrs. M. A. Alps whereon grew some highly prized common

barberries. Birds deposited the seeds under the trees along the banks of the ditch and from there the bushes were being spread over the section served by this irrigation system.

One large bush, located on the Poudre River, north west of Ft. Collins, and near the foot hills, proved to be the source of a great many rust spreading barberries scattered along the banks of this stream and in adjacent field.

Mr. Alfred Bosbec, living north of Loveland volunteers the information that since the removal of his barberry hedge he has been able to grow wheat in a field that formerly proved almost a total loss when put into grain because of the rust attack.

The Wolhurst farm, south of Denver, had a flourishing barberry hedge located near the farmhouse on the
bench above the river. From here the bushes has
spread down to the river and had commenced their marck
along its banks. Since the removal of this hedge and
the escapes from the surrounding territory very little
rust has been experienced in the section where formerly it
gave a great deal of trouble.

A great many instances, similar to the above could be cited over various parts of the state, and it is with great satisfaction that we note the decreasing losses thru rust.

DISCUSSION OF THE YEAR'S WORK.

Considerable has been accomplished during the past year in theway of eradicating the common barberry from the state of Colorado. We feel quite safe in agying that practically all the bushes of the state have been discovered, although of course there may be a few in hiding in certain parts of the state which have not been thoroly gone into, if worked at all. The main problem at present is to completely do away with those bushes found and prevent the shooting up of sprouts every year. This we hope to be able to accomplish in most instances by means of chemical eradication. The comparative few that can not be dealt with in this way will be killed off by thoro and persistant digging of roots wherever their location is indicated by the presence of sprouts.

During the year three scouts were employed in the field and for various lengths of time, one of them being the State Leader. Each one was equipeed with a car and assighed certain territory to cover, this proved the most economical system of survey so far practiced in view of the fact th t considerable more money was spent in Administration and Overhead than in previous years. The actual field expenses averaged \$.151 per running mile and \$.174 sq. mile surveyed. This same method was employed during the season of 1920 and it is possible that

the actual field expenses for that year were slightly under those of 1922. However, the nature of the work that season differed somewhat from that performed the past year and after everything is taken into consideration the average field expenses for the two years appear practically the same for the two years, i. e., those of 1920 as compared to those of 1922. However, too definite conclusions should not be drawn from such comparisons as the controlling factors are so varying with different years.

Little was done toward publicity during the year for the reason that there was nothing to be gained by it.

The work had been carried on practically all over the state and every section was familiar with its nature and purpose.

The chemical treatment of bushes was not carried on to the degree that we would have wished because of the fact that the field men were not in possession of the details necessary to the carrying on of this phase of the work effectively. However, a start was made and future work will be guided by past experience.

One beautiful example of the spread of rust from the barberry was observed near the town of Burlington in Kit Carson County. On recommendation of the county agent this section was not surveyed the previous year because he know that there were no barberry bushes in the community. Last year an outbreak of rust in the

grain fields was called to the attention of the State

Leader and an investigation was made. As a result a

prosperous barberry was discovered in the center of the

epidemic from which the spread of rust could be traced.

A few other bushes were located over the state by this

same method and it is our plan to watch closely

for rust outbreaks next year.