

STEWARDSHIP INCENTIVES PLAN

For:

Cumberland Lode Forest

NEAL ARMSTRONG, Conservator
MARGARET HASCAL, Owner
13687 West Virginia Drive
Lakewood, CO 80228
(303) 987-3534

★ KEN ARMFIELD 543-1094
8.8 acres

Part of
NE1/4 NE1/4, Sec 7, T1S, R71W, S.P.M.
and
NW1/4 NW1/4, Sec 12, T1S, R72W, S.P.M.

Prepared By:

Douglas J. Stevenson
Colorado State Forest Service
936 Lefthand Canyon
Boulder, CO 80302
(303) 442-0428

June 6, 1997

This management plan has been prepared at my request to guide my forest management activities which I voluntarily apply on my property. Activities recommended in this plan are appropriate to my objectives and will benefit the natural resources on my property.

Neal Armfield, Conservator

Date

TABLE OF CONTENTS

Subject	Page
I. FOREST STEWARDSHIP MANAGEMENT PLAN	1
II. TABLE OF CONTENTS	2
III. OBJECTIVES	4
IV. GENERAL DESCRIPTION	
A. Property Location	4
1. Area	4
2. Boundary Monuments	4
3. Access	4
B. Topography	5
1. Slope and Aspect	5
2. Elevation	5
3. Geology	5
4. Soils	6
C. Local Markets	6
D. Typology and Stand Identification	6
E. Land Use	8
1. Current	7
2. Historical	8
F. Desired Conditions	8
G. Impact on Neighbors and Nearby Communities	9
V. INVENTORY (Species, Stocking and Age by Product Classes; Insect and Disease Presence, Damage and Risk; Fire Hazard and Risk Factors; Operational Limitations of Soil and Terrain; Access Roads and Trails, Limita- tions; Riparian and Wetland Features; Silvicultural Objectives; Cultural Features; Noxious Weeds; Wildlife Species, Habitat, Condition and Impact; Endangered Species; Scenic and Aesthetic Qualities; Known Archeo- logical Sites)	9
VI. PRESCRIPTION BY MANAGEMENT UNIT (Silvicultural Prac- tice, Implementation; Type and Amount of Product; Restrictions and Special Requirements; Additional Roads, Trails (if Applicable); Work Standards; Imple- mentation Priority)	11
VII. IMPLEMENTATION SCHEDULE (Recommended Practices, Year Implemented, Units Completed)	13
(Records and Maps; Treatments, Dates Completed, Volume	
VIII. Harvested; Price Received, Management Costs	14

STEWARDSHIP INCENTIVES PLAN

3

IX. MAP	15
X. APPENDICES	
I. Tree Species	16
A. Douglas-fir	19
B. Limber Pine	19
C. Ponderosa Pine	20
II. Dwarf-mistletoe	23
A. Service in Action	24
B. Dwarf-mistletoe Addendum	26
III. Soils	27
A. Fern Cliff Series	28
B. Peyton Series	30
IV. Glossary	32
V. Bibliography	40

OBJECTIVES

The forestry objectives for this property are:

1. Consistent with the Stewardship Incentives Programs, to improve health and vigor of the forest and enhance its productivity.
2. Practice silviculture and multiple use management, giving particular attention to protection of wildlife habitat and recreational qualities of the area.
3. Preserve the aesthetic qualities of the property.
4. Protect soil and water resources.

GENERAL DESCRIPTION

Property Location

The property is located about at the four-mile marker on Magnolia Road, straddling the road. The address is 3900 Magnolia Road.

Area

The property contains 8.8 acres, as determined from CSFS' ARC/INFO data base. 1.0 acres is occupied by a cabin, parking area and mine tailings, but still counts as forested for Stewardship Incentives Program purposes.

A breakdown is shown below:

Forested	7.8 Acres	
	<u>1.0 Acres</u>	House and Yard
	8.8 Acres	STEWARDSHIP ACRES

Access

There is an existing driveway entering the property from Magnolia Road. A different driveway provides access to the west end of the property. The east end of the property is extremely steep with difficult access. The east drive is the only fire access to the cabin; in the event of a serious fire, escape would be cut off.

The mine and tailings in front of it would be the best safety zone in the area.

The west end of the property is flatter and somewhat protected from below by Magnolia Road. There is less fire risk here and trucks can access the area for thinning and maintenance work.

TOPOGRAPHY

Slope and Aspect

Aspect is primarily north and northeast. On most sites, slope is around 60%, with slopes up to 100% in places.

Elevation

The lowest point is along the north property line where Calhoun Gulch leaves the property at an elevation of about 7900 feet above sea level. The highest point is at the southwest corner of the property with an elevation of about 8300 feet above sea level.

Geology

Precambrian rocks now about 1.8 billion years old were intruded about 1.7 billion years ago by the Boulder Creek Granodiorite Formation. This is the bedrock throughout the property.

A north-northwest trending fault of Precambrian Age passes near, but does not cross, the property. It has occasionally been reactivated.

Lower Paleozoic rocks (Cambrian through Mississippian) are missing in this area. It is thought that these rocks once existed, but were eroded away during Early Pennsylvanian times when the area was uplifted on the northeast flank of the Ancestral Front Range uplift, one of several northwest-trending mountain ranges that comprised the late Paleozoic Ancestral Rocky Mountains. These mountains (Ouachita Orogeny) resulted from the reactivation of Precambrian structures when Africa collided with South America and the southern edge of North America. Gravel and sediments washing off the Ancestral Front Range were deposited as the Fountain Formation, which was later uplifted to form the Flatirons. By the late Paleozoic, the Ancestral Front Range was eroded to a set of low hills.

In the Early Cretaceous, the area began to subside and was eventually buried under almost 10,000 feet of marine sediment (Pierre Shale).

In the Late Cretaceous-Early Tertiary (about 67.5 million years ago), the Laramide Orogeny uplifted a mountain range with much the same configuration as the present day Front Range. The mineralization that created the Colorado Mineral Belt and our local mineral deposits (including the Cumberland) occurred at this time. Erosion about balanced uplift so relief was never great, much less than at present. By the Late Eocene uplift ceased, leaving a low-profile range of hills. Most of the faulting and eastward tilting that raised the Flatirons into position occurred during the Laramide Orogeny.

Intrusive volcanic activity occurred to the east during the Paleocene, creating Valmont Dike and the basalt formation now being mined by Andesite Rock Company. No Paleocene igneous activity involved this property.

During the Oligocene this region was reduced to a plain, similar to eastern Colorado today with an elevation of about 3000 feet. In the Miocene, thermal uplift and east-west expansion formed the Rio Grande Rift and began the rise of the modern Front Range, which continues to rise today.

The property has never been glaciated (The nearest glacier reached the flat at Barker Reservoir east of Nederland.); although, Wisconsin-age and later climate fluctuations have had major impacts on plant life and continue to have an effect, even now.

Soils

The steeper, north-facing slopes are on Fern Cliff soils. These grade into Peyton soils toward the west end of the property.

LOCAL MARKETS

Sawtimber markets in Boulder and vicinity are severely limited and do not apply on this property, anyway. Firewood markets are weaker than they were ten years ago, but still readily able to handle expected cutting, of which there will be little due to steepness of the terrain.

TYOLOGY AND STAND IDENTIFICATION

Western forests are typed by the dominant tree species occurring, provided that species makes up a plurality of stocking. When no species makes up at least 20% of the stand, the type is listed as "mixed." The stand consists of ponderosa pine throughout.

Size classes are as follows:

- Class 1A: Seedlings (Less than 4.5 feet tall).
- Class 1B: Saplings (4.5 feet tall to 5.0 inches DBH).
- Class 2A: Small poles (5.0 to 7.0 inches DBH).
- Class 2B: Large poles (7.0 to 9.0 inches DBH).
- Class 2C: Near-merchantable (9.0 to 11.0 inches DBH).
- Class 3A: Small sawtimber (11.0 to 15.0 inches DBH).
- Class 3B: Medium sawtimber (15.0 to 21.0 inches DBH).
- Class 4: Large sawtimber (21.0+ inches DBH).
- Class 5: Large old growth (21.0+ inches DBH and dating from pre-settlement).

A stand is classified by adding together stocking figures, starting with the highest class, until a minimum level of 32.5 trees per acre, 10 square feet of basal area per acre or 200 board feet per acre is obtained over a minimum 3.0-acre area. This means that trees larger than the listed size class may occur in small numbers. Classes 2C through 5 are typed by board foot volume; classes 1B through 2B are typed by basal area and class 1A is typed by stem count.

There are no Class 4 or Class 5 stands on your property.

Pre-settlement stands in Boulder County are those with stand birthdates of 1850 or earlier (Age: about 140 years). Stand birthdates are determined by taking mean age weighted by volume, basal area or stem count, as above, and subtracting that from the current year, rounding the result to the nearest decade. It is possible for a younger, faster-growing class of trees to overtake an older, slower-growing class and change the stand birthdate, without any other change in the stand.

Multiple classes: Real stands are rarely even-aged or all-aged, but consist of in-between mixes. Second and third classes of trees are typed as if they were separate stands, but the result is listed along with the dominant class. Second and third classes frequently differ from the dominant class.

Typology: Your timber is ponderosa pine/Douglas-fir small sawtimber (Class 3A), heavily stocked, birthdate: 1890.

LAND USE

Current

The property is being used as a cabin site and for recreation. Mr. Armfield is interested in protecting the health and appearance of the property and, particularly, in fire protection.

Historical

The current forest typology originated following a major climatic shift at the end of the Pleistocene, about 12,000 years ago. Species that now occur at 9000 to 9,500 feet of elevation, such as lodgepole pine, Engelmann spruce and subalpine fir, grew here. Aside from being wetter than it is now and having a greater tendency to mixed-species stands, it probably wasn't a whole lot different during the Ice Age than it is now.

During the Altithermal, about 7000 to 9000 years ago, the climate became even warmer and dryer than it is now. Most of this property was probably in grass and ponderosa pine during the Altithermal.

The current forest originated about 1760 following a major fire. This is surmised from the existence of a class of Douglas-firs, on nearby property, all dating from about the same decade. The only known agent capable of large-scale land clearing over thousands of acres, is fire.

About 1853 a large fire burned the area west of Boulder, including most of western Boulder County. According to local legend, this fire was set by Arapahos, angry at being cheated by whites. One suspects the legend sprang up later so that whites would not have to take the blame for settlers' carelessness.

The property was horse-logged in the 1870s.

The township was originally surveyed in 1870 and 1888. I don't have a copy of the original survey for this township, so I can't relate the details. A dependent resurvey was done in 1926 and 1927 by Albert Rich, Arthur Brown, Roy Chase and Emil Voigt. The Surveyor General's map dated February 19, 1930 shows Magnolia Road in its current location.

DESIRED CONDITIONS

Healthy, vigorous, fully-stocked stands of trees is a goal of the Stewardship Incentives Programs. This condition need not be achieved immediately, but progress should be made in this direction.

It would be desirable to increase the area's usefulness to wildlife, indirectly enhancing recreational values. In particular, nesting sites for cavity-nesting birds and shelter piles for small animals would be beneficial.

IMPACT ON NEIGHBORS & NEARBY COMMUNITIES

The property is surrounded by U. S. Forest Service land. The only area visible from a public road is that portion immediately adjoining Magnolia Road. From across the valley cutting would be visible through binoculars, but would not be apparent unless the viewer was told where to look. Most people will not realize that anything is happening.

INVENTORY

This is a heavily-stocked, small sawtimber ponderosa pine stand with associated Douglas-fir. Scattered other species, such as aspen and Rocky Mountain juniper occur, but not in significant numbers. Sawtimber stocking totals about 1200 board feet per acre, on average. The stand averages 130 square feet of basal area per acre, or about 300 trees. It originated about 1875, apparently following cutting.

The north side of the road is too steep for fire trucks, except for those parked on the mine tailing.

Harvesting operations are not anticipated due to the steep terrain. Thinning and firewood salvage are recommended for that part of the property west of Magnolia Road and Defensible Space for the entire property east of Magnolia Road.

There are no Federal wetland areas in this stand.

Silvicultural objectives for this stand are:

1. Thin western stand to 100 basal area. Salvage firewood. It should be possible to accomplish this at little, or no, net cost by selling wood to a local firewood cutter. Remaining slash may be piled or chipped, if desired (Piling and chipping are labor-intensive and expensive; you may not want to do this part of the work.
2. Conduct a Defensible Space around the cabin. It may be possible to obtain permission from the U. S. Forest Service to carry out this practice on adjoining Federal land, as well. This should include thinning and removal of fire ladders in the area below the driveway to provide an escape route.
3. Create nesting sites for cavity-nesting birds. This can be accomplished by girdling selected trees and allowing them to die. It also means preserving suitable habitat trees that are already dead. This can also be

accomplished by putting up bird houses designed and positioned for the particular species.

Threatened and Endangered Wildlife

The U. S. Fish and Wildlife Service lists the following species for Boulder County:

American peregrine falcon, Falco peregrinus, Endangered
Bald eagle, Haliaeetus leucocephalus, Endangered
Whooping crane, Grus americana, Endangered
Eskimo curlew, Numenius borealis, Endangered
White-faced ibis, Plegadis chihi, Category 2
Mountain plover, Charadrius montanus, Category 1
Northern goshawk, Accipiter gentilis, Category 2
Black tern, Chlidonias niger, Category 2
Mexican spotted owl, Strix occidentalis lucida, Threatened
Loggerhead shrike, Lanius ludovicianus, Category 2
Boreal toad, Bufo boreas boreas, Category 2
Black-footed ferret, Mustela nigripes, Endangered
Preble's meadow jumping mouse, Zapus hudsonius preblei,
Category 2
Fringed-tailed myotis, Myotis thysanodes pahasapensis,
Category 2
North American wolverine, Gulo gulo luscus, Category 2
Swift fox, Vulpes velox, Category 2
Greenback cutthroat trout, Oncorhynchus clarki stomias,
Threatened
Plains topminnow, Fundulus sciadicus, Category 2
Rocky Mountain capshell, Acroloxus coloradensis, Category 2
Regal fritillary butterfly, Speyeria idalia, Category 2
Lost ethmiid moth, Ethmia monachella, Category 2

The following plants are also listed:

Bell's twinpod, Physaria bellii, Category 2
Larimer aletes, Aletes humilis, Category 2
Ute ladies'-tresses orchid, Spiranthes diluvialis,
Threatened
Colorado butterflyweed, Gaura neomexicana coloradensis,
Category 1
Showy prairie gentian, Eustoma grandiflorum, Category 2
Pale moonwort, Botrichium pallidum, Category 2
Purple lady's slipper orchid, Cypripedium fasciculatum,
Category 2

The peregrine falcon and bald eagle have been observed in Boulder County numerous times since 1987. The white-faced ibis has been observed just across the county line at Continental Pond in Weld County in the fall of 1994.

The purple lady's slipper has been observed several times since 1987.

The Mexican spotted owl occurred in Boulder County historically, but has not been seen here since the Threatened and Endangered Species Act was passed in 1973. The nearest known nest is located south of Denver in Douglas County. A detailed search of Coal Creek, Boulder and Lefthand canyons and their tributaries is being done this year; if results are negative, Mexican spotted owl will probably be removed from the Boulder County list.

The northern goshawk is favored by the many age classes of trees created by rotational cutting. As long as buffer zones are left around nests (30 acres) and cutting activities avoid a fledging area (400 acres) during the fledging season, there should be no problems. Though this bird may occur in Boulder County, I am not aware of it.

To the best of my knowledge, the black-footed ferret is listed only because its prey (prairie-dogs) is found here. I do not know of any sightings. The ferret is a creature of the plains and would not pose a problem for most mountain projects.

The whooping crane was included on the list because it might come here during migration. Again, I am not aware of any sightings. The same applies to the Eskimo curlew.

The Ute ladies'-tresses only occurs below 7000 feet; there is no need to worry about it on this property.

The black tern is a shorebird and is a concern around creeks and reservoirs, but not in the mountains, away from water.

The black-footed ferret, preble's meadow jumping mouse, fringed-tailed myotis, Colorado butterflyweed and showy prairie gentian occur only in the plains. The mountain plover, northern goshawk, Mexican spotted owl and purple lady's slipper occur only in the mountains (except for a single spotted owl sighting in Adams County).

Other listed species could occur in either mountains or plains, especially areas where forest and prairie intermix.

There are no known archeological sites within this stand.

PRESCRIPTION BY MANAGEMENT UNIT

There are two management units in this stand:

1. The most-urgent need is for a Defensible Space practice around the cabin to reduce the risk from wildfire. First, thin the stand surrounding the cabin out to a distance of 75 feet so that adjoining tree crowns do not

touch and cannot spread fire from tree to tree to the cabin. Next, prune low-hanging branches and saplings so fire cannot climb from the ground into the tree tops. Clean up accumulations of burnable debris and slash out to a distance of fifty feet (75 feet on the down-hill side). I would also suggest enclosing porches so that burnable leaves and needles cannot collect beneath them and re-siding the cabin with something less flammable than the raw lumber now used for siding.

To reduce the danger from fire, the rest of the lot property east (north) of Magnolia Road needs to be thinned with a fifty-foot strip downhill from the drive being treated as described above.

2. That part of the property west (south) of Magnolia Road should be thinned. It may be possible to do this commercially as a firewood sale. CSFS can help by laying out the sale area, marking the "cut" trees and tallying the volume. We can then find a cutter, prepare the agreement and supervise the sale. Most of this cost is subsidized and sale proceeds normally more than cover the rest. You won't make a big profit, but at least the work gets done. Slash should be left on site and scattered unless there is an opportunity for chipping.

Defensible Space is a fire-safety practice that seeks to create an area around a building (particularly a house) or a road that will not readily burn, or if it burns will do so quietly in a way that will not endanger fire fighters' lives or equipment. This allows a strike team or pumper crew to defend a structure that they would otherwise have to abandon in order to protect their own lives. It involves thinning the stand around the building and below the road so tree crowns do not touch and cannot transmit fire from tree to tree to the structure. It also involves pruning low-hanging limbs and removal of selected bushes and small trees to eliminate path-ways that fire can use to climb into the tree crowns, or climb up the side of a building. Also included (but not funded) are safety practices such as keeping woodpiles (spark catchers) away from the building, enclosing areas beneath decks so debris cannot collect there and cleaning needles and leaves off rooves and out of gut-ters. Accumulations of debris are removed and slash is chipped or hauled away.

The practice is eligible for 65% of actual cost, up to \$750 in SIP cost-sharing (See above.). Even without cost-sharing, it is well worth the cost, particularly to a house that is vulnerable to fire driven by a south or southwest wind.

IMPLEMENTATION SCHEDULE and RECORD

RECOMMENDED PRACTICES, YEAR IMPLEMENTED, UNITS COMPLETED

1997:

Conduct a Defensible Space practice around the house. The Colorado State Forest Service can assist by marking trees to cut, helping find contractors, etc.

1998:

Extend Defensible Space area to include the rest of the property north of Magnolia Road, giving particular attention to the area immediately below the drive. This may be eligible for cost-sharing of 65% up to \$200 per acre.

1999:

Prepare firewood sale for area west of Magnolia Road; sell.

2000:

Administer firewood sale.

2001:

No activities.

2002:

Revise and update this plan. Forests are not static. They change over time. Also, your desires for the property may change, too, and it is unlikely that the practices recommended will actually be completed exactly on schedule.


2003-2006:

No activities.

2007: Have new plan prepared.

RECORDS and MAPS; TREATMENTS, DATES COMPLETED, VOLUME HARVESTED;
PRICE RECEIVED, MANAGEMENT COSTS

The following section is expandable and meant to serve as a log of
forest management work being done.



APPENDIX I

Tree Species Observed on Cumberland Lode Forest

DOUGLAS-FIR (Pseudotsuga menziesii (Mirb.) Franco¹)

BOTANICAL FEATURES

Leaves $\frac{3}{4}$ " to $1\frac{1}{4}$ " long, yellow-green or blue-green, more or less flattened, standing out from all sides of the twig or with a tendency to be somewhat 2-ranked; apex rounded-obtuse or rarely acute, stomatiferous below, persistent for 8 or more years.

Cones 3" to 4" long, pendent, ovoid-cylindric, with exserted, 3-lobed, forklike, appressed or strongly reflexed bracts; seeds triangular, terminally winged; about 42,000 seeds to the pound, dewinged.

Buds fusiform, sharp-pointed, lustrous brown.

Bark on young stems smooth except for resin blisters; at length becoming 6" to 24" thick on old trees, and then divided into thick reddish-brown ridges separated by deep irregular fissures. In a few instances the bark is "tight" (fine-textured) on old trees and corky on others, particularly those of the mountain form.

GENERAL DESCRIPTION

Douglas-fir, monarch of Pacific Northwest forests, was first observed by Menzies on Vancouver Island when he accompanied the British naval captain Vancouver on an expedition to the Pacific Coast in the early 1790s. For more than a quarter of a century this tree was variously classified as a spruce, hemlock, true fir, and even as a pine; in fact logs exported by the Hudson's Bay outpost near the mouth of the Columbia River were listed in European ports as "Oregon pine," a name which has persisted in the trade to this day, especially in Australia. It remained for David Douglas, a Scottish botanist sent out by the Royal Horticultural Society in 1825, to study this tree, to show that it was sufficiently different to be considered as separate from other previously described conifers; later Carrière coined the new generic name Pseudotsuga. This name was a rather unfortunate choice, since it literally means "false hemlock." The common name, Douglas-fir (The names red fir and yellow fir have been used by loggers and lumbermen to differentiate locally certain specimens on the basis of ring width, color and softness.), commemorates Douglas, and in addition serves to distinguish this species from the true fir (Abies).

Douglas-fir is a dimorphic species with two more or less distinct forms (Several European workers have claimed that there are three species of Douglas-fir; this is based largely upon needle structure. Studies by W. E. Kilgore at the New York State College of Forestry have failed to substantiate this viewpoint.). One of

¹Harlow, William M. and Harrar, Ellwood S. Textbook of Dendrology, Fifth Edition, McGraw-Hill, 1968.

these is restricted to the forests of the Pacific slope, and the other to those of the Rocky Mountain region.

The Rocky Mountain form of Douglas-fir is considered distinct from the coast form by some taxonomists, who accordingly classify it as Pseudotsuga glauca Mayr. or Pseudotsuga menziesii var. glauca (Beissn.) Franco. However, in certain sections, the two types intergrade. Usually the foliage of the Rocky Mountain tree is blue-green, but sometimes trees with blue-green foliage and others with yellow-green leaves are found standing together. Similarly, although yellow-green crowns are typical of the coast form, some trees show a blue-green coloration. The principal botanical difference between these two forms lies in the structure of their cones. Rocky Mountain trees have small cones rarely 3 inches in length, with much-exserted and strongly reflexed bracts. By contrast, the cones of the coast form are often 4 in. long and have straight, more or less appressed bracts.

Douglas-fir comprises about 50 percent of the standing timber of our western forests. It produces more timber than any other American species and at the present time furnishes about one-fifth of the total annual cut.

ROCKY MOUNTAIN FORM

The Rocky Mountain form of Douglas-fir rarely exceeds a height of more than 130 ft or a diameter of 3 ft. It occurs in both pure and mixed stands with ponderosa pine, western larch, and grand fir. Other associates include western hemlock, western white and lodgepole pines, Engelmann spruce, white fir and aspen. Douglas-fir is more tolerant than these except the hemlock and spruce.

Although most abundant on moist sites, Rocky Mountain Douglas-fir is quite drought resistant and is often found on arid areas with ponderosa pine. It is frost-resistant and hardy in the East and is a common ornamental of that region. The trees are grown for timber in Europe and have been planted successfully in many parts of the world.

RANGE

Western United States and British Columbia. Altitudinal distribution: sea level to 5,000 ft along the coast; 4,000 to 6,000 ft inland; 10,000 ft in the southern Rocky Mountains.

LIMBER PINE (Pinus flexilis James)²

DISTINGUISHING CHARACTERISTICS

Needles in 5's about 2½" long, clustered near the branch ends, dark green, stout, rigid, stomatiferous on all surfaces; resin canals dorsal. **Cones** 3 to 10 in. long, cylindrical, the scales thickened, and slightly reflexed at the apex; **seeds** large, with rudimentary wings or wingless. **Bark** on young stems smooth, silvery white to light gray or greenish gray; that on old trunks dark brown to nearly black, separated by deep fissures into rectangular to nearly square, superficially scaly plates or blocks.

GENERAL DESCRIPTION

Limber pine was first observed near Pike's Peak by Dr. Edwin James, an army surgeon attached to Long's Mountain Expedition of 1820. Like other relatively inaccessible trees of high altitudes, limber pine is primarily of importance in protection of valuable watersheds. Ordinarily the tree attains but small proportions, varying from 30 to 50 ft in height and from 15 to 24 in. d.b.h. (max. 85 by 7 ft). The bole is stout, noticeably tapered, and supports a number of large plumelike often drooping branches. The result is an extensive crown which not infrequently reaches to within a few feet of the ground. Young trees develop a long, sparsely branched taproot which is later supplemented by several laterals.

RANGE

East slopes of the Rocky Mountains in southern British Columbia and southern Alberta, south along the mountains to Arizona and New Mexico; west to the mountains of southern California, and north along the Sierra Nevada to northern California; east through Nevada and Idaho (one outpost is found in the Black Hills of South Dakota). Altitudinal distribution: 4,000- to 10,000-ft elevation in Montana, Wyoming, and Idaho; 4,500 to 11,500 ft in Colorado; 8,000 to 11,800 ft in southern California.

²Harlow, William M. and Harrar, Ellwood S., Textbook of Dendrology, Fifth Edition, McGraw-Hill, New York, 1968.

PONDEROSA PINE (Pinus ponderosa Laws.)³

BOTANICAL FEATURES

Needles in 3's, or 2's on the same tree, 5" to 11" long, dark gray-green to yellow-green, flexible, persistent until the 3rd season. Crushed needles have a turpentine odor similar to that of most other pines.

Cones 3" to 6" long, ovoid to ellipsoidal, sessile, solitary or clustered; usually leaving a few basal scales attached to the twig, when shed; apophysis dark reddish brown to dull brownish yellow, transversely ridges and more or less diamond-shaped; umbo dorsal, with a slender, often deciduous prickle; seeds ¼" long, ovoid, slightly compressed toward the apex, brownish purple; wings moderately wide, about 1" long; about 12,000 (6,900-23,000) seeds to the pound.

Twigs stout, exhaling a turpentine odor when bruised; buds usually covered with droplets of resin.

Bark brown to black and deeply furrowed on vigorous or young trees (bull pines); yellowish brown to cinnamon-red and broken into large flat, superficially scaly plates separated by deep irregular fissures on slow-growing and old trunks.

GENERAL DESCRIPTION

This is the most important pine in western North America, and in the United States is found in commercial quantities in every state west of the Great Plains. At present it furnishes more timber than any other American pine and in terms of total annual production of lumber by species is second only to Douglas-fir.

Ponderosa pine is a large tree 150 to 180 feet high and 3 to 4 ft in diameter (On the best sites, 300-year-old dominant trees average about 175 ft high and 48 in. d.b.h.) (max. 262 by 8.6 ft). Even though this species commonly forms open parklike forests, the boles are ordinarily symmetrical and clear for one-half or more of their length; short conical or flat-topped crowns are characteristic of old trees. Four-year-old trees may have tap-roots four to five feet long. Moderately deep wide-spreading laterals develop as the trees get older. Ponderosa pine is not exacting in its soil requirements, but trees on thin, dry soils are usually dwarfed. Its occurrence on dry sites with the nut pines and certain of the junipers is indicative of its great resistance to drought. This species attains its greatest development, however, on the relatively moist but well-drained western slopes of the Siskiyou and Sierra Nevada Mountains of southern Oregon and California, respectively.

³Harlow, William M. and Harrar, Ellwood S., Textbook of Dendrology, Fifth Edition, McGraw-Hill, 1968.

Ponderosa pine occurs in pure and mixed coniferous stands. Excellent pure forests are found in the Black Hills of South Dakota, the Blue Mountains of Oregon, the Columbian Plateau northeast of the Sierra Nevada, and in northern Arizona and New Mexico. It is also commonly the most abundant tree in mixed coniferous stands; east of the summit of the Cascade Range in Washington and Oregon it occurs with western larch, Douglas-fir, and occasionally lodgepole pine; in the central Rocky Mountains with Douglas-fir; and in California with Jeffrey and sugar pines, incense-cedar, Douglas-fir, and white fir. On the Fort Lewis plains in western Washington, near Puget Sound, ponderosa pine is occasionally found in association with Douglas-fir and Oregon white oak.

Small quantities of seed are produced annually, but large crops are released only at intervals of three to five years. Under forest conditions germination as high as 50 percent may be anticipated, but in the nursery this figure can be increased to 80 percent. Seedlings can exist under the canopy of the parent trees, even though they grow quite slowly, and in such conditions often attain a height of only 3 to 4 ft during the first 15 to 20 years. Reproduction is best in clearings made by fire or logging. The seedlings will grow on sterile sites and have been planted extensively in the Nebraska sand hills and elsewhere. Ponderosa pine is classed as intolerant.

The rapidity of growth has a marked effect on the general appearance of the trees of this species. Young, vigorous specimens commonly develop dense crowns of dark foliage, and bark which is dark brown to nearly black, more or less corky, and deeply furrowed. In contrast, the foliage of old-growth or slow-growing trees is yellow-green, and the bark yellow-brown to cinnamon-red and plated. Those of the first type are generally called "bull" or "blackjack pines," and to some woodsmen ponderosa pine and bull pine are different trees. Fast-growth bull pines 150 years of age found near Cle Elum, Washington, measured 30 to 40 in. in diameter, while more typical ponderosa pines occurring in the same vicinity were only 10 to 14 inches in diameter at the same age. The growth of this species varies considerably with locality. In California, trees 120 years of age averaged 23 in. d.b.h., while in Arizona trees of the same age were only 16 in., and in the black hills 10½ in. Trees over 500 years of age are seldom encountered (Keen considers that this pine may reach an age of 800 years, while Mills reported a tree in southwestern Colorado with 1047 rings.). Severe damage is caused by bark beetles, and ponderosa pine is also attacked by more than 100 other kinds of insects. Fires kill seedlings and cause considerable damage even to large trees. Severe fires in the past have completely destroyed hundreds of thousands of acres of ponderosa pine forest. Other destructive agents include mistletoe and fungi.

The common name ponderosa pine is identical with the species name. Previously called western yellow pine, logs of this tree were also

sold under such names as Arizona white pine, California white pine, and western soft pine, since the wood resembles that of the white pines rather than that of the hard, moderately heavy wood of the southern yellow pines. Finally, the name ponderosa pine was adopted by the U.S. Forest Service, and it is now accepted by the industry.

RANGE

Western North America. Altitudinal distribution: 5,000 to 8,000 ft in Arizona, 3,300 to 6,000 ft in Montana and South Dakota, 2,000 to 7,000 ft in northern Idaho, sea level to 6,200 ft in British Columbia and Washington, sea level (Columbia River Valley) to 7,000 ft in Oregon, 300 to 7,000 ft in northern California, 4,000 to 9,000 ft in southern California; for the most part a tree of relatively low elevations.

APPENDIX II

Dwarf-mistletoe

DWARF-MISTLETOE⁴**Quick Facts**

Dwarf mistletoe causes a serious forest problem in many parts of Colorado.

Hosts for mistletoe include most members of the pine family.

The seeds of mistletoe are dispersed in August and early September.

The ultimate effect of dwarf mistletoe is premature death of the affected tree.

Dwarf mistletoes (Arceuthobium spp.) are a major problem in Colorado forests on ponderosa and lodgepole pine. Other members of the pine family, Douglas-fir, pinyon and limber pine are damaged occasionally. Nursery and ornamental plantings seldom are attacked; however, this parasite can be introduced into an area by the planting of collected stock infected with dwarf mistletoe.

Dwarf mistletoes are small, leafless, parasitic flowering plants. The seeds, explosively discharged from the fruit, are very sticky and adhere to any surface they strike. Seeds that adhere to young branches of susceptible trees germinate and the mistletoe plant penetrates the bark. These seeds generally are dispersed in August and September.

This parasite is easily identified by the yellow to green or brownish-green segmented shoots that protrude from the infected part of the tree. These perennial shoots are 2 to 6 inches (5-15 centimeters) long and $\frac{1}{8}$ - to $\frac{1}{4}$ -inch (.3-.6-cm) in diameter.

The "roots" of the dwarf mistletoe are imbedded in the bark and phloem of the tree. The parasite produces secondary root-like structures called "sinkers" that become imbedded deeper in the wood as the twig adds its annual growth rings. These "roots" provide the parasite with nutrients obtained from the living tissues of its host.

Symptoms

The first symptom of dwarf mistletoe infection is a slight swelling of the bark at the site of infection. As the "roots" of the parasite become more extensive in the host, a distorted branching habit or witches' broom may form. The witches' broom diverts food from uninfected parts of the tree, subsequently reducing vigor and causing premature death of the tree. Infected trees that do not develop witches' brooms usually have visible mistletoe shoots protruding

⁴Swift, C. E. and Dickens, L. E. Dwarf-Mistletoe, Colorado State University Extension Service, Service in Action Leaflet No. 2.925.

from the infected area; however, shoots are not formed until two to three years after infection.

Control

Pruning is the best control measure available for reducing or eliminating dwarf mistletoe infections in ornamental trees or urban forests. Trees severely infected in the upper branches or those with only a few live branches should be cut. Trees with high, unreachable mistletoe infections will continue to rain seeds on nearby trees if not cut down.

Lightly infected trees can be freed from the parasite by pruning off all infected branches. All branches to be pruned should be cut off flush with the trunk. The entire branch should be removed. The trees should be examined every two or three years and any infected branches pruned off. The mistletoe shoots die as soon as the branch is cut, consequently burning pruned-off branches is not necessary.

If the mistletoe on a branch is close to the trunk the infection may have already entered the trunk. Shoots will form on the trunk even if the branch is removed. When pruning infected limbs, the following guidelines should be used to insure the trunk is free from infection. Trees with infections closer than indicated should be cut down to remove a future source of infection.

Branch diameter (outside bark)	Distance of infection on branch from trunk
Under 1.0 inch (2.5 centimeters)	6 inches (15.2 cm)
1.1 - 2.0 inches (2.8 - 5.1 cm)	8 inches (20.3 cm)
2.1 - 3.0 inches (5.3 - 7.6 cm)	10 inches (25.4 cm)
3.1 - 4.0 inches (7.9 - 10.2 cm)	12 inches (30.5 cm)

In some cases a highly desirable tree with a trunk infection cannot be removed for aesthetic or other reasons. In these instances, the mistletoe shoots must be knocked off periodically as they appear to prevent further spread.

In heavily infested areas, nonsusceptible trees can be planted to replace cut trees. Ponderosa pine areas can be planted to:

Douglas-fir	Pinyon pine	White fir
Limber pine (sic)	Blue spruce	Rocky Mountain juniper

In lodgepole pine areas, the following trees can be substituted:

Engelmann spruce	Subalpine fir	Douglas-fir
------------------	---------------	-------------

Hardwoods such as ash, birch and aspen, also can be planted in affected areas because dwarf mistletoes do not attack hardwood trees.

DWARF-MISTLETOE ADDENDUM⁵

Three species of dwarf-mistletoe occur in Boulder County. They are ponderosa pine dwarf-mistletoe (Arceuthobium vaginatum), lodgepole pine dwarf-mistletoe (A. americanum) and limber pine dwarf-mistletoe (A. cyanocarpum), each named for its primary host.

Besides its primary host, each dwarf-mistletoe species attacks the other two pine species as a secondary host (Limber pine is attacked by ponderosa pine dwarf-mistletoe as a secondary host and should not have been listed on the Service in Action leaflet as suitable for planting on ponderosa pine sites.). Only in rare circumstances are other species of trees affected.

Dwarf-mistletoe infects and eventually kills its primary host. Secondary hosts are much more resistant to attack. There are numerous examples of secondary host trees standing in the middle of heavy dwarf-mistletoe infections without becoming infected.

Dwarf-mistletoe control is achieved in forest situations by clear-cutting the infected patch, allowing the stand to regenerate from natural seeds from adjacent stands. If the patch is a large one, the clearcut may have to be completed in several stages so that a seed source remain nearby until the stand regenerates.

In urban settings, or with ponderosa pine dwarf-mistletoe, which is large enough to see easily, it is often feasible to prune dwarf-mistletoe out of infected trees. Due to dwarf-mistletoe's incipient stage, this process must be repeated for at least three consecutive years.

Planting with susceptible tree species before the overstory stand is free of dwarf-mistletoe will result in reinfection. Seedlings are small and not usually infected during the first few years, so if control efforts are continued until all dwarf-mistletoe is gone, an extra two or three years' growth can be obtained by planting after the first year's cleaning.

⁵Stevenson, Douglas J. Dwarf-mistletoe Addendum, Colorado State Forest Service, Boulder, 1994.

APPENDIX III

Soil Series on Cumberland Lode Forest

Fern Cliff Soil Series⁶

The Fern Cliff series is made up of deep, well-drained soils. These soils formed in loamy mixed alluvium on short fans and valley side slopes in the mountain area. Slopes are 15 to 60 percent. Elevations are 6,300 to 8,200 feet. The native vegetation is mainly a forest of ponderosa pine and Douglas-fir with a sparse understory of grass. Annual precipitation is 18 to 24 inches. Mean annual air temperature is 43° to 47° F., and the frost-free season is about 80 to 120 days.

In a representative profile the surface layer is dark grayish-brown stony sandy loam about 3 inches thick. The subsurface layer, about 17 inches thick, is light-gray stony sandy loam. The upper part of the subsoil, about 9 inches thick, is light-gray and yellowish-brown stony sandy loam and sandy clay loam. The sandy clay loam is in thin layers and bands in the sandy loam. The lower part of the subsoil is light brownish-gray and yellowish-brown stony sandy clay loam and sandy loam about 31 inches thick. Below this is light yellowish-brown sandy loam that contains many stones.

Fern Cliff soils have moderate to moderately rapid permeability. Available water capacity for the profile is moderate. Roots can penetrate to a depth of 60 inches or more. Moderate amounts of stone are on the surface and throughout the profile.

Reaction in the upper part of the surface layer is slightly acid, and in the subsurface layer it is medium acid. In the subsoil and substratum it is slightly acid.

These soils are used for pasture, for recreation and forestry, and for homesites.

Typical profile for Fern Cliff stony sandy loam in Fern Cliff-Allens Park-Rock outcrop complex, 15 to 60 percent slopes, located 1,400 feet south and 2,400 feet west of the northeast corner of sec.5, T. 1 N., R. 71 W.:

- O1 - 4 inches to 2, undecomposed organic material, chiefly needles, bark and twigs.
- O2 - 2 inches to 0, partially decomposed organic matter like that of the horizon above.
- A1 - 0 to 3 inches, dark grayish-brown (10YR 4/2) stony sandy loam, very dark brown (10YR 2/2) when moist; strong, fine, crumb structure; soft, very friable; 15 to 20 percent stone; slightly acid; clear, smooth boundary.
- A2 - 3 to 20 inches, light-gray (10YR 7/2) sandy loam, grayish

⁶Moreland, Donald E. and Moreland, Ronald C., Soil Survey of Boulder County Area, Colorado, USDA - Soil Conservation Service, Denver, 1975.

brown (10YR 5/2) when moist; weak, fine, platy structure that parts to moderate fine granular; soft, very friable; 15 to 20 percent stone; medium acid; gradual, wavy boundary.

A&B - 20 to 29 inches, light-gray (10YR 7/2) stony heavy sandy sandy loam, grayish brown (10YR 5/2) when moist; weak, fine, subangular blocky structure; horizon contains thin, discontinuous, yellowish-brown (10YR 5/4) sandy clay loam lamellae and seams that are dark yellowish brown (10YR 4/4) when moist; in some places a soft matrix and very hard lamellae, and in other places a very friable matrix and friable lamellae; thin, nearly continuous clay films on ped faces in lamellae; 20 percent of soil horizon is stone; horizon is slightly acid; diffuse, wavy boundary.

B&A - 29 to 60 inches, yellowish-brown (10YR 5/4) stony clay loam in ½- to 2-inch thick discontinuous lamellae; these lamellae are dark yellowish brown (10YR 4/4) when moist, and interspersed between them is light brownish-gray (10YR 4/2) heavy sandy loam material like that of the horizon above; dark grayish brown (10YR 4/2) when moist; lamellae have moderate, medium, subangular blocky structure, and interspersed material is massive; lamellae are very hard and friable and interspersed material is slightly hard and very friable; lamellae have thin, continuous clay films on ped faces; this horizon is 20 percent stone; slightly acid; gradual, wavy boundary.

C - 60 to 80 inches, light yellowish-brown (2.5Y 6/3) very stony sandy loam, light olive brown (2.5Y 5/3) when moist; massive; slightly hard, very friable; 60 percent stone; slightly acid.

The A1 horizon ranges from 0 to 4 inches in thickness, and in some places it is absent. The A2 horizon ranges from loamy sand to sandy loam in texture. Content of coarse fragments ranges from 5 to 35 percent throughout the solum, but reaches as high as 60 or 70 percent in the C horizon. Depth to bedrock is 60 inches or more.

Fern Cliff-Allens Park-Rock outcrop complex, 15 to 60 percent slopes (FcF). - This complex is made up of about 30 percent Fern Cliff stony sandy loam, about 30 percent Allens Park gravelly sandy loam, and about 20 percent Rock outcrop.

Fern Cliff soils are on mountain slopes and short fans. Allens Park soils are on ridges and side slopes, and on saddles between the ridges. Rock outcrop is throughout the Area, but mainly on ridges.

Included with this complex in mapping are minor areas of Fern Cliff stony loamy sand, Juget soils, Peyton soils, and narrow bands of alluvial soils along drainageways. These included soils make up about 20 percent of each mapped area.

Runoff is medium to rapid on areas of this complex. The erosion hazard is high.

All of the acreage of this complex is woodland. Timber cutting is somewhat limited by the steep slope and the slow growth of trees. Some areas are used for grazing. Many areas are now used as sites for cabins and homes, and for recreational uses and wildlife habitat. (Capability unit VIIe-1, nonirrigated; tree suitability group 1)

Peyton Soil Series⁷

The Peyton series is made up of deep, well-drained soils. These soils formed on upland hills and valley side slopes in weathered loamy and sandy material that has been locally transported. Slopes are 5 to 20 percent. Elevations are from 5,800 to 7,500 feet. The vegetation is mainly tall grasses and, in places, ponderosa pine. Annual precipitation is 18 to 24 inches. Mean annual air temperature is 44° to 48° F., and the frost-free season is about 80 to 120 days.

In a representative profile the surface layer is dark-gray very gravelly loamy sand about 11 inches thick. The subsoil, about 32 inches thick, is brown gravelly sandy clay loam that grades to gravelly coarse sandy loam. The substratum extending to a depth of 60 inches or more is pale-brown gravelly sandy loam. Soil reaction is neutral.

Peyton soils have moderate permeability. Available water capacity for the profile is moderate. Roots can penetrate to a depth of 40 to 60 inches or more.

These soils are used for pasture and homesites.

Representative profile of Peyton very gravelly loamy sand in an area of Peyton-Juget very gravelly loamy sands, 5 to 20 percent slopes, is located 1,640 feet north and 100 feet east of southwest corner of sec. 11, T. 1 N., R. 71 W.:

- A1 - 0 to 11 inches, dark-gray (10YR 4/1) very gravelly loamy sand, very dark brown (10YR 2/2) when moist; weak, fine, granular structure; slightly hard, very friable; neutral; clear, smooth boundary.
- B1t - 11 to 16 inches, brown (10YR 5/3) gravelly sandy clay loam, dark brown (10YR 3/3) when moist; weak, medium,

⁷Moreland, Donald E. and Moreland, Ronald C. Soil Survey of Boulder County Area, Colorado, USDA - Soil Conservation Service, Denver, 1975.

subangular blocky structure; hard, very friable; thin, patchy clay films on ped faces; neutral; clear, smooth boundary.

B2t - 16 to 30 inches, brown (10YR 5/3) gravelly sandy clay loam, brown (10YR 4/3) when moist; moderate, medium subangular blocky structure; very hard, friable; thin, nearly continuous clay films on ped faces; neutral; clear, smooth boundary.

B3t - 30 to 43 inches, brown (10YR 5/3) gravelly coarse sandy loam, brown (10YR 4/3) when moist; weak, medium and coarse subangular blocky structure; hard, friable; thin, patchy clay films on ped faces; neutral; gradual, smooth boundary.

C - 43 to 60 inches, pale-brown (10YR 6/3) gravelly sandy loam, brown (10YR 5/3) when moist; massive; very hard, friable; neutral.

The A horizon ranges from 6 to 12 inches in thickness and from heavy loamy sand to sandy loam in texture. Structure of the B2 horizon ranges from weak and moderate subangular blocky to weak prismatic. The C horizon ranges from sandy loam to loamy sand in texture. Amount of coarse fragments in the soil ranges from 10 to 25 percent, although in places the surface layer contains up to 60 percent coarse fragments.

Peyton-Juget very gravelly loamy sands, 5 to 20 percent slopes (PgE). - This complex is on the uplands.... These are open park-like areas mainly more than 20 acres in size. This complex consists of about 65 percent Peyton very gravelly loamy sand and about 20% Juget very gravelly loamy sand. Peyton very gravelly loamy sand is in the main part of the area. Juget very gravelly loamy sand is on the edges of the area and on narrow ridges extending into the areas. The Peyton soil has the profile described as representative of the Peyton series. The Juget soil has a profile much like that described as representative of the Juget series, but the surface layer is very gravelly loamy sand.

Included with this complex in mapping are small areas of rock outcrop and a few small areas of Allens Park soils. Also included along some drainageways are areas of a deep, dark-colored soil formed in alluvium. A few small areas around lakes or ditches have a seasonal high water table. The included soils make up about 15% of each mapped area.

Runoff is slow to medium on this complex. The erosion hazard is moderate to high. Tillage of most areas is limited by the amount of gravel.

All the acreage of this complex is used for pasture, recreational uses and wildlife habitat. Some areas are used for homesites. (Capability unit VIIs-1, nonirrigated; tree suitability group 2)

APPENDIX IV

Glossary

SILVICULTURE TERMINOLOGY^a**Advance Regeneration (Reproduction) syn****Advance Growth**

Seedlings or saplings that develop or are present in the understory.

Afforestation

Establishment of a forest or stand in an area not previously forested.

Age Class (Cohort)

A distinct aggregation of trees originating from a single natural disturbance or regeneration cutting.

Artificial Regeneration (Reproduction)

Creation of a new age class by renewal of a tree crop by direct seeding, or by planting seedlings or cuttings.

Burning, Prescribed

The application of fire, usually under existing stands and under specific conditions of weather and fuel moisture, in order to control vegetation to meet goals of silviculture or hazard reduction.

Cleaning

A release treatment made in an age class not past the sapling stage in order to free the favored trees from less desirable individuals of the same age class which overtop them or are likely to do so.

Cohort

See Age Class.

Composition, Stand

The proportion of each tree species in a stand expressed as a percentage of either the total number, basal area, or volume of all tree species in the stand.

Crop Tree

Any tree that is selected to become a component of a future final harvest.

Crown

The part of a tree or woody plant bearing live branches and foliage.

Crown Class

A class of tree based on crown position relative to the crowns of adjacent trees.

Codominant

Trees with crowns forming the general level of the main canopy in even-aged groups of trees, receiving full light from above and comparatively little from the sides.

^aLoftis, David L., "Silviculture Terminology" SAF Silviculture Working Group Newsletter, Society of American Foresters, Bethesda, Maryland, 1993, pp. 1-3.

Dominant

Trees with crowns extending above the general level of the main canopy of even-aged groups of trees, and receiving full light from above and partly from the sides.

Intermediate

Trees with crowns extending into the lower portion of the main canopy of even-aged groups of trees, but shorter in height than codominants. They receive little direct light from above and none from the sides.

Overtopped (Suppressed)

Trees of varying levels of vigor that have their crowns completely covered by the crowns of one or more neighboring trees.

Crown Cover

The ground area covered by the crowns of trees or woody vegetation as delimited by the vertical projection of crown perimeters and commonly expressed as a percentage of total ground area (syn. Canopy Cover).

Crown Density

The amount, compactness, or depth of foliage of the crowns of trees and/or shrubs.

Cutting Cycle

The planned interval between partial harvests in an uneven-aged stand (See Thinning Interval.).

Even-Aged Stand

A stand of trees containing a single age class in which the range of tree ages is usually less than 20% of rotation.

Harvesting Method

A cutting method by which a stand is harvested. Emphasis is on meeting logging requirements rather than silvicultural objectives (See Regeneration Methods.).

Improvement Cutting

A cutting made in a stand past the sapling stage primarily to improve composition and quality by removing less desirable tree species.

Ingrowth

Trees that during a specified period have grown past an arbitrary lower limit of (usually) diameter or height. Ingrowth is usually measured as basal area or volume per unit area.

Intermediate Treatments (Tending)

A collective term for any treatment designed to enhance growth, quality, vigor, and composition of the stand after establishment of regeneration and prior to final harvest (See Tending, Stand Improvement.).

Liberation

A release treatment made in a stand not past the sapling stage in order to free the favored trees from competition of older, overtopping trees.

Natural Regeneration

An age class created from natural seeding, sprouting, suckering, or layering.

Nurse Tree (Nurse Crop)

A tree, group or crop of trees, shrubs, or other plants, either naturally occurring or introduced, used to nurture or improve the form of a more important tree or crop during youth by protecting it from frost, insolation, or wind.

Overstory Removal

The cutting of trees comprising an upper canopy layer in order to release trees or other vegetation in an understory.

Precommercial Thinning (PCT)

A thinning that does not yield trees of commercial value, usually designed to improve crop spacing.

Regeneration (Reproduction) Method

A cutting method by which a new age class is created. The major methods are Clearcutting, Seed Tree, Shelterwood, Selection, and Coppice (See Harvesting Method.).

Even-Aged Methods**Clearcutting**

A method of regenerating an even-aged stand in which a new age class develops in a fully-exposed microclimate after removal, in a single cutting, of all trees in the previous stand. Regeneration is from natural seeding, direct seeding, planted seedlings, and/or advance reproduction. Harvesting may be done in groups or patches (**Group or Patch Clearcutting**), or in strips (**Strip Clearcutting**). In the Clearcutting System, the management unit or stand in which regeneration, growth, and yield are regulated consists of the individual clearcut stand (See Group Selection).

Clearcutting with Reserves

A clearcutting method in which varying numbers of reserve trees are not harvested to attain goals other than regeneration.

Seed Tree

An even-aged regeneration method in which a new age class develops from seedlings that germinate in fully-exposed micro-environments after removal of all the previous stand except a small number of trees left to provide seed. Seed trees are removed after regeneration is established.

Seed Tree with Reserves

A seed tree method in which some or all of the seed trees are retained after regeneration has become established to attain goals other than regeneration.

Shelterwood

A method of regenerating an even-aged stand in which a new age class develops beneath the partially-shaded micro-environment provided by residual trees. The sequence of treatments can include three distinct types of cuttings: 1) an optional preparatory harvest to enhance conditions for seed production; 2) an establishment harvest to prepare the seed bed and create a new age class; and 3) a removal harvest to release estab-

lished regeneration from competition with the overwood. Harvesting may be done uniformly throughout the stand (**Uniform Shelterwood**), in groups or patches (**Group Shelterwood**), or in strips (**Strip Shelterwood**).

Shelterwood with Reserves

A variant of the Shelterwood Method in which some or all of the shelter trees are retained, well beyond the normal period of retention, to attain goals other than regeneration. The resulting stand may be two-aged or tend towards an uneven-aged condition as a consequence of both an extended period of regeneration establishment and the retention of reserve trees that may represent one or more age classes.

Two-Aged Methods

Methods designed to maintain and regenerate a stand with two age classes (See Shelterwood with Reserves and Coppice with Reserves.).

Uneven-Aged (Selection) Methods

Methods of regenerating a forest stand, and maintaining an uneven-aged structure, by removing some trees in all size classes either singly, in small groups, or in strips.

Group Selection

A method of regenerating uneven-aged stands in which trees are removed, and new age classes are established, in small groups. The maximum width of groups is approximately twice the height of the mature trees, with small openings providing micro-environments suitable for tolerant regeneration and the larger openings providing conditions suitable for more intolerant regeneration. In the Group Selection system, the management unit or stand in which regeneration, growth, and yield are regulated consists of a landscape containing an aggregation of groups (See Clearcutting.).

Single Tree Selection

A method of creating new age classes in uneven-aged stands in which individual trees of all size classes are removed more-or-less uniformly throughout the stand to achieve desired stand structural characteristics.

Coppice Methods

Methods of regenerating a stand in which the majority of regeneration is from stump sprouts or root suckers.

Coppice

A method of regenerating a stand in which all trees in the previous stand are harvested and the majority of regeneration is from sprouts or root suckers.

Coppice with Reserves

A coppice method in which reserve trees are retained to attain goals other than regeneration. The method normally creates a two-aged stand.

Regeneration (Reproduction) Period

The time between the initial regeneration cutting and the successful re-establishment of a new age class by natural means, planting, or direct seeding.

Regular Uneven-Aged (Balanced) Stand

A stand in which three or more distinct age classes occupy approximately equal areas and provide a balanced distribution of diameter classes.

Release (Release Operation)

A treatment designed to free young trees from undesirable, usually overtopping, competing vegetation. Treatments include cleaning, liberation, and weeding (See Stand Improvement.).

Reserve Trees

Trees pole-sized or larger, retained after the regeneration period under the Clearcutting, Seed Tree, Shelterwood, or Coppice Methods. syn. Standards.

Salvage Cutting

The removal of dead trees or trees being damaged or killed by injurious agents other than competition, to recover value that would otherwise be lost.

Sanitation Cutting

The removal of trees to improve stand health and to reduce actual or anticipated spread of insects and disease (See Stand Improvement.).

Sapling

A tree, usually young, that is larger than a seedling but smaller than a pole.

Silviculture

The art and science of controlling the establishment, growth, composition, health, and quality of forests and woodlands. Silviculture entails the manipulation of forest and woodland vegetation in stands and on landscapes to meet the diverse needs and values of landowners and society on a sustainable basis.

Silvicultural System

A planned process whereby a stand is tended, harvested, and re-established. The system name is based on the number of age classes (See Even-Aged, Two-Aged, Uneven-Aged.), and/or the regeneration method used (See Clearcutting, Seed Tree, Shelterwood, Selection, Coppice, Coppice with Standards.).

Site Class

A classification of site quality, usually expressed in terms of ranges of dominant tree height at a given age or potential mean annual increment at culmination.

Site Preparation

A hand or mechanized manipulation of a site designed to enhance the success of regeneration. Treatments may include chopping, discing, bedding, raking, burning and scarifying. All treatments are designed to modify the soil, litter, vegetation and to create microclimate conditions conducive to the establishment and growth of desired species.

Site Quality (Productivity)

The productive capacity of a site, usually expressed as volume production of a given species.

Size Classes

Tree sizes recognized by distinct ranges, usually of diameter or height.

Stand

A contiguous group of trees sufficiently uniform in age class distribution, composition, and structure, and growing on a site of sufficiently uniform quality, to be a distinguishable unit (See Mixed, Pure, Even-Aged, and Uneven-Aged Stands.).

Mixed Stand

A stand in which there is a mixture of species.

Pure Stand

A stand composed of essentially a single species.

Stratified Mixture

A stand in which different species occupy different strata of the total crown canopy.

Stand Density

A quantitative, absolute measure of tree occupancy per unit area in such terms as numbers of trees, basal area, or volume.

Stand Improvement

A term comprising all intermediate cuttings made to improve the composition, structure, condition, health and growth of even- or uneven-aged stands.

Stocking

An indication of growing-space occupancy relative to a pre-established standard. Common inducers of stocking are based on percent occupancy, basal area, Relative Density, and Crown Competition Factor.

Stratum (Canopy Layer)

A distinct layer of vegetation within a forest community.

Tending

See Intermediate Treatments.

Thinning

A cutting made to reduce stand density if trees primarily to improve growth, enhance forest health, or to recover potential mortality.

Crown Thinning (Thinning From Above, High Thinning)

The removal of trees from the dominant and codominant crown classes in even-aged stands, or in even-aged groups within uneven-aged stands, in order to favor the best trees of those same crown classes.

Free Thinning

The removal of trees in even-aged or uneven-aged stands to control stand spacing and favor desired trees using a combination of thinning criteria without regard to crown position.

Low Thinning (Thinning From Below)

The removal of trees from the lower crown classes to favor those in the upper crown classes.

Mechanical Thinning (Geometric Thinning)

The thinning of trees in either even- or uneven-aged stands involving removal of trees in rows, strips, or by using fixed spacing intervals.

Selection Thinning (Dominant Thinning)

The removal of trees in even-aged stands, or in even-aged groups within even-aged stands, in the dominant crown class in order to favor the lower crown classes.

Thinning Interval

The period of time between successive thinning entries, usually used in connection with even-aged stands (See Cutting Cycle.).

Tolerance, Shade

The relative capacity of a plant to become established and grow in the shade.

Two-Aged Stand

A stand composed of two distinct age classes that are separated in age by more than 20 percent of rotation.

Undercutting (Root Pruning)

The root pruning of seedlings in a nursery bed.

Uneven-Aged Stand

A stand of trees of three or more distinct age classes, either intimately mixed or in small groups.

Uneven-Aged System

A planned sequence of treatments designed to maintain and regenerate a stand with three or more age classes (See Single Tree Selection, Group Selection.).

Weeding

A release treatment in stands not past the sapling stage that eliminates or suppresses undesirable vegetation regardless of crown position.

Wrenching

The disturbance of seedling roots in a nursery bed (e.g. with a tractor-drawn blade) with the objective of stimulating the development of a fibrous root system.

APPENDIX V
Bibliography

ADDITIONAL READING

Bilodeau, Sally W., et. al., "Geology of Boulder, Colorado, United States of America," Bulletin of the Association of Engineering Geologists, Vol. XXIV, No. 3, 1987, pp. 289-332.

Dilworth, J. R., Log Scaling and Timber Cruising, Oregon State University Book Stores, 1970.

Dunfee, Chuck, et. al., S-390 Fire Behavior, Boise Interagency Fire Center, Boise, Idaho, 1981.

Hawksworth, Frank G. and Wiens, Delbert, Biology and Classification of Dwarf-mistletoes (Arceuthobium), Agriculture Handbook No. 401, Rocky Mountain Forest and Range Experiment Station, Forest Service, Fort Collins, Colorado, 1972.

Hoover, Robert L. and Wills, Dale L., Managing Forested Lands for Wildlife, Colorado Division of Wildlife in Cooperation with USDA Forest Service, Denver, 1984.

Moreland, Donald C. and Moreland, Ronald E., Soil Survey of Boulder County Area, Colorado, United States Department of Agriculture, Soil Conservation Service, 1975.

Shigo, Alex L., A New Tree Biology, Shigo and Trees Associates, Durham, New Hampshire, 1986.

Spellenberg, Richard, The Audubon Society Field Guide to North American Wildflowers, Western Region, Alfred A. Knopf, New York, 1979.

United States Forest Service, Silvics of Forest Trees of the United States, Agricultural Paper # 271.