

THESIS

THE 2006 WARM FIRE: EFFECTS ON HABITAT AND PREY SPECIES  
OF THE NORTHERN GOSHAWK

Submitted by

Jeffrey S. Lambert

Department of Forest and Rangeland Stewardship

In partial fulfillment of the requirements

For the Degree of Master of Science

Colorado State University

Fort Collins, Colorado

Spring 2015

Master's Committee:

Advisor: Dan Binkley

Richard Reynolds

Julie Savidge

Copyright by Jeffrey Scott Lambert 2015

All Rights Reserved

## ABSTRACT

### THE 2006 WARM FIRE: EFFECTS ON HABITAT AND PREY SPECIES OF THE NORTHERN GOSHAWK

Reductions in the frequency of fire in Southwestern ponderosa pine and mixed-conifer forests since initiation of forest management early in 20<sup>th</sup> century changed the composition and structure of the forest habitats of the northern goshawks (*Accipiter gentilis*), a food-limited species of conservation concern, and the birds and mammals it feeds on. A conservation strategy for the species in the goshawk's food web in these forest types recommends restoring the mix of predator and prey habitats that historically characterized these forests—characteristics that were sustained by frequent low-severity surface fire. Thus, the effects of fire severities (high- and low-severity) and lack of fire on today's habitats may influence the abundances of bird and mammal goshawk prey. . The 2006 Warm Fire burned 235 km<sup>2</sup> of ponderosa pine and mixed-conifer forest on the North Kaibab Ranger District in northern Arizona in late June and early July 2006. Forest habitat metrics such as live tree and snag densities, cone production, canopy and ground cover estimates were collected from 2007-2010 on 60 0.5km transects to compare the effects of high- and low-severity fire and no fire on a suite of 13 important prey species of the goshawk. I describe habitat changes resulting from the different fire severities in ponderosa pine and mixed conifer forests. I estimated abundances for 13 bird and mammal prey species in forests burned by different fire severities and tested predictive models designed to gain an understanding which habitat characteristics, affected by fire, best predicted individual bird and mammal abundances. Red squirrels and golden-mantled ground squirrels showed the most sensitivity to fire, while chipmunks were evenly distributed across fire severities. Hairy woodpeckers and northern flickers, in contrast, benefitted from high-severity fire, due to increased snags, cavity nesting opportunities, and foraging. American robins and Steller's jays were also evenly distributed across fire severities. High-severity fire had a significant impact on forest ecosystems. Changes in forest structure were found to be detrimental to

some species while creating short- and possibly intermediate-term benefits for others. Lowering the risk of high-severity fire by restoring composition and structure should also protect the habitats of goshawk and the prey species most sensitive to fire. The effects of low-severity fire were mixed, suggesting that it may be possible to return fire to ecosystem without a significant impact to many birds and mammals in the short term. Frequent low-severity fire could also help to maintain sub-canopy and understory grass / shrub openings by removing excess growth and ground debris.

## ACKNOWLEDGEMENTS

I first extend my appreciation to my advisor, Dr. Dan Binkley of the Department of Ecosystem Science and Sustainability and my Forest Service supervisor, Dr. Richard Reynolds of the Rocky Mountain Research Station. Richard encouraged me to pursue this research opportunity, and has supported me throughout my program as an employer, mentor, and friend. Dan provided me with an opportunity to formalize this research and I would like to thank him for his encouragement, professional advice, life lessons, and the significant amount of time he invested in my academic development. I would also like to thank Dr. Julie Savidge of the Department of Fish, Wildlife, and Conservation Biology for her academic guidance and for serving on my graduate committee. I owe much to Richard, Dan, and Julie for their support during my program. I would also like to thank Dr. Scott Baggett and Ben Bird for their statistical consultations and advice throughout this project.

This study would not have been possible without the support of many field technicians who assisted me with data collection and logistics on the Kaibab Plateau. In particular I would like to thank Wyatt Nimitz (2007) and Carrie Lambert (2008-2010) for waking up early many mornings and assisting me with bird and mammal surveys. I would also like to thank Alison Cofer, Heather Danaceau, Mary Francis, Tara Hatfield, Jean Lawrence, Katie Parmentier, Mike Price, John Seyfried, Leath Tonino, and Tim Webber for their support throughout this study. I appreciate all of their hard work and enthusiasm over the years.

I express my deepest gratitude to my family and friends for their encouragement and support. I am especially grateful to my wife Carrie for her love and encouragement. Carrie also provided me with invaluable assistance collecting data throughout the study. I am also thankful for my son Riley who provided me with the proper perspective about what is important in life.

This project was funded and supported by Region 3 of the U.S. Forest Service, the Rocky Mountain Research Station, and the Joint Fire Science Program. Housing and logistics during the field season were provided by the North Kaibab Ranger District. I thank you all.

## TABLE OF CONTENTS

ABSTRACT.....	ii
ACKNOWLEDGEMENTS.....	iv
LIST OF TABLES.....	vii
LIST OF FIGURES.....	viii
INTRODUCTION.....	1
METHODS.....	6
STUDY AREA.....	6
SAMPLING DESIGN.....	7
VEGETATION SAMPLING.....	9
PREY SAMPLING.....	10
DATA ANALYSIS.....	11
RESULTS.....	14
TREE AND SNAG DENSITY AND CANOPY COVER.....	14
GROUND COVER.....	15
SAPLINGS COUNTS.....	16
CONE PRODUCTIVITY.....	16
BIRD AND MAMMAL ABUNDANCE.....	17
PREDICTIVE MODELS.....	17
DISCUSSION.....	20

CONCLUSIONS.....	25
REFERENCES.....	43
APPENDICES.....	49

LIST OF TABLES

TABLE 1-TRANSECT CLASSIFICATION BY FOREST TYPE AND FIRE SEVERITY.....28  
TABLE 2-PRE- AND POST-FIRE TRANSECT CLASSIFICATION.....29  
TABLE 3-PREY SPECIES ABUNDANCE RELATED TO HABITAT AND FIRE SEVERITY.....30-33  
TABLE 4-TRANSECT METRICS BY FOREST TYPE AND FIRE SEVERITY.....34  
TABLE 5-SPECIES COUNT DATA.....35  
APPENDIX A.....50-51  
APPENDIX B.....52-55  
APPENDIX C.....56  
APPENDIX D..... 57

## LIST OF FIGURES

FIGURE 1-KAIBAB PLATEAU STUDY AREA.....	36
FIGURE 2.1-TOTAL TREE DENSITY.....	37
FIGURE 2.2-TREE DENSITY >20CM DBH (LARGE).....	37
FIGURE 3-CANOPY COVER.....	38
FIGURE 4.1-TOTAL SNAG DENSITY.....	39
FIGURE 4.2-SNAG DENSITY >20CM DBH (LARGE).....	39
FIGURE 5.1-OCCULAR ESTIMATES OF MIXED-CONIFER GROUND COVER.....	40
FIGURE 5.2-OCCULAR ESTIMATES OF PONDEROSA PINE GROUND COVER.....	40
FIGURE 6.1-MIXED CONIFER SAPLING COUNTS.....	41
FIGURE 6.2-PONDEROSA PINE SAPLING COUNTS.....	41
FIGURE 7.1-MIXED CONIFER CONE PRODUCTIVITY.....	42
FIGURE 7.2-PONDEROSA PINE CONE PRODUCTIVITY.....	42

## INTRODUCTION

Wildfire is a major disturbance in ecosystems across the United States (Agee 1993) and is an important driver of landscape change (Fulé et al. 1997). In the Southwestern United States historic fire regimes included frequent low-severity surface fires in ponderosa pine (*Pinus Ponderosa*) and low elevation dry (warm, dry) mixed-conifer forests (White and Vankat 1993, Fulé et al 2000), less frequent mixed-severity fires in mid-elevation wet (cool, mesic) mixed-conifer forests, and infrequent, high-severity fires in higher elevation spruce-fir forests (White and Vankat 1993). In Arizona and New Mexico, pure ponderosa pine and mixed-conifer forests occupy nearly 2.5 and 1 million hectares of forest area, respectively (Brown 1994, Korb et al. 2013). As a dominant species in Southwestern forests, ponderosa pine can influence forest ecosystems and fire behavior (Moore et. al. 1999). As a consequence of frequent low-severity surface fire, ponderosa pine forests in northern Arizona have been described as low-density, open forests consisting of aggregations of old pine trees interspersed with aggregations of younger pine trees in a grass/shrub matrix (White 1985), while mixed-conifer forests have been classified into two sub-categories: warm/dry and cool/moist forests (Romme et al. 2009). Dry mixed-conifer forests are typically located at lower elevations and are dominated by ponderosa pine and Douglas-fir, but can include white fir (*Abies concolor*) and aspen (*Populus tremuloides*) (Korb et al. 2013). Historically, these forests were characterized by large old trees, relatively open with low trees densities, and heterogeneous patches of uneven-aged trees (Binkley et al. 2008, Fule et al. 2009). Snag, log, and woody debris abundance was thought to be similar to ponderosa pine forests (Moore et al. 2004). Cool/moist mixed-conifer forests, in comparison, are found at higher elevations and include a greater abundance of Douglas-fir (*Pseudotsuga menziesii*) and other conifers adapted to these conditions (Romme et al. 2009). Cool/moist forests are more spatially heterogeneous regarding patch sizes that differ in tree ages and densities than ponderosa pine and dry mixed-conifer forest types due to a mixed-severity fire regime (Binkley et al. 2008, Fule et al. 2009).

After more than a century of extensive fire exclusion starting in the early 1900's, along with other management practices, ponderosa pine and mixed-conifer forest have been altered by increases in tree densities, a shift to more shade-tolerant tree species, accumulations of small trees, accelerated mortality of old trees, and diminished understory (herbs, shrubs) productivity (Covington and Moore 1994, Fule et al. 2009, Churchill et al. 2013). The buildup of fuel has made these forests vulnerable to high-severity crown fires that not only kill trees but exposes landscapes to erosion, exotic plant invasion, and other negative consequences (Covington et al. 1994, Griffis et al. 2001).

Historically, the fire return interval in Southwestern ponderosa pine ranged from 2-25 years (Swetnam and Baisan 1996, Heyerdahl et al. 2001). Frequent fires in this ecosystem favored shade intolerant, fire resistant tree species, minimal fuel build up, and open forest conditions that promoted productive understory growth (Fule and Laughlin 2007). Pre-settlement mixed-conifer forests (both dry and wet mixed-conifer) were shaped by mixed-severity fire regimes consisting of areas of surface and areas of stand-replacing fires that could occur in the same event. These ecosystem types contained trees of various ages and species composition resulting from variation in fire intensity, frequency, and area burned (Fule et al 2003). Pre-settlement dry mixed-conifer forests were historically dominated by ponderosa pine, Douglas-fir and white fir and had reported fire return intervals of 4-14 and 18-32 years (similar to ponderosa pine forests) in two studies (White and Vankat 1993, Brown et al. 2001, Korb et al. 2013). Historically, wet mixed-conifer forests had sub-decadal to century long fire return intervals upwards of 200 years (White and Vankat 1993, Romme et al. 2009). Wet mixed-conifer forests were characterized by cooler mesic conditions, increased tree density and canopy cover, infrequent intense fires, and fire intolerant overstory trees (White and Vankat 1993). Because of long fire return intervals in high elevation Engelmann spruce-sub-alpine fir landscapes, fire suppression has had little effect on these ecosystems (White and Vankat 1993). Due to more frequent surface fire, dry mixed-conifer forests were more open with groups of small trees similar to ponderosa (Fule et al. 2003, Romme et al. 2009). Moist higher elevation forests experienced less frequent moderate to high severity fires that created large patches (cohorts) of similar-aged trees (Fule et al. 2003, Romme et al. 2009).

In many instances alteration of the historic fire regimes, especially in ponderosa pine and dry mixed-conifer forests, created conditions that currently favor more frequent and large high severity wildfires. Forest ecosystems that were typically shaped by low-intensity and mixed severity fires have been transformed into ones that are susceptible to high-severity, stand-replacing crown fires.

The northern goshawk's (*Accipiter gentilis*), a U.S. Forest Service sensitive species of special conservation concern, response to habitat changes due to forest management and disturbances such as fire is of particular interest throughout Western U.S. forests. The goshawk is a large forest dwelling raptor that occupies mature temperate and boreal forests throughout the Holarctic (Squires and Reynolds 1997). Goshawk home ranges vary from 15-30 km<sup>2</sup> and are generally "fixed" in their locations, with minor yearly fluctuations due to their nesting and foraging requirements (Reynolds et. al. 1994). Goshawks occupy a variety of forest types that are composed of different vertical and horizontal structures throughout their N.A. range. Typically, goshawks in the Southwest U.S. nest in mature trees with large open branches and hunt in the sub-canopy space, along forest edges, and in herbaceous and shrub openings (Reynolds et al. 1992, 2006, 2013). Nest sites are characterized by mature and old trees and relatively dense canopies that provide nest protection from predators and environmental extremes. Nesting sites comprise < 1% (0.12 km<sup>2</sup>) of the total goshawk home range and have low vegetative structural diversity compared to forest conditions in their much larger home ranges, which is used for foraging, roosting, and by young hawks post-fledging (Reynolds et. al. 2006). On the Kaibab Plateau mature and old canopy forests with open understories are used more often than any other forest types by breeding goshawks (Bright-Smith and Mannan 1994, Reynolds et al. 2006). Adapted to foraging in the lower vegetation column of forests, a goshawk's hunting ability is thought to be impaired when forest understories become filled with small trees or tall dense brush (Reynolds and Meslow 1982) due to a lack of fire and other ecological processes. Loss of nesting and foraging habitat due to high-severity fire may also decrease survival and nesting productivity through loss of prey habitats and lowered prey abundance. Due to the complexity of forest fire regimes, most of which have been altered throughout the goshawks

geographic range, understanding fire effects on its food webs, on its habitats, its reproduction, and its population viability is an important conservation concern across the southwestern United States.

Goshawks on the Kaibab Plateau have a diverse diet, foraging upon a suite of more than 20 species of ground and tree squirrels, rabbits and hares, woodpeckers, grouse, turkeys, and passerines (Reynolds 1983, Reynolds and Meslow 1984, Reynolds et al. 1994, Squires and Reynolds 1997, Salafsky et al. 2005, 2007). Prey counts conducted from 1999-2002 suggested that prey species abundance is a factor limiting goshawk reproduction on the Kaibab Plateau (Salafsky 2004, Salafsky et al. 2005, 2006). Quantifying the effects of fire on bird and mammal prey of goshawks is difficult since there is a high degree of species-specific responses to fire and variation in detectability. Bock and Block (2005) found that season, scale, size and severity of fire influenced bird and mammal responses to fire disturbances. Saab et al. (2002) noted that fire can create unique nesting and foraging habitats by increasing diversity in vegetation type, canopy closure, and climate. Studying the effects of fire on the birds and mammals in the goshawk food web is important because goshawks are both food limited (Salafsky et al. 2005, 2007) and a species of special concern in the Southwest forests. Uniquely suited to preying on a varied suite of birds and mammals in Kaibab plateau forests, goshawk productivity has been closely tied to annual fluctuations in prey abundance (Salafsky 2004, Salafsky et al. 2005, 2006), and prey abundance depends of availabilities and productivities of varied habitats that support the suite of prey.

From 8 June 2006 to containment on 4 July 2006, the Warm Fire, which started as a lightning strike, burned 235 km<sup>2</sup> of ponderosa pine and mixed-conifer forest on the North Kaibab Ranger District in northern Arizona. For 18 days post-ignition, the Warm Fire met District and Federal Wildland Fire Management criteria to be used as a wildland fire-use fire, during which time approximately 77 km<sup>2</sup> were burned at low to moderate-severity. On 25 June 2006 weather conditions changed and the fire exceeded its maximum manageable area and active suppression was initiated. An additional 158 km<sup>2</sup> of forest burned between 25 June and 4 July 2006, before full containment. The majority of the forest that burned after 25 June 2006 had moderate-high and high-severity fire damage, while some small patches dispersed throughout the fire burned at low and moderate-low severity. In total, approximately 64% of the Warm

Fire (12% of the North Kaibab Ranger District) burned at moderate-high to high-severity and 36% (7% of the North Kaibab Ranger District) burned at moderate-low to low-severity, resulting in some level of alteration to almost 20% of the forests on the ranger district.

Changes in live forest structure caused by the Warm Fire provided an opportunity to explore the impact of fire on the composition and structure of habitats and species composition and abundance of bird and mammal prey of goshawks. To evaluate how fire severity affected the flora and fauna of the Kaibab Plateau, I conducted surveys of (1) the vegetation composition and structure of post-Warm Fire habitats and (2) estimated bird and mammal abundances in ponderosa pine and mixed-conifer forests on transects in areas unburned, burned at low-severity, and burned at high-severity fire in the Warm Fire.

*Study Objectives:*

1. Describe and compare the vegetation composition and structure in relation to fire severity (no burn, low-severity, and high-severity) in the ponderosa pine and mixed-conifer forest types.
2. Estimate the composition and abundance of bird and mammal species by forest type (ponderosa pine, mixed-conifer) and by fire severity.
3. Model the effects of selected forest metrics and fire severity on birds and mammals and determine the best models for predicting species abundance.

## METHODS

### Study Area

The 1,285 km<sup>2</sup> study area (Figure 1) includes the North Kaibab Ranger District of the Kaibab National Forest above 2,182 m elevation on the Kaibab Plateau in northern Arizona. The eastern, southern, and western edges of the plateau are formed by steep slopes, creating a distinct boundary between the forests and surrounding shrub-steppe ecosystems. The northern edge of the Plateau slopes gradually through a broad ecotone from shrub-steppe ecosystems (1,750 m elevation) to spruce-fir forests at the highest elevations (maximum 2,800 m). Mixed-conifer forests occupy about 275 km<sup>2</sup> (20%) of the study area at the highest elevations, ponderosa pine (*Pinus ponderosa*) forests occupy 714 km<sup>2</sup> (55%) at mid-elevation ranges, and pinyon-juniper (*Pinus edulis*, *Juniperus spp.*) woodlands occupy 106 km<sup>2</sup> (<10%) at the lower elevations. Quaking aspen (*Populus tremuloides*) forests cover about 112 km<sup>2</sup>, interspersed among the other forest woodland types (Joy 2002), comprising up to 14% of the tree basal area on the Plateau (Binkley et al. 2006). Ponderosa pine forests of the Kaibab Plateau are characterized by relatively low tree diversity (White 1985). Common ponderosa pine understory grasses included: *Bromus spp.*, wild rye (*Elymus elymoides*), mountain muhly (*Mulenbergia montana*), and *Poa spp.*; understory forbs include: *Antennaria spp.*, *Chenopodiaceae spp.*, Wright's deervetch (*Lotus wrightii*), and *Lupinus spp.*; and understory shrubs include: Oregon grape (*Berberis repens*), Fendler's ceanothus (*Ceanothus fendleri*), and New Mexican locust (*Robinia neomexicana*). Mixed-conifer forests of the Kaibab Plateau have higher tree diversity than ponderosa pine (White and Vankat 1993). Species commonly found in mixed-conifer forests include: white fir (*Abies concolor*), subalpine fir (*Abies lasiocarpa*), Englemann spruce (*Picea engelmannii*), ponderosa pine, quaking aspen, and Douglas fir (*Pseudotsuga menziesii*). Common mixed-conifer understory grasses include: *Bromus spp.*, wild rye, *Festuca spp.*, and *Poa spp.*; understory forbs include: *Antennaria spp.*, *Chenopodiaceae spp.*, wild strawberry, *Fragaria vesca*, and *Lupinus spp.*; and understory shrubs include: Oregon grape, Fendler's ceanothus, common juniper (*Juniperus communis*), and *Ribes spp.*

Almost all of the North Kaibab Ranger District had received tree harvests prior to the Warm Fire. Organized tree harvests on the NKRD began in the early 1900s and were primarily limited to cutting dead and dying trees (single-tree selection harvest) (Sesnie and Bailey 2003). In the 1960s, small patch cuts began in higher elevation mixed-conifer forests, but were discontinued in the early 1970s. In the 1980s intensive stand management using shelterwood / seed-tree cuts occurred to create even-aged forest patches in ponderosa and mixed-conifer forests, but by the early 1990s this protocol was discontinued (Sesnie and Bailey 2003). While pre-1960s selection harvests occurred over much of the NKRD, intensive stand management occurred in scattered patches on the Kaibab plateau (Sesnie and Bailey 2003).

### **Sampling Design**

To evaluate the effects of the 2006 Warm Fire on habitat composition and structure, and associated prey species, I measured vegetation composition and structure and conducted bird and mammal counts on 60 transects placed in six strata (unburned, low-severity, and high-severity fire in ponderosa pine and mixed-conifer forests that had previously been treated with two different silvicultural prescription, single-tree harvested and shelter-wood/seed-tree harvest) over 4 post-fire years (2007-10). With vegetation burn severity maps (North Kaibab Ranger District, NKRD 2007), I aggregated moderate-high and high-severity burn categories into one high-severity strata, and moderate-low and low-severity burn categories into one low-severity strata (North Kaibab Ranger District, NKRD 2007). I sampled a total of 60 0.5km transects (Table 1); 40 in unburned forest (23 in ponderosa pine, 17 in mixed-conifer), 8 burned by low-severity fire (4 in ponderosa pine, 4 in mixed-conifer), and 12 burned by high-severity fire (3 in ponderosa pine, 9 in mixed-conifer).

I selected 49 of my 60 transects from 60 previously-established transects used in a study of co-variation between prey abundance and goshawk fecundity (Table 2) (Salafsky 2004). I randomly chose and sampled 40 transects from 51 of the original 60 pre-existing transects that had been unburned in the Warm Fire. I sampled all 9 of the original 60 pre-existing transects that were burned by the fire. Finally, during the summer of 2007, I established 11 new transects in forests burned at high (n=6) and low (n=5)

severity, bringing the total transects I sampled to 60. Previously established transects were used in my study to (1) minimize transect establishment time and to maximize data collection opportunities and (2) to provide for future comparisons of pre- and post-fire data to model prey abundance and goshawk reproduction over time. By the 1980s nearly the entirety of the North Kaibab Ranger District had received single-tree (locally known as “pick and pluck”) selection harvests. In the 1970s and 80s a more intensive stand management was introduced whereby 0.12-0.16 km<sup>2</sup> areas were harvested with shelter-wood and seed-tree prescriptions (Sesnie and Bailey. 2003). By 1990, approximated 30% of the North Kaibab Ranger District had been shelter-wood or seed-tree harvested (Sesnie and Bailey 2003). Thus, in the Salafsky (2004) study and in my study, transects were allocated among single-tree and shelter-wood/seed-tree harvested forests according to the approximate proportional areas of these past treatments. However, for the purposes of my study, I did not distinguish between the two harvest treatments and assumed that bird and mammal abundances reported here reflected average abundances in the two treatments. Ten of the transects I sampled in unburned ponderosa pine transects had received seed-tree harvest treatments before the Warm Fire, five in unburned mixed-conifer forests had also received seed-tree treatments, and six in high-severity mixed-conifer forest had received seed-tree treatments.

The 11 new transects that I established were located and classified using Warm Fire vegetation burn severity maps (North Kaibab Ranger District 2007). The eastern edge of the Warm Fire, characterized by narrow ridges and steep drainages, presented a challenge when selecting new transects due to the need for uniformly viewable areas on each side of transects. This requirement was identical to that used for establishing and sampling all of the pre-existing transects (Salafsky 2004). Prior to establishment, all potential new transect locations were visited to confirm burn severity as indicated by the burn severity maps (North Kaibab Ranger District 2007), forest type, management classification, and to address logistical and safety considerations. All pre-existing transects were also visited to confirm fire severity classification, forest type, and management classification. An equal number of transects were located in each forest type (n=30). Fire size (proportional to each severity category), habitat (unburned, burned) continuity, and transect viewable area constrained the number of transects in each strata. All

transects were located completely inside one forest type and fire severity category. Transect width varied by the detectability specific to each bird and mammal species and reflected the criteria used in the Salafsky (2004) study. Each transect consisted of 11 reference stakes marked at 50 meter intervals, four of which were randomly selected along each transect to serve as center points for vegetation plots.

### **Vegetation Sampling**

I determined the mean density of live trees per km<sup>2</sup> on each transect by averaging the tree densities on each of the four plot centers on each transect. I used the variable-radius plot technique with an angle gauge and a basal area factor (BAF) of 20 ft<sup>2</sup>/acre (4.59 m<sup>2</sup>/ha) to determine tree densities. Live tree densities were determined from the center of each plot by collecting the diameter at breast height (dbh) of each live tree counted 'in' with the angle gauge. I then calculated the number of trees per km<sup>2</sup> from measurements collected and averaged the four plot values to determine a mean density for each transect. This procedure was used to calculate the mean density of all live and dead trees >20cm dbh.

Tree canopy cover estimates were estimated at each plot center using a spherical densitometer. The unburned plot canopy cover estimates that I used were collected by Susan Salafsky (personal communication) prior to the Warm Fire. Cover estimates for all pre-existing plots burned by the Warm Fire and all newly established plots were collected post-fire. Mean canopy cover estimates for each transect were determined by averaging the cover estimates calculated for each plot.

Cone counting was conducted to index potential food supplies for birds and mammals (cone productivity). In late July and early August of each year, the closest tree  $\geq 30$  cm dbh in each quadrant (NE, SE, SW, and NW) from the plot center was selected for cone counting. A total of four trees were sampled on each plot. If the closest tree  $\geq 30$  cm dbh was non-cone producing the count for that tree was recorded as zero. Current year cones were counted by placing a 1m<sup>2</sup> frame on the ground at eight cardinal directions (N, NE, E, SE, S, SW, W, NW) under the crown of each Douglas fir, Engelmann spruce, and ponderosa pine tree. To assess cone production of subalpine and white fir trees, which do not accumulate cones on the ground, cones were counted in the canopies of each tree using binoculars. Mean cone count estimates for each transect were determined by averaging the counts obtained from each plot.

Understory ground cover measurements were collected from 2007-2010 from the four 20m radius plots on each transect. Ocular estimates of live ground cover (grasses, forbs, and shrubs; tree seedlings and sapling excluded) were collected annually on all plots and the dominant forbs and shrubs were identified each year. Numbers of sapling (conifer and aspen)  $\geq 1.5$ m in height and  $\leq 10$ cm dbh in each plot were tallied. Understory plot measurements were collected by the same trained teams of two technicians during the study in order to minimize observer bias.

### **Prey Sampling**

From 2007-2010, I conducted distance sampling to estimate the abundance of bird and mammal species commonly found in goshawk diets on the Kaibab Plateau (Reynolds et al. 1994, Salafsky et al. 2005). Bird species sampled included American robin (*Turdus migratorius*), Dusky grouse (*Dendragapus obscurus*), Clark's nutcracker (*Nucifraga columbiana*), downy woodpecker (*Picoides pubescens*), hairy woodpecker (*Picoides villosus*), mourning dove (*Zenaida macroura*), northern flicker (*Colaptes auratus*), red-naped sapsucker (*Sphyrapicus nuchalis*), Steller's jay (*Cyanocitta stelleri*), three-toed woodpecker (*Picoides dorsalis*), and Williamson's sapsucker (*Sphyrapicus thyroideus*). Mammal species sampled included black-tailed jackrabbit (*Lepus californicus*), chipmunk (*Tamias spp.*), cottontail rabbit (*Sylvilagus spp.*), golden-mantled ground squirrel (*Callospermophilus lateralis*), Kaibab squirrel (*Sciurus aberti kaibabensis*), red squirrel (*Tamiasciurus hudsonicus*), and rock squirrel (*Spermophilus variegatus*). I conducted counts on 0.5km line transects following assumptions and sampling protocols in Buckland et al. (2001) and procedures used by Salafsky (2004). I sampled each transect once in 2007 (late summer) then twice (early summer and late summer) annually from 2008-2010. Early summer counts occurred between 28 May and 24 June and late summer counts between 23 July and 14 August. All surveys were conducted by two observers (2007: J. Lambert and W. Nimitz, 2008 – 2010: J. Lambert and C. Lambert). Each observer was experienced at identifying bird and mammal species both visually and aurally and each was trained to the same counting methods. During each sampling period six transects were sampled daily (3 transects per observer). Sampling was not conducted during rain or heavy wind due to reduced animal activity and detectability. The order of transect surveys was

randomized each sample period to minimize observer and time-of-day bias. Prey sampling began 0.5 hours after sunrise and was completed within 3.5 hours. Individuals seen or heard were identified to species and the perpendicular distance from the transect line to the individual was determined using a laser rangefinder (accuracy  $\pm 1$  m). The few individuals that could not be visually or aurally located accurately were not tallied.

### **Data Analysis**

Vegetation measurements at the transect level were evaluated using the least square means procedure in GLIMMIX with  $\alpha \leq 0.05$  to identify significant differences in vegetation within each forest type across fire severity categories (PROC GLIMMIX, SAS Institute version 9.4, 2012). The Tukey-Kramer adjustment for multiple comparisons was used to assess the differences in the means of canopy cover, live tree density, live large tree density (>20cm dbh), snag density, large snag density (>20cm dbh), live ground vegetation, saplings per plot, and cone production by conifer species (Ott and Longnecker 2001). All data were modeled in GLIMMIX using a beta distribution for percentage estimates and a negative binomial distribution for count data (PROC GLIMMIX, SAS Institute version 9.4, 2012). Common data transformations were applied to the data but none improved analysis so data were imported into SAS in an untransformed format. Data presented in figures and tables are also untransformed.

I used the `gdistsamp` (generalized distance sampling) function in the UNMARKED (Fiske and Chandler 2011) package of the statistical software R (The R Foundation, Version 3.0.1, 2013) to estimate the abundance of all birds and mammals. The `gdistsamp` function fits the generalized distance sampling model of Royle et al. (2004), which allows abundance and detection parameters to be modeled as functions of covariates (Royle et al. 2004). The UNMARKED package utilizes replicate sampling to estimate the abundance ( $\lambda$ ) of the sampled population (the number of individuals that could potentially use a survey area) and probability that a given bird or mammal species is available ( $\phi$ ) to be counted. I used the negative binomial distribution to model abundance and all observation distances were right truncated to conform to Salafsky (2004) protocols. Distance sampling estimates a detection function that

compensates for differences in detection probability among species, habitats, and distances. Thus, density estimates were not confounded by differences in detectability due to the different forest types that were sampled (Buckland et al. 1993).

The UNMARKED package has several advantages over other estimation programs that potentially improve the accuracy of abundance estimates and predictions. UNMARKED relaxes the assumption that all individuals on a survey line are detected [ $g(0) = 1$ ], and allows for temporary immigration and emigration of mobile species in and out of survey sites (Chandler et al. 2011). By accounting for temporary immigration and emigration of mobile species to and from unknown locations, UNMARKED estimates the abundance ( $\lambda$ ) of individuals that could potentially use a survey plot, given a set of conditions (Chandler et al. 2011), however these estimates do not represent the number of individuals per unit of area. The second element of the model estimates the probability that an individual will be available to be counted ( $\phi$ ) on a plot given a set of temporal and spatial conditions (Chandler et al. 2011). Abundance ( $\lambda$ ) was modeled by fire severity and availability ( $\phi$ ) (both reported in Appendix A) was modeled by forest type and year. Density (D) of each species per km<sup>2</sup> (reported in Appendix B) can be obtained by multiplying the abundance estimate by the availability estimate ( $\lambda \phi$ ).

Buckland et al. (1993, 2001) recommends that approximately 60 observations are needed for reliable estimates of abundance. Thus, I estimated abundance only for species with >60 detections per year and tested models of the effects of fire severity differences on these species. To achieve a minimum of 60 observations for some under-counted species, counts were aggregated to estimate abundance at the guild (i.e., woodpeckers) level while counts of a few other species were aggregated across within-year sample periods. To evaluate the best models for predicting bird and mammal abundances, I used Akaike's Information Criterion (AIC) and Akaike weights (Burnham and Anderson 2002) generated with the R statistical software package UNMARKED. Models were developed a priori to evaluate how fire severity and vegetation characteristics that I hypothesized to be biologically important (e.g. tree and snag densities, canopy cover, food resources, ground cover) influenced bird and mammal abundances (Table 3 for a list of models). I used a  $\Delta$ AIC of 2 to identify competitive models (Burnham and Anderson 2002).

Year was included in each model as a surrogate for unmeasured variables (e.g., precipitation, temperature). Because AIC comparisons require data sets of equal sample size, I was unable to use the 2007 count data to test models because I only sampled transects once that season.

## RESULTS

### **Tree and Snag Density and Canopy Cover**

Relative to no-burn and low-severity fire, total live tree density, large (>20cm dbh) live tree density, canopy cover, and conifer cone production were, as expected, lower in both ponderosa pine and mixed-conifer forests burned at high severity (Figures 2-3, Appendix C). The total and large (>20cm dbh) snag densities in high severity burns were higher in both ponderosa pine and in mixed-conifer forests than in low severity and unburned forests; however, these means were significantly different only for total snags in mixed-conifer and large snags in ponderosa pine (Figure 4.1, 4.2, Appendix C). Although there were wider ranges of total live trees and canopy cover in unburned versus low-severity burn in both forest types, there were no significant differences in mean numbers of total and large live trees, and canopy cover from unburned forests. In ponderosa pine forests, densities of total and large snags were significantly higher in low severity versus unburned forests, but not in mixed-conifer (Figures 2-4, Appendix C).

Mixed-conifer.—Mixed-conifer forest transects burned by high-severity fire had an average of 709 fewer live trees/ha and 63% lower canopy cover compared to low-severity fire and an average of 960 fewer live trees/ha and 58% less canopy cover compared to unburned transects (Table 4). Transects burned by high-severity had an average of 470 fewer large live trees/ha compared to low-severity fire and an average of 429 fewer large live trees/ha compared to unburned mixed-conifer forests. Post-fire there were more snags on transects burned by high-severity than low-severity and unburned transects. There were significantly more large snags on mixed-conifer transects burned by high-severity fire than unburned transects, but there was no significant difference in large snag densities on high versus low-severity fire transects. On average, transects burned by high-severity fire had 411 more snags/ha than low-severity and 375 more snags/ha than unburned transects. Transects that were burned by high-severity fire also had an average of 117 more large snags/ha on unburned transects. No statistical differences in

total live tree, total snag densities, and canopy cover were found on mixed-conifer transects burned by low-severity fire versus unburned transect due to minimal low-severity ground fire effects in these forests.

**Ponderosa pine.**—Ponderosa pine forest transects burned by high-severity fire had an average of 495 fewer total live trees/ha and 50% less canopy cover compared to low-severity fire and an average of 343 fewer live trees/ha and 41% less canopy cover compared to unburned transects (Table 4). Transects burned by high-severity had an average of 197 fewer large live trees/ha compared to low-intensity fire and an average of 174 fewer large live trees/ha compared to unburned ponderosa pine forests. Post Warm fire I found an average of 804 more snags/ha on transects burned by high-severity fire than unburned transects. However contrary to expectations, there was no statistical difference in total snag density between transects burned by high- and low-severity fire. Large snag density on transects burned by high-severity fire averaged 191 more snags/ha compared to low-severity fire, and 212 more snags/ha compared to unburned transects. On average there were also 21 more snags/ha on low-severity fire transects compared to unburned transects. No other statistical differences in tree or snag densities and canopy cover were found when forests burned by low-severity fire were compared to unburned transects.

### **Ground Cover**

The percentage of live ground cover was greater on mixed-conifer transects burned by high-severity fire compared to low-severity fire and unburned transects. In contrast, there was no significant difference in the percentage of ground cover when ponderosa pine forests were compared across fire severities (Table 4, Figure 5.1, 5.2). The dominant flora (forbs and shrubs) found on transects were identified to species and reported by forest type and fire severity (Appendix D).

**Mixed-conifer.**—Mixed-conifer ground plots burned by high-severity fire averaged 17% more ground cover than low-severity fire and 18% more ground cover than unburned transects. No statistical difference in ground cover was found between transects burned by low-severity fire and unburned transect.

Ponderosa pine.—No statistical difference in the percent of ground cover was found when transects were compared across fire severity types, possibly due to low vegetative growth on pine transects in the first year (2007) of observation (Figure 5.2, Appendix C).

### **Sapling Counts**

There were more saplings in the 20m radius plots in mixed-conifer and ponderosa pine forest burned by high-severity fire than in low-severity. There were more saplings in mixed-conifer forests burned by high-severity fire than in unburned forests, but there were no significant difference between pine forests burned by high-severity fire versus unburned. There were also fewer saplings in mixed-conifer and ponderosa pine transects burned by low-severity fire compared to unburned forest transects (Table 4, Figure 6.1, 6.2).

Mixed-conifer.—Mixed-conifer burned by high-severity fire had an average of 47 more saplings per transect than low-severity fire and 43 more saplings per transect than unburned transects. Transects burned by low-severity fire had an average of 4 fewer saplings per transect than unburned transects.

Ponderosa pine.—Ponderosa pine burned by high-severity fire had an average of 12 more saplings per transect than low-severity fire transects. However no significant difference in the number of saplings was found when transects burned by high-severity fire were compared to unburned transects. Transects burned by low-severity fire had 10 fewer saplings per transect than unburned transects.

### **Cone Productivity**

Mixed-conifer and ponderosa pine transects burned by high-severity fire had significantly fewer cones than low-severity fire and unburned transects. No significant difference in the number of cones were found along transects burned by low-severity fire versus unburned transects (Table 4, Figure 7.1, 7.2).

Mixed-conifer.—Mixed-conifer transects burned by high-severity fire had an average of 122 fewer cones per transect than low-severity fire and 297 fewer cones per transect than unburned transects. There was no significant difference in the number of cones counted on mixed-conifer transects burned by low-severity fire compared to unburned transects.

Ponderosa pine.—Ponderosa pine transects burned by high-severity fire had an average of 267 fewer cones per transect than low-severity fire and 239 fewer cones per transect than unburned transects. There was no significant difference in the number of cones counted on ponderosa pine transects burned by low-severity fire compared to unburned transects.

### **Bird and Mammal Abundance**

A total of 17 bird and mammal species were observed at least once during the four study years (Table 5). Of the birds and mammals reported in the diets of goshawks on the NKRD (Reynolds et al. 1994, Salafsky et al. 2005), only black-tailed jackrabbits were not observed. However, cotton-tailed rabbits, dusky grouse, rock squirrels, red-naped sapsuckers, and black-tailed jackrabbits were observed four or fewer times during transect surveys. Of the seven species with  $\geq 60$  observations per year (Table 5, those species included in predictive models), the hairy woodpecker (low:  $p=.46$ , unburned:  $p<.01$ ) and northern flicker (low:  $p=.02$ , unburned:  $p=.05$ ) were the only species that were more abundant in burned (high- or low-severity) than unburned forests. In contrast, golden-mantled ground squirrels (low:  $p=.03$ , high:  $p<.01$ ) were less abundant in burned (high- and low-severity) than unburned forests. Similarly, red squirrels (low:  $p<.01$ , unburned:  $p<.01$ ) were less abundant in forests burned by high-severity fire than low-severity and unburned forests. American robin, chipmunk, and Steller's jay abundances were similar in burned and unburned forests (Appendix A).

### **Predictive Models**

Models of fire severity differences on vegetation composition and structure show that fire severity was an important factor for predicting the abundance of some birds and mammals, but not all species (Table 3). The top model predicting American robin abundances contained year, forest type, and live ground cover covariates and had a weight of evidence of 0.98. American robin abundance was highest in mixed-conifer forests sites with greater ground vegetation cover. The top model predicting chipmunk abundance also contained year, forest type, and live ground cover and had a weight of evidence of 0.56. A second competing model, ( $\Delta AIC \leq 2$ ) contained year and forest type covariates and carried a weight of evidence of 0.28. While the inclusion of ground cover contributed some weight to the top

model, the second model with fewer covariates is more parsimonious and therefore may be a better predictor of chipmunk abundance. In the top model, chipmunk species were found to be most abundant in ponderosa pine then mixed-conifer forests. The top model for Steller's jay's had a weight of evidence of 0.23. Three other competing models, including the null model, carried weight suggesting that we may not have had a sufficiently good model to predict jay abundance. Northern flicker abundance was predicted best by the model containing year and canopy cover covariates with a weight of evidence of 0.44. Two other competing models also carried weight (year, ground cover,  $W_t=0.27$ ; and year, forest type, ground cover,  $W_t=0.17$ ). Either of these models, especially the second which had fewer covariates, could be an equal or better predictor of abundance. In the top model, flickers were most abundant in forests with less canopy cover, while in the second and third models they were most abundant in forests with more ground vegetation cover.

Fire severity was an important factor for predicting the abundance of three species showing the most sensitivity to fire effects. Fire severity was included in the best model for golden-mantled ground squirrels, hairy woodpeckers, and red squirrels. The top models for estimating abundances of golden-mantled ground squirrel ( $W_t = 0.64$ ), hairy woodpecker ( $W_t = 0.51$ ), and red squirrel ( $W_t = 1.00$ ) contained year, fire severity, and forest type. Golden-mantled ground squirrels had a second competing model containing year, forest type, and ground vegetation cover with a weight of evidence of 0.30. Since these competing models contained the same number of covariates, either model may equally predict abundance. In the top model, golden-mantled ground squirrels were more abundant in unburned ponderosa pine forests. Red squirrels were equally abundant in mixed-conifer forests unburned or burned by low-severity fire. In contrast, hairy woodpeckers were more abundant in forests burned at high- and low-severity. However, a second competing model with year and fire severity (no forest type) had an almost equal weight of evidence of 0.49, suggesting that forest type contributed minimal weight to the top model. This second model had fewer covariates and may be an equally acceptable or better predictor of hairy woodpecker abundance. Fire severity was in the top model predicting the total abundance of individuals in the woodpecker guild. This model contained year and fire severity and had a weight of evidence of

0.72. A second competing model included year, fire severity, and forest type with a weight of evidence of

0.28. Woodpecker guild abundance was highest in forests burned by high-severity fire with some evidence in the second competing model that abundance in this guild was higher in ponderosa pine than mixed-conifer forests.

## DISCUSSION

Forests are dynamic ecosystems that undergo change through vegetation growth, senescence, and mortality, and natural (i.e., fire, insects, diseases, and wind) and anthropogenic disturbances (i.e., tree harvests). These changes have both short and long term impacts on habitats and the animals that depend upon them (Bartels and Thompson 1993). The 2006 Warm Fire started on 4 June and burned 235 km<sup>2</sup> of ponderosa pine and mixed-conifer forest on the North Kaibab Ranger District. During the initial 18 days, the Warm Fire was managed as a wildland-use fire during which 77 km<sup>2</sup> was burned mostly at low- to moderate-severity. However, on 25 June the weather changed and winds pushed the fire south burning approximately 158 km<sup>2</sup> mostly at high-severity until containment on 4 July. As expected, the high-severity burn portions of the Warm Fire had the greatest influence on habitat composition and structure, reducing canopy cover, number of live trees, and conifer cone production. However, these losses were countered by a significant flush of new ground cover, especially in mixed-conifer forests, and creation of an abundance of snags, which are important habitats to many birds and mammals—especially to species in the woodpecker guild. Most of the soil's O horizon (organic material), which contains food resources for birds and mammals, was also consumed in high-severity fire, exposing the mineral soil in areas (personal observation). Mixed-conifer forests burned at high-severity had a greater percentage of live ground cover post-fire than forests burned by either low-severity or unburned areas, a response consistent with the idea that fire can be a driver of understory diversity and structure (Laughlin et al. 2006, McMasters 2010). Higher levels of herbaceous ground vegetation in areas of high-severity fire results from stimulation of fine root production, decomposition, nitrogen mineralization, and other micro-environmental processes (Covington et al. 1997, Turner et al. 1997). Grass, forb, and shrub cover can also increase with a reduction in tree competition (reduced shading) frequently diversifying the plant community (Moore and Deiter 1992, Covington et al. 1997). In low-severity burned areas an insufficient amount of the O horizon may have been consumed by

the fire, limiting mineral nutrient returns to the soil, constraining a flush of vegetative growth observed in some high-severity burns (Laughlin and Abella 2007). Sapling density, especially aspen, was high in areas burned at high-severity fire in the Warm Fire; a finding similar to other studies of fire effects on aspen regeneration (Smith et al. 2011). Large clumps of small diameter pine saplings were present on 15 or more unburned ponderosa pine transects. In the absence of regular surface fire, establishment and growth of patches and clumps of dense saplings, both beneath overstory trees and in opening between tree groups restrict the detection and access to ground prey (ground squirrels, rabbits, hares, grouse, turkey) by winged predators such as the northern goshawk that hunt in the sub-canopy spaces of forests (Reynolds et al. 1992, 1996, Moore et al. 1999, Covington 2003, Koprowski et al. 2006). Loss of the grass, forb, and shrub vegetation (which support many different plants, insects, birds, and mammals) due to ingrowth and shading by dense regeneration, especially in openings between tree groups, can result in reductions in biodiversity, trophic interactions, and biological productivity and efficiency (Moore et al. 2006, Reynolds et al. 2006, Kalies et al. 2012, Reynolds et al. 2013).

Red squirrel and golden-mantled ground squirrel abundances were lower in areas burned by high-severity fire, likely because of the loss of cone producing large live trees, nesting and denning sites (woody debris, logs, stumps, live tree nests), food resources (e.g., conifer cone middens, herbaceous plants), and live tree canopy. Direct mortality from fire, loss of cover, increased exposure to predators, and increased distance from re-colonizing populations are other possible factors affecting their post-burn abundances in high-severity fire areas (Converse et al. 2006, Koprowski et al. 2006, Kalies et al. 2012). Depending on scale, due to positive post-fire influences on understory productivity, food resources, and habitats of high-severity fire can benefit other small mammals such as mice (*peromyscus* spp.) and voles (*Microtus* spp.) (Kalies et al. 2012), which were not included in my study.

Low-severity fire, that consumes ground fuels and reduces tree regeneration, can benefit many mammal and bird species by restoring the composition, structure, and fine-scale habitat patchiness, including snags, logs, and woody debris, that typified ponderosa pine and mixed-conifer forests before fire suppression (Carey and Harrington 2001, Covington and Moore 1994, Kennedy and Fontaine 2009,

Kalies et al. 2012, Larson and Churchill 2012, Reynolds et al. 2013). These and other studies have shown that tree squirrels, rabbits, turkeys, and grouse respond favorably to habitat patchiness and diversity (tree, shrub, and grass / forb cover) created by low-severity ground fires (Allen et al. 1982, Schroeder 1984, Reynolds et al. 1992, Kalies et al. 2012). Chipmunks in this study were not affected by fire at any intensity level; suggesting that they may be capable of utilizing a wide range of habitats. Because chipmunks and red squirrels abundances were unaffected by low-severity fire, and were similar to abundances in unburned forests, restoration of vegetation composition and pattern with regular low-severity ground fire could have both short and long-term benefits for bird and mammal communities in mixed-conifer and ponderosa pine forests on the Kaibab Plateau, while causing minimal impact on some species.

Post-fire increases in large diameter fire-killed snags and canopy cover reduction benefit most woodpecker species and other birds that occupy more open forests (Murphy and Lehnhausen 1998, Jackson et al. 2002, Bock and Block 2005, Saab et al. 2009). In this study, modeled woodpecker species responded positively to high-severity fire. The abundance of northern flickers, a cavity nester, was associated with decreased canopy cover; a response that was not unexpected because this woodpecker forages in openings associated with both burned and unburned open canopied forests (Kotliar et al 2007). Large snags provide suitable nesting habitats and an abundant food resources for woodpeckers and other cavity-nesting and bark-gleaning species (Saab and Vierling 2001, Saab et al. 2005), but at least for bark-gleaning species the flush of sub-bark insects can be short-lived. Indeed, Bock and Block (2005) reported that hairy woodpecker home ranges varied by season and fire severity. Wintering foraging ranges were smaller two years post-fire, when food supplies were abundant, than six years post-fire. Northern flickers followed a similar pattern. Over time, as food supplies diminishes, woodpeckers traveled longer distances to find food, eventually dispersing away from burned

areas. Thus, the beneficial effects of high-severity fire for species in the woodpecker guild can be short-lived, especially relative to the length time required for the forest to regrow and for food supplies and cavity-nesting opportunities to replenish.

The effects of low-severity fire on bird abundance were mixed. Of the four bird species I was able to model; hairy woodpecker and northern flickers were the only species to show a distinctly positive relationship to high- (and low-: hairy woodpeckers) severity fire. American robins and Steller's jay's abundances were similar in burned and unburned forests in this study.

Many of the birds and mammals that I surveyed are habitat generalists, a behavioral strategy allowing them to occupy a variety of habitats. For example, ground cover was an important predictor of robin abundance whereas fire severity had no effect. While other studies have shown that robins can respond positively to post-fire vegetative growth and species composition (Sackett and Haase 1998, Kotliar et al 2007, George and Zack 2008), robin abundance estimates in my study area were similar across burn severities, despite fire severity. While vegetative re-growth in newly opened areas following high-severity fire may contribute to the abundance of robins, grass, forb, and shrub openings found throughout ponderosa pine and mixed-conifer forests in my study also provide foraging opportunities in unburned areas as well. The cavity excavating northern flicker that forages in grassy open-canopied forests (Saab and Vierling 2001, Saab et al. 2005, Kotliar et al. 2007, Wiebe et al. 2008), was also uniformly abundant across the fire strata in my study. Canopy openings and increased ground vegetation (a covariate found in a competing model) were both important predictors of flicker abundance, suggesting co-linearity between these two covariates. The Steller's jay, a habitat and foraging generalists with a diverse omnivorous diet (Greene et al. 1998), was equally abundant across burn severities and forest types in my study. Other studies examining the effects of fire on

Steller's jays and American robins showed both positive and inconclusive responses to post-fire habitat changes due to fire severity as well (Bock and Block 2005, Kotliar et al. 2007, Dickson et al. 2009).

Of the seven birds and mammals I modeled, all but red squirrels and golden mantled ground squirrels demonstrated adaptable use of habitat conditions-- including conditions resulting from forest type and fire severity. The high number of species included in this study that were habitat generalists contributed to the difficulty of identifying effects of fire severity on the bird and mammal communities in ponderosa pine and mixed-conifer forests. By competing models that included habitat covariates specific to each species, I hoped to explore the relationship between fire, habitat change, and abundance of each species. What I did not consider in this study was the effects of past forest management on the predictive models. During the past century, nearly the entire North Kaibab Ranger District received single tree harvests. In the 1960s through 1990 more intensive management was introduced including small < 4ha clearcuts and 12-16 ha shelterwood/seed tree harvests (Sesnie and Bailey 2003). I attempted to establish a sample of transects in the three burn/no burn categories in order to equally sample the most ubiquitous past tree harvesting prescriptions, single-tree and seed-tree, on the North Kaibab Ranger District. Unfortunately, due to the specific location of the Warm Fire, several burn severity areas did not contain equal areas of these management prescriptions suitable for situating transects; achieving a balance of management prescription by forest type was not possible. Given the differences in tree densities in single-tree harvested areas versus seed-tree harvested areas, one might expect some corresponding differences in bird and mammal abundances in the two treatments. The extent to which treating the two prescriptions as one on my final abundance estimates is unknown without further examination.

## CONCLUSIONS

Fire is the most important natural disturbance in southwestern pine and mixed-conifer forests (Covington and Moore 1994, Fulé et al. 1997). Frequent, low-severity surface fire was part of the evolutionary history of ponderosa pine and lower elevation dry mixed-conifer forests (Savage and Swetnam 1990, Kalies et al. 2012, Larson and Churchill 2012, Reynolds et al. 2013) and high-severity crown fires seldom occurred and was confined to small patches (Pyne 1996). Frequent low severity surface fire in these forest types maintained an open grass-forb-shrub matrix and a spatial pattern of trees aggregated into clumps, groups, and patches, with scattered individual trees (White 1985, Cooper 1961, Larson and Churchill 2012, Reynolds, et al. 2013). Both the tree species composition (fire-tolerant and shade-intolerant trees) and spatial pattern resulted from a feed-back between fire and vegetation. Frequent surface fire resulted in a characteristic composition and structure that in turn supported a low severity, surface fire regime (Larson and Churchill 2012). This historical composition and structure manifested in a fine-scale (<4ha), highly interspersed mix of habitats (grass-forb-shrub vegetation), dense groups of trees with interlocking crowns, snags, logs, and woody debris; (Reynolds et al. 2013) that supported a variety of bird and mammal species (rabbits, ground squirrels, grouse, turkeys, and tree squirrels).

In my study, each of the seven species (four birds, three mammals) with sufficient numbers of detections to model responded to fire or lack of fire in a variety of ways. Low-severity fire was significantly detrimental to only golden-mantled ground squirrels while three of the seven species (chipmunks, robin, and Steller's jays) showed no difference in abundance across the three burn strata. Furthermore, two species (red squirrel and golden-mantled ground squirrel) showed negative responses to high-severity fire while two other species (hairy

woodpeckers, northern flickers) showed a positive response to high-severity fire. This summary of responses suggested that high-severity fire on the North Kaibab Ranger District could be more detrimental to the larger suite of important goshawk prey than low-severity burn. Similar findings have been summarized in a literature review of the overall bird community response to high-severity fire in ponderosa pine forests, where only two species responded positively to high-severity, four species responded positively to moderate-severity fire, and 9 bird species responded negatively to high-severity fire (Bock and Block 2005). Other studies of low-severity fire, including prescribed fire, reported either few changes in cavity-nesting bird abundance (Horton and Mannan 1988) or declines in species common in areas with heavy forest cover, where birds favored more open forest conditions (Marshall 1963).

While knowledge of the effects of low-severity fire on birds and mammals in ponderosa pine and mixed-conifer is increasing, more information is needed to provide robust management recommendations (Bock and Block 2005). Nonetheless, during the last century vegetative species composition and structure in southwestern forests has changed to fire intolerant and shade tolerant tree species where forest openings are overgrown (Cooper 1960, Covington and Moore 1994). Heavy accumulations of ground and ladder fuels, and increased frequencies and sizes of stand-replacement fires also characterize these forests (Cooper 1960, Covington and Moore 1994). Grass-forb-shrub vegetation productivity, biodiversity, and bird and mammal habitats have also declined due to tree shading and increased habitat homogeneity (Kalies et al. 2012, Reynolds et al. 2013). In response, there is a demand for recommendations to restore the historical species compositions, structures, and fine-scale spatial patterns, which facilitate the return of low-severity fire regimes and habitat heterogeneity that characterized the evolutionary environment of native plants and animals in these forests (Larson and Churchill 2012). While my

ability to estimate abundances was limited by species detectability, to a small sample of the suite of species in Kaibab goshawk diets, low-severity fire was found to benefit or have a neutral impact on many of species in this study. Restoration of a fine-scale mix of habitats (small tree groups interspersed within a grass-forb-shrub matrix, snags, and logs) are needed to support populations of the majority of bird and mammals important to many food webs.

Table 1. Transect classification by forest type and fire severity.

	Unburned		Sub-total	Burned				Sub-total	Grand Total n=60
Forest Type	Pine n=23	Mixed Conifer n=17	n=40	Pine n=7		Mixed Conifer n=13		n=20	
Fire Severity				High n=3	Low n=4	High n=9	Low n=4		

Table 2. Pre- and Post-fire transect classification.

No. Transects	60 pre-existing		11 established post-fire		Grand Total n=60
Burn Status	Unburned	Burned	Burned		
No. Transects	n=51	n=9	n=11		
Selected for sampling	n=40	n=9	n=11		
Fire Severity		High n=9	High n=3	Low n=8	

Table 3. Prey species abundance related to habitat and fire severity.

<b>American Robin</b>				
Model	K	AIC	$\Delta$ AIC	$W_t$
Year + Forest Type + Ground Cover	8	1901.71	0.00	0.98
Year + Ground Cover	7	1910.17	8.46	0.01
Year + Forest Type	7	1913.30	11.59	0.00
Year + Severity + Forest Type	9	1915.15	13.44	0.00
Year + Severity	8	1921.83	20.12	0.00
Year + Snag Density (>20 cm dbh)	7	1924.73	23.03	0.00
Year	6	1925.94	24.23	0.00
Year + Tree Density	7	1927.31	25.60	0.00
Year + Tree Density (>20 cm dbh)	7	1927.47	25.76	0.00
Null	4	1936.11	34.41	0.00
<b>Chipmunks</b>				
Model	K	AIC	$\Delta$ AIC	$W_t$
Year + Forest Type + Ground Cover	8	1774.36	0.00	0.56
Year + Forest Type	7	1775.71	1.34	0.28
Year + Severity + Forest Type	9	1779.00	4.63	0.05
Year + Tree Density (>20 cm dbh)	7	1779.18	4.81	0.05
Year + Snag Density (>20 cm dbh)	7	1780.23	5.87	0.03
Year + Canopy Cover	7	1781.53	7.16	0.02
Year + Ground Cover	7	1784.31	9.95	0.00
Year	6	1785.37	11.00	0.00
Year + Severity	8	1785.54	11.18	0.00
Year + Cones	7	1787.21	12.85	0.00
Null	4	1788.65	14.29	0.00

Model rankings were based upon Akaike's Information Criteria (AIC), AIC difference ( $\Delta$ AIC), and Akaike's model weight ( $W_t$ ). Only 2008 – 2010 data could be included in the predictive models and evaluated using AIC.

Table 3 (continue). Prey species abundance related to habitat and fire severity.

<b>Golden-mantled Ground Squirrel</b>				
Model	K	AIC	$\Delta$ AIC	$W_i$
Year + Severity + Forest Type	9	1449.19	0.00	0.64
Year + Forest Type + Ground Cover	8	1450.72	1.53	0.30
Year + Forest Type	7	1454.09	4.90	0.06
Year + Severity	8	1465.20	16.01	0.00
Year + Tree Density (>20 cm dbh)	7	1468.38	19.19	0.00
Year + Canopy Cover	7	1469.00	19.82	0.00
Year + Ground Cover	7	1473.23	24.05	0.00
Year	6	1476.62	27.43	0.00
Year + Cones	7	1478.40	29.21	0.00
Null	4	1485.88	36.69	0.00
<b>Hairy Woodpecker</b>				
Model	K	AIC	$\Delta$ AIC	$W_i$
Year + Severity + Forest Type	9	1683.01	0.00	0.51
Year + Severity	8	1683.12	0.11	0.49
Year + Snag Density (>20 cm dbh)	7	1704.34	21.33	0.00
Year + Tree Density	7	1711.38	28.37	0.00
Year + Tree Density (>20 cm dbh)	7	1713.96	30.95	0.00
Year + Canopy Cover	7	1714.21	31.20	0.00
Year	6	1715.41	32.40	0.00
Year + Forest Type	7	1716.92	33.91	0.00
Null	4	1739.45	56.44	0.00

Model rankings were based upon Akaike's Information Criteria (AIC), AIC difference ( $\Delta$ AIC), Akaike's model weight ( $W_i$ ). Only 2008 – 2010 data could be included in the predictive models and evaluated using AIC.

Table 3 (continue). Prey species abundance related to habitat and fire severity.

**Northern Flicker**

Model	K	AIC	$\Delta$ AIC	$W_t$
Year + Canopy Cover	7	3886.02	0.00	0.44
Year + Ground Cover	7	3886.96	0.94	0.27
Year + Forest Type + Ground Cover	8	3887.87	1.85	0.17
Year + Tree Density (>20 cm dbh)	7	3889.46	3.44	0.08
Year + Severity	8	3892.02	6.00	0.02
Year + Severity + Forest Type	9	3893.97	7.95	0.01
Year + Snag Density (>20 cm dbh)	7	3895.66	9.64	0.00
Year	6	3896.28	10.26	0.00
Year + Forest Type	7	3897.14	11.12	0.00
Null	4	3910.16	24.14	0.00

**Red Squirrel**

Model	K	AIC	$\Delta$ AIC	$W_t$
Year + Severity + Forest Type	9	1744.94	0.00	1.00
Year + Tree Density (>20 cm dbh)	7	1757.62	12.69	0.00
Year + Canopy Cover	7	1767.14	22.20	0.00
Year + Severity	8	1769.29	24.35	0.00
Year + Forest Type	7	1770.19	25.25	0.00
Year + Forest Type + Ground Cover	8	1772.08	27.14	0.00
Year + Cones	7	1778.40	33.46	0.00
Year	6	1780.56	35.62	0.00
Year + Snag Density (>20 cm dbh)	7	1782.07	37.13	0.00
Year + Ground Cover	7	1782.27	37.33	0.00
Null	4	1795.42	50.48	0.00

Model rankings were based upon Akaike's Information Criteria (AIC), AIC difference ( $\Delta$ AIC), Akaike's model weight ( $W_t$ ). Only 2008 – 2010 data could be included in the predictive models and evaluated using AIC.

Table 3 (continue). Prey species abundance related to habitat and fire severity.

**Steller's Jay**

Model	K	AIC	$\Delta$ AIC	W <sub>t</sub>
Year + Forest Type	7	1577.72	0.00	0.23
Null	4	1577.95	0.22	0.20
Year	6	1578.70	0.98	0.14
Year + Forest Type + Ground Cover	8	1579.60	1.88	0.09
Year + Tree Density (>20 cm dbh)	7	1579.80	2.07	0.08
Year + Cones	7	1580.51	2.79	0.06
Year + Ground Cover	7	1580.66	2.94	0.05
Year + Tree Density	7	1580.67	2.94	0.05
Year + Severity	8	1580.95	3.23	0.05
Year + Severity + Forest Type	9	1581.14	3.42	0.04

**Woodpecker Guild**

Model	K	AIC	$\Delta$ AIC	W <sub>t</sub>
Year + Severity	8	2028.77	0.00	0.72
Year + Severity + Forest Type	9	2030.66	1.89	0.28
Year + Snag Density (>20 cm dbh)	7	2048.31	19.54	0.00
Year + Tree Density	7	2057.74	28.97	0.00
Year + Canopy Cover	7	2059.48	30.71	0.00
Year + Tree Density (>20 cm dbh)	7	2059.92	31.15	0.00
Year	6	2060.03	31.26	0.00
Year + Forest Type	7	2061.07	32.30	0.00
Null	4	2083.04	54.27	0.00

Model rankings were based upon Akaike's Information Criteria (AIC), AIC difference ( $\Delta$ AIC), Akaike's model weight (W<sub>t</sub>). Only 2008 – 2010 data could be included in the predictive models and evaluated using AIC.

Table 4. Transect metrics by forest type and fire severity.

Mixed Conifer - High Severity (n=9)					Ponderosa Pine - High Severity (n=3)				
	Mean	SD	Min	Max		Mean	SD	Min	Max
Overall Tree Density	9.46	12.15	0	32.6	Overall Tree Density	1.24	2.14	0	3.71
Tree Density >20cm dbh	9.46	12.15	0	32.6	Tree Density >20cm dbh	1.24	2.14	0	3.71
Overall Snag Density	476.93	283.04	21.32	794.33	Overall Snag Density	810.66	451.48	501.43	1328.76
Snag Density >20cm dbh	154.82	119.97	21.32	360.25	Snag Density >20cm dbh	215.05	77.86	166.75	304.87
% Canopy Cover	4.78	3.6	0	10	% Canopy Cover	3.33	3.51	0	7
% Ground Vegetation	72.1	14.61	35.83	92.83	% Ground Vegetation	63.94	19.46	14	84.78
Sapling Regeneration	58.75	6.79	0	150	Sapling Regeneration	33.33	33.33	0	100
Cones Production	41.33	102.82	0	480	Cones Production	20.83	65.72	0	229
Mixed Conifer Low Severity (n=4)					Ponderosa Pine - Low Severity (n=4)				
	Mean	SD	Min	Max		Mean	SD	Min	Max
Overall Tree Density	718.91	243.47	479.48	1057.48	Overall Tree Density	496.4	340.99	266.02	997.83
Tree Density >20cm dbh	479.24	286.77	170.85	840.96	Tree Density >20cm dbh	198.06	64.28	141.93	279.55
Overall Snag Density	65.69	49.15	0	103.42	Overall Snag Density	135.69	211.62	0	451.37
Snag Density >20cm dbh	55.02	42.39	0	103.3	Snag Density >20cm dbh	24	25.71	0	52.44
% Canopy Cover	67.75	2.87	64	71	% Canopy Cover	53.75	13.07	38	68
% Ground Vegetation	54.77	17.69	11.3	78.11	% Ground Vegetation	53.61	11.39	33.77	78.75
Sapling Regeneration	11.31	2.95	2.5	30.75	Sapling Regeneration	21.41	6.11	0	95
Cones Production	163.75	231.84	14	939	Cones Production	287.38	297.1	18	1132
Mixed Conifer - Unburned (n=17)					Ponderosa Pine - Unburned (n=23)				
	Mean	SD	Min	Max		Mean	SD	Min	Max
Overall Tree Density	969.01	629.42	141.48	2186.67	Overall Tree Density	343.91	467.01	20.4	2009.17
Tree Density >20cm dbh	438.41	189.55	119.72	682.83	Tree Density >20cm dbh	175.22	136.48	20.4	522.63
Overall Snag Density	101.72	183.16	0	772.59	Overall Snag Density	6.81	18.73	0	90.57
Snag Density >20cm dbh	37.53	39.26	4.64	129.06	Snag Density >20cm dbh	2.88	4.2	0	16.35
% Canopy Cover	62.18	24.48	14	84	% Canopy Cover	44.48	24.8	3	88
% Ground Vegetation	54.46	15.01	23.49	81.75	% Ground Vegetation	56.23	21.84	12.75	95.75
Sapling Regeneration	15.67	1.43	0	45.45	Sapling Regeneration	31.67	3.32	0	106
Cone Production	337.87	428.13	0	2663	Cone Production	259.72	298.66	0	1100

Tree and Snag densities are displayed in trees per hectare. All other means are averages for 2007-2010. n is the number of transects in a forest type and fire severity

Table 5. Species Count Data.

<i>Species Common Name</i>	<b>Species Counts by Fire Severity</b>											
	High				Low				Unburned			
	2007	2008	2009	2010	2007	2008	2009	2010	2007	2008	2009	2010
American Robin	0	29	79	29	3	0	12	10	27	60	53	92
Black-tailed Jackrabbit	0	0	0	0	0	0	0	0	0	0	0	0
Blue Grouse	0	0	0	0	0	0	0	0	1	0	0	1
Chipmunk spp.	1	1	11	20	13	11	16	19	38	78	76	92
Clark's Nutcracker	1	23	7	3	4	0	2	2	21	17	7	3
Cottontail Rabbit	0	0	0	0	0	0	0	0	0	0	1	1
Downy Woodpecker	6	10	5	0	2	2	2	6	12	4	5	3
Golden-mantled Ground Sqr.	0	1	1	5	6	3	7	12	25	56	98	106
Hairy Woodpecker	37	51	39	21	19	22	33	13	41	38	39	20
Kaibab Squirrel	0	0	1	0	2	4	1	4	7	20	8	11
Mourning Dove	5	3	14	0	0	1	4	0	8	14	7	7
Northern Flicker	51	52	102	54	25	23	26	19	154	141	170	145
Red-naped Sapsucker	0	0	0	0	0	0	0	0	4	0	0	0
Red Squirrel	0	1	2	1	6	10	29	32	127	83	123	118
Rock Squirrel	0	0	0	0	0	0	0	0	0	0	0	2
Steller's Jay	5	8	18	2	8	6	10	9	46	45	50	45
Three-toed Woodpecker	0	4	2	1	0	0	1	2	5	0	4	4
Williamson's Sapsucker	0	0	0	0	0	1	2	0	6	12	7	7

Count data for 2007 only includes one late summer sample period while 2008-2010 include two sample periods.




Figure 1. Kaibab Plateau Study Area




Figure 2.1. Total tree density. Horizontal **bold** lines represent the category medians. Dashed lines represent the group means. Ranges are represented by the whiskers and outliers by ‘o’.




Figure 2.2. Tree density >20 cm dbh (Large). Horizontal **bold** lines represent the category medians. Dashed lines represent the group means. Ranges are represented by the whiskers and outliers by ‘o’.




Figure 3. Canopy cover. Horizontal **bold** lines represent the category medians while the dashed lines represent the group means. Category ranges are represented by the whiskers and outliers by 'o'.

**Total Snag Density by Forest Type and Fire Severity**




Figure 4.1. Total snag density. Horizontal **bold** lines represent the category medians. Dashed lines represent the group means. Category ranges are represented by the whiskers and outliers by 'o'.

**Snag Density >20cm dbh by Forest Type and Fire Severity**




Figure 4.2. Snag density >20 cm dbh (Large). Horizontal **bold** lines represent the category medians. Dashed lines represent the group means. Ranges are represented by the whiskers and outliers by 'o'.




Figure 5.1. Ocular estimates of mixed conifer ground cover. Transects are by year and fire severity.




Figure 5.2. Ocular estimates of ponderosa pine ground cover. Transects are by year and fire severity.




Figure 6.1. Mixed conifer sapling counts. Transects are by year and fire severity.




Figure 6.2. Ponderosa pine sapling counts. Transects are by year and fire severity.




Figure 7.1. Mixed conifer cone productivity. Transects are by year and fire severity.




Figure 7.2. Ponderosa pine cone productivity. Transects are by year and fire severity.

## REFERENCES

- Agee, J.K., 1993. Fire Ecology of Pacific Northwestern Forests. Washington, DC: Island Press.
- Allen, S.A., Lancia, R.A., Betsill, C.W., 1982. Habitat preferences of cottontail rabbits in an intensive farm and a traditional farm. Annual Conference Southeast Fish and Wildlife Agencies 36, 614-626.
- Bartels, M.A., Thompson, D.P., 1993. *Spermophilus lateralis*. Mammalian Species 440, 1-8.
- Binkley, D., Romme, B., Cheng, T., 2008. Historical forest structure on the Uncompahgre Plateau: Informing restoration prescriptions for mountainside stewardship. Colorado State University, Colorado Forest Restoration Institute, Fort Collins, Colorado, USA.
- Binkley, D., Moore, M., Romme, W., Brown, P., 2006. Was Aldo Leopold right about the Kaibab deer herd? Ecosystems 9, 227-241.
- Bock, C.E., Block, W.M., 2005. Response of birds to fire in the American Southwest. Pp 1093-1099 in C.J. Ralph and T.D. Rich (editors). Bird conservation implementation and integration in the Americas: proceedings of the third international Partners in Flight Conference. General Technical Report PSW-GTR-191. In: USDA Forest Service Pacific Southwest Research Station, Albany, California, USA.
- Bright-Smith, D.J., Mannan, R.W., 1994. Habitat use by breeding male Northern Goshawks in Northern Arizona. Studies in Avian Biology 16, 58-65.
- Brown, D.E., (editor). 1994. Biotic communities: southwestern United States and northwestern Mexico. University of Utah Press, Salt Lake City, Utah.
- Brown, P.M., Kaye, M.W., Huckaby, L., Baisan, C., 2001. Fire history along environmental gradients in the Sacramento Mountains, New Mexico: Influences of local patterns and regional processes. Ecoscience 8, 115-126.
- Buckland, S.T., Anderson, D.R., Burnham, K.P., Laake, J.L., 1993. Distance sampling: estimating abundance of biological populations. Chapman 7 Hall, London, U.K.
- Buckland, S.T., Anderson, D.R., Burnham, K.P., Laake, J.L., Borchers, D.L., Thomas, L., 2001. Introduction to Distance Sampling, Oxford: Oxford University Press.
- Burnham, K.P., Anderson, D.R., 2002. Model selection and inference: a practical information-theoretic approach, 2<sup>nd</sup> Ed. Springer-Verlag, New York, NY. U.S.A.
- Carey, A.B., Harrington, C.A., 2001. Small mammals in young forests: implications for management for sustainability. Forest Ecology and Management. 65(4), 1014-1027.
- Chandler, R.B., Royle, J.A., King, D.I., 2011. Inference about density and temporary emigration in unmarked populations. Ecology 92(7), 1429-1435.
- Churchill, D.J., Larson, A.J., Dahlgreen, M.C., Franklin, J.F., Hessburg, P.F., Lutz, J.A., 2013. Restoring forest resilience: from reference spatial patterns to silvicultural prescriptions and monitoring. Forest Ecology and Management 291, 442-457.

- Converse, S.J., Block, W.M., White, G.C., 2006. Small mammal population and habitat responses to forest thinning and prescribed fire. *Forest Ecology and Management* 228, 263-273.
- Cooper, C.F., 1960. Changes in vegetation, structure, and growth of southwest pine forests since white settlement. *Ecological Monographs* 30, 129-164.
- Cooper, C.F., 1961. Pattern in ponderosa pine forests. *Ecology* 42, 493-499
- Covington, W.W., 2003. The evolutionary and historical context. Pp 26-47. In P. Friederici (ed.). *Ecological Restoration of Southwestern Ponderosa Pine Forests*. Washington, DC: Island Press.
- Covington, W.W., Moore, M.M., 1994. Southwestern ponderosa forest structure: Changes since Euro-American settlement. *Journal of Forestry* 92(1), 39-47.
- Covington, W.W., Fule, P.Z., Moore, M.M., Hart, S.C., Kolb, T.E., Mast, N.J., Sackett, S.S., Wagner, M.R., 1997. Restoring ecological health in ponderosa pine forests of the southwest. *Journal of Forestry* 95, 23-29.
- Dickson, B.G., Noon, B.R., Flather, C.H., Jentsch, S., Block, W.M., 2009. Quantifying the multi-scale response of avifauna to prescribed fire experiments in the southwest United States. *Ecological Applications* 19(3), 608-621.
- Fiske, I., Chandler, R., 2011. UNMARKED: An R Package for Fitting Hierarchical Models of Wildlife Occurrence and Abundance. *Journal of Statistical Software* 43(10), 1-23.
- Fule, P.Z., Laughlin, D.C., 2007. Wildland fire effects on forest structure over an altitudinal gradient, Grand Canyon National Park, USA. *Journal of Applied Ecology* 44, 136-146.
- Fule, P.Z., Covington, W.W., Moore, M.M., 1997. Determining reference conditions for ecosystem management of southwestern ponderosa pine forests. *Ecological Applications* 7, 895-908.
- Fule, P.Z., Korb, J.E., Wu, R., 2009. Changes in forest structure of a mixed-conifer forest, southwestern Colorado, USA. *Forest Ecology and Management* 258(7), 1200-1210.
- Fule, P.Z., Heinlein, T.A., Covington, W.W., Moore, M.M., 2000. Continuing fire regimes in remote forests of Grand Canyon National Park. In 'Proceedings: Wilderness science in a time of change' (Eds. DN Cole and SF McCool) Pp 242-248. Report RMRS-P-15. In: USDA Forest Service Rocky Mountain Research Station, Ogden, Utah, USA.
- Fule, P.Z., Crouse, J.E., Heinlein, T.A., Moore, M.M., Covington, W.W., VanKamp, G., 2003. Mixed-severity fire regime in high-elevation forest of the Grand Canyon, Arizona, USA. *Landscape Ecology* 18, 465-486.
- George, T.L, Zack, S., 2008. Bird occupancy and richness in ponderosa forests with contrasting forest structure and fire history. *Canadian journal of Forest Research* 38(5), 936-942.
- Greene, E., Davison, W., Muehter, V.R., 1998. Steller's jay (*Cyanocitta stelleri*). In *The Birds of North America*, No. 343 (A. Poole, Ed). The Birds of North America Online, Ithaca, New York.
- Griffis, K.L., Crawford, J.A., Wagner, M.R., Moir, W.H., 2001. Understory response to management treatments in northern Arizona ponderosa pine forests. *Forest Ecology and Management* 146, 239-245.

- Hoffmeister, D.F., 1986. Mammals of Arizona. The University of Arizona Press. Arizona Fish and Game Dept. Tucson, Arizona. Pp 206-213.
- Heyerdahl, E.K., Brubaker, L.B., Agee, J.K., 2001. Spatial controls of historical fire regimes: A multi-scale example from the interior west, USA. *Ecology* 82, 660-678.
- Horton, S.P., Mannan, R.W., 1988. Effects of prescribed fire on snags and cavity-nesting birds in southeastern Arizona pine forests. *Wildlife Society Bulletin* 16, 37-44.
- Jackson, J.A., Ouellet, H.R., Jackson, B.J., 2002. Hairy Woodpecker (*Picoides villosus*). In *The Birds of North America*, No. 702 (A. Poole, Ed). The Birds of North America Online, Ithaca, New York.
- Joy, S.M., 2002. Northern Goshawk habitat on the Kaibab National Forest in Arizona: factors affecting nest locations and territory quality. Ph.D. dissertation, Colorado State University, Ft. Collins, CO, USA.
- Kalies, E.L., Dickson, B.G., Chambers, C.L., Covington, W.W., 2012. Community occupancy responses of small mammals to restoration treatments in ponderosa pine forests, northern Arizona, USA. *Ecological Applications* 22(1), 204-217.
- Kennedy, P.L., Fontaine, J.B., 2009. Synthesis of knowledge of the effects of fire and fire surrogates on wildlife in U.S. dry forests. Special Report 1096. Oregon State University Extension Service, Corvallis, Oregon, USA.
- Koprowski, J.L., Leonard, K.M., Zugmeyer, C.A., Jolley, J.L., 2006. Direct effects of fire on endangered Mount Graham Red Squirrels. *The Southwestern Naturalist* 51, 59-63.
- Korb, J.E., Fule, P.Z., Wu, R., 2013. Variability of warm/dry mixed conifer forests in southwestern Colorado, USA: Implications for ecological restoration. *Forest Ecology and Management* 304, 182-191.
- Kotliar, N.B., Kennedy, P.L., Ferree, K., 2007. Avifaunal responses to fire in southwestern montane forests along a burn severity gradient. *Ecological Applications* 17(2), 491-507.
- Larson, A.J., Churchill, D., 2012. Tree Spatial Patterns in fire-frequent forests of western North America, including mechanisms of pattern formation and implications for designing fuel reduction and restoration treatments. *Forest Ecology and Management* 267, 74-92.
- Laughlin, D.C., Abella, S.R., 2007. Abiotic and biotic factors explain independent gradients of community composition in ponderosa pine forests. *Ecological Modeling* 205, 231-240.
- Laughlin, D.C., Moore, M.M., Bakker, J.D., Casey, C.A., Springer, J.D., Fule, P.Z., Covington, W.W., 2006. Assessing targets for the restoration of herbaceous vegetation in ponderosa pine forests. *Restoration Ecology* 14, 548-560.
- Marshall, J.T., 1963. Fire and birds in the mountains of southern Arizona. *Proceedings, Annual Tall Timbers Fire Ecology Conference* 2, 135-141.
- McMaster, M.A., 2010 Effects of Fire and Post-Fire Seeding on Plant Communities in a Ponderosa Pine Forest. M.S. Thesis. Northern Arizona University, Flagstaff, AZ.
- Moore, M.M., Deiter, D.A., 1992. Stand density index as a predictor of forage production in northern Arizona pine forests. *Journal of Range Management* 45, 267-271.

- Moore, M.M., Covington, W.W., Fule, P.Z., 1999. Evolutionary environment, reference conditions, and ecological restoration: A southwestern ponderosa pine perspective. *Ecological Applications* 9, 1266-1277.
- Moore, M.M., Huffman, D.W., Fule, P.Z., Covington, W.W., Crouse, J.E., 2004. Comparison of historical and contemporary forest structure and composition on permanent plots in southwestern ponderosa pine forests. *Forest Science* 50, 162-176.
- Murphy, E.C. and Lehnhausen, W.A., 1998. Density and foraging ecology of woodpeckers following a stand-replacement fire. *Journal of Wildlife Management* 62, 1359-1372.
- North Kaibab Ranger District, 2007. Warm Fire Vegetation Burn Severity Maps. USDA Forest Service, North Kaibab Ranger District, Kaibab National Forest, Fredonia, AZ.
- Ott, R.L., Longnecker, M., 2001. An introduction to statistical methods and data analysis, 5<sup>th</sup> edition. Duxbury, Pacific Grove, CA.
- Pyne, S.J., Andrews, P.L., Laven, Richard, D., 1996. Introduction to Wildland Fire. Wiley-Interscience publication.
- R Development Core Team, 2008. R: A language and environment for Statistical computing. R Foundation for Statistical Computing ver 3.0.1, Vienna, Austria. ISBN 3-900051-07-0, URL <http://www.R-project.org>.
- [Reynolds, R. T.](#), 1983. Management of western coniferous forests habitat for nesting accipiter hawks. General Technical Report. RM-102. USDA Forest Service. Rocky Mountain Forest and Range Experiment Station, Fort Collins, CO.
- Reynolds, R.T., Meslow, E.C., 1984. Partitioning of food and niche characteristics of coexisting Accipiter in Oregon. *Journal of Wildlife Management* 46, 124-138.
- Reynolds, R.T., Wiens, J.D., Salafsky, S.R., 2006. A review and evaluation of factors limiting Northern Goshawk populations. *Studies in Avian Biology* 31, 260-273.
- Reynolds, R.T., Joy, S.M., Leslie, D.G., 1994. Nest productivity, fidelity, and spacing of Northern Goshawks in Arizona. *Studies in Avian Biology* 16, 106-113.
- Reynolds, R.T., Wiens, J.D., Joy, S.M., Salafsky, S.R., 2005. Sampling considerations for demographic and habitat studies of Northern Goshawks. *Journal of Raptor Research* 39, 274-285.
- Reynolds, R.T., Sanchez Meador, A.J., Youtz, J.A., Nicolet, T., Matonis, M.S., Jackson, P.L., DeLorenzo, D.G., Graves, A.D., 2013. Restoring Composition and Structure in Southwestern Frequent-Fire Forests: A science-based framework for improving ecosystem resiliency. General Technical Report RMRS-GTR-310. In: USDA Forest Service Rocky Mountain Research Station, Fort Collins, Colorado, USA.
- Reynolds, R.T., Graham, R.T., Reiser, M.H., Bassett, R.L., Kennedy, P.L., Boyce Jr., D.A., Goodwin, G., Smith, R., Fisher, E.L., 1992. Management Recommendations for the Northern Goshawk in the Southwestern United States. General Technical Report RMRS-GTR-217. In: USDA Forest Service Rocky Mountain Research Station, Fort Collins, Colorado, USA.

- Romme, W.H., Floyd, M.L., Hanna, D., 2009. Historical range of variability and current landscape conditions analysis: South Central Highlands Section, Southwestern Colorado & Northwestern New Mexico. Colorado State University, Colorado Forest Restoration Institute, Fort Collins, CO, and USDA Forest Service, Rocky Mountain Region, Golden, CO.
- Royle, J.A., Dawson, D.K., Bates, S., 2004. Modeling Abundance Effects in Distance Sampling. *Ecology* 85(6), 1591-1597.
- Sackett, S.S., Haase, S.M., 1998. Two case histories for using prescribed fire to restore ponderosa pine ecosystems in Northern Arizona. Pp 380-389 in T.L. Pruden and L.A. Brennan (eds.). *Fire in Ecosystem Management: Shifting the Paradigm from Suppression to Prescription*. Proceedings of the 20<sup>th</sup> Tall Timbers Fire Ecology Conference, 7-10 May 1996, Boise, ID. Tall Timbers Research Station. Tallahassee, Florida, USA.
- Sallabanks, R., James, F.C., 1999. American Robin (*Turdus migratorius*). In *The Birds of North America*, No. 462 (A. Poole, Ed). The Birds of North America Online, Ithaca, New York.
- SAS Institute, 2012. SAS/PROC GLIMMIX, Version 9.4. SAS Institute, Inc., Cary, NC, USA.
- Savage, M., Swetnam, T.W., 1990. Early 19<sup>th</sup> Century fire decline following sheep pasturing in a Navajo ponderosa pine forest. *Ecological Society of America* 71(6), 2374-2378.
- Schroeder, R.L., 1984. Habitat suitability index models: blue grouse. Washington: U.S. Fish and Wildlife Service, U.S. Department of the Interior. FWS/OBS-82/10.81.
- Sesnie, S., Bailey, J., 2003. Using history to plan the future of old-growth ponderosa pine. *Journal of Forestry* 101, 40-47.
- Smith, E.A., O'Loughlin, D., Buck, J.R., St.Clair, S.B., 2011. The influences of conifer succession, physiographic conditions and herbivory on quaking aspen regeneration after fire. *Forest Ecology and Management* 262, 325-330.
- Squires, J.R., Reynolds, R.T., 1997. Northern Goshawk (*Accipiter gentilis*). In A. Poole and F.Gill (EDS.). *The birds of North America*, No. 298. The Birds of North America Inc., Philadelphia, PA, USA.
- Saab, V., Vierling, K.T., 2001. Reproductive success of Lewis's Woodpecker in burned and cottonwood riparian forests. *Condor* 103, 491-501.
- Saab, V.A., Russell, R.E., Dudley, J.G., 2009. Nest-site selection by cavity-nesting birds in relation to post-fire salvage logging. *Forest Ecology and Management* 257(1), 151-159.
- [Saab, V. A.](#), Powell, H.D.W., Kotliar, N.B., Newlon, K.R., 2005. Variation in fire regimes of the Rocky Mountains: Implications for avian communities and fire management. *Studies in Avian Biology* 30, 76-96.
- [Saab, V. A.](#), Brannon, R., Dudley, J.G., Donohoo, L., Vanderzanden, D., Johnson, V., Lachowski, H., 2002. Selection of fire-created snags at two spatial scales by cavity-nesting birds. In *Proceedings of The Symposium on The Ecology and Management of Dead Wood in Western Forests*, Nov 2-4, 1999, Reno, Nevada, editors Shea, P. J.; W.F. Laudenslayer, Jr.; B. Valentine; C. P. Weatherspoon, and T. E. Lisle. General Technical Report PSW-GTR-181. In: USDA Forest Service Pacific Southwest Research Station, USA.

[Salafsky, S. R.](#), 2004. Covariation between prey abundance and northern goshawk fecundity on the Kaibab Plateau, Arizona. M.S. Thesis, Colorado State University, Fort Collins, Colorado, USA.

[Salafsky, S. R.](#), [R. T. Reynolds](#), Noon, B.R., 2005. Patterns of temporal variation in goshawk reproduction and prey resources. *Journal of Raptor Research* 39(3), 237–246.

[Salafsky, S. M.](#), [R. T. Reynolds](#), Noon, B.R., Wiens, J.A., 2007. Reproductive responses of northern goshawks to variable prey populations. *Journal of Wildlife Management* 71, 2274-2283.

[Squires, J. R.](#), Reynolds, R.T., 1997. Northern goshawk. Squires, J.R., and R.T. Reynolds. 1997. Northern goshawk (*Accipiter gentilis*). In *The Birds of North America*, No. 298. A. Poole and F. Gill (editors). The Academy of Natural Sciences Philadelphia, PA, and the American Ornithologists' Union. Washington, D.C. USA.

Swetnam, T.W., Baisan, C.H., 1996. Historical fire regime patterns in the southwestern United States since AD 1700. Pp. 11-32 in C.D. Allen (ed.). 2<sup>nd</sup> La Mesa Fire Symposium; Los Alamos, NM. General Technical Report RM-GTR-286. USDA Forest Service Rocky Mountain Research Station, Fort Collins, Colorado, USA.

Turner, M.G., Romme, W.H., Gardner, R.H., Hargrove, W.W., 1997. Effects of fire size and pattern on early succession in Yellowstone National Park. *Ecological Monographs* 67(4), 411-433.

White, A.S., 1985. Pre-settlement regeneration patterns in a Southwestern ponderosa pine stand. *Ecology* 66, 589-594.

White, M.A., VanKat, J.L., 1993. Middle and high elevation coniferous forest communities of the North Rim region of Grand Canyon National Park, Arizona, USA. *Vegetatio* 109, 161-174.

Wiebe, K.L., Moore, W.S., 2008. Northern Flicker (*Colaptes auratus*). In *The Birds of North America*, No. 166a (A. Poole, Ed). *The Birds of North America Online*, Ithaca, New York. USA.

## APPENDICES

APPENDIX A

						Mixed Conifer					Ponderosa Pine			
	Severity	Abundance	SE	LCL	UCL	Year	Availability	SE	LCL	UCL	Availability	SE	LCL	UCL
<b>American Robin</b>	High	5.46	1.56	3.12	9.54	2007	0.04	0.01	0.03	0.07	0.02	0.01	0.01	0.03
	Low	2.81	1.16	1.25	6.29	2008	0.06	0.01	0.04	0.09	0.03	0.01	0.02	0.04
	Unburned	2009	0.10	0.02	0.07	0.14	0.04	0.01	0.03	0.07	0.04	0.01	0.03	0.07
		2010	0.09	0.02	0.07	0.13	0.04	0.01	0.02	0.06	0.04	0.01	0.02	0.06
<b>Chipmunks</b>	High	9.18	3.40	4.44	18.98	2007	0.02	0.01	0.01	0.04	0.07	0.02	0.04	0.11
	Low	9.25	3.27	4.62	18.51	2008	0.02	0.01	0.01	0.03	0.06	0.01	0.04	0.09
	Unburned	2009	0.02	0.01	0.01	0.04	0.07	0.02	0.05	0.11	0.07	0.02	0.05	0.11
		2010	0.03	0.01	0.02	0.05	0.09	0.02	0.06	0.13	0.09	0.02	0.06	0.13
<b>Clark's Nutcracker</b>	High	3.99	2.09	1.43	11.15	2007	0.03	0.02	0.01	0.10	0.03	0.02	0.01	0.10
	Low	1.60	1.11	0.41	6.27	2008	0.03	0.01	0.01	0.07	0.02	0.01	0.01	0.08
	Unburned	2009	0.01	0.01	0.00	0.03	0.01	0.01	0.00	0.03	0.01	0.01	0.00	0.03
		2010	0.01	0.00	0.00	0.02	0.00	0.00	0.00	0.02	0.00	0.00	0.00	0.02
<b>Downy Woodpecker</b>	High	4.83	2.76	1.57	14.83	2007	0.03	0.02	0.01	0.11	0.03	0.02	0.01	0.11
	Low	4.44	2.83	1.28	15.46	2008	0.01	0.01	0.00	0.05	0.01	0.01	0.00	0.04
	Unburned	2009	0.01	0.01	0.00	0.04	0.01	0.01	0.00	0.04	0.01	0.01	0.00	0.03
		2010	0.01	0.01	0.00	0.03	0.01	0.01	0.00	0.03	0.01	0.00	0.00	0.03
<b>Golden-mantled Ground Squirrel</b>	High	1.19	0.82	0.31	4.56	2007	0.01	0.00	0.00	0.02	0.04	0.01	0.02	0.07
	Low	2.91	1.43	1.12	7.60	2008	0.01	0.00	0.00	0.02	0.04	0.01	0.02	0.07
	Unburned	2009	0.01	0.00	0.01	0.03	0.06	0.02	0.03	0.10	0.06	0.02	0.03	0.10
		2010	0.01	0.00	0.01	0.03	0.07	0.02	0.04	0.11	0.07	0.02	0.04	0.11
<b>Hairy Woodpecker</b>	High	9.52	2.30	5.93	15.29	2007	0.09	0.02	0.05	0.14	0.09	0.03	0.05	0.16
	Low	7.99	2.47	4.35	14.65	2008	0.05	0.01	0.03	0.08	0.06	0.02	0.03	0.10
	Unburned	2009	0.05	0.01	0.03	0.08	0.06	0.02	0.03	0.10	0.06	0.02	0.03	0.10
		2010	0.02	0.01	0.01	0.04	0.03	0.01	0.01	0.04	0.03	0.01	0.01	0.05
<b>Kaibab Squirrel</b>	High	0.07	0.11	0.00	1.67	2007	0.03	0.05	0.00	0.36	0.16	0.20	0.01	0.78
	Low	0.43	0.60	0.03	6.52	2008	0.05	0.07	0.00	0.50	0.23	0.28	0.01	0.87
	Unburned	2009	0.02	0.02	0.00	0.21	0.09	0.12	0.01	0.63	0.09	0.12	0.01	0.63
		2010	0.03	0.04	0.00	0.34	0.15	0.19	0.01	0.77	0.15	0.19	0.01	0.77

APPENDIX A

						Mixed Conifer					Ponderosa Pine			
	Severity	Abundance	SE	LCL	UCL	Year	Availability	SE	LCL	UCL	Availability	SE	LCL	UCL
<b>Mourning Dove</b>	High	3.59	2.14	1.11	11.56	2007	0.02	0.01	0.01	0.07	0.05	0.04	0.01	0.21
	Low	0.75	0.62	0.15	3.81	2008	0.01	0.01	0.00	0.05	0.04	0.03	0.01	0.15
	Unburned	0.96	0.67	0.25	3.80	2009	0.02	0.01	0.01	0.07	0.05	0.04	0.01	0.20
		2010	0.01	0.00	0.00	0.02	0.01	0.01	0.00	0.07	0.01	0.01	0.00	0.07
<b>Northern Flicker</b>	High	6.86	1.30	4.73	9.96	2007	0.13	0.02	0.09	0.18	0.13	0.03	0.09	0.19
	Low	3.76	0.94	2.30	6.15	2008	0.06	0.01	0.04	0.08	0.06	0.01	0.04	0.09
	Unburned	4.81	0.86	3.39	6.83	2009	0.08	0.01	0.06	0.12	0.08	0.02	0.06	0.12
		2010	0.06	0.01	0.04	0.09	0.06	0.01	0.04	0.09	0.06	0.01	0.04	0.09
<b>Red Squirrel</b>	High	0.11	0.07	0.04	0.35	2007	0.28	0.07	0.16	0.42	0.05	0.01	0.03	0.08
	Low	6.30	2.16	3.22	12.33	2008	0.10	0.03	0.06	0.17	0.02	0.00	0.01	0.03
	Unburned	5.02	1.15	3.20	7.86	2009	0.17	0.04	0.10	0.27	0.03	0.01	0.02	0.04
		2010	0.17	0.04	0.10	0.27	0.03	0.01	0.02	0.04	0.03	0.01	0.02	0.04
<b>Steller's Jay</b>	High	3.60	1.36	1.72	7.53	2007	0.03	0.01	0.01	0.05	0.04	0.02	0.02	0.09
	Low	3.84	1.52	1.77	8.36	2008	0.01	0.00	0.01	0.03	0.02	0.01	0.01	0.04
	Unburned	4.27	1.32	2.33	7.84	2009	0.02	0.01	0.01	0.03	0.03	0.01	0.01	0.05
		2010	0.01	0.00	0.01	0.02	0.02	0.01	0.01	0.02	0.02	0.01	0.01	0.04
<b>Three-toed Woodpecker</b>	High	1.61	1.64	0.22	11.80	2007	0.03	0.03	0.00	0.19	0.01	0.01	0.00	0.08
	Low	2.33	2.37	0.32	17.06	2008	0.01	0.01	0.00	0.09	0.00	0.00	0.00	0.03
	Unburned	1.40	1.25	0.24	8.10	2009	0.02	0.02	0.00	0.14	0.01	0.01	0.00	0.05
		2010	0.02	0.02	0.00	0.14	0.01	0.01	0.00	0.14	0.01	0.01	0.00	0.05
<b>Williamson's Sapsucker</b>	High	0.00	0.00	0.00	0.00	2007	0.03	0.04	0.00	0.29	0.00	0.01	0.00	0.06
	Low	3.86	5.06	0.30	50.25	2008	0.03	0.04	0.00	0.30	0.01	0.01	0.00	0.06
	Unburned	5.55	6.80	0.50	61.28	2009	0.02	0.03	0.00	0.23	0.00	0.00	0.00	0.05
		2010	0.02	0.02	0.00	0.19	0.00	0.00	0.00	0.19	0.00	0.00	0.00	0.04

Appendix A. Abundance estimates by fire severity, and probability of availability by year and forest type, for all species. Birds and mammals with  $\geq 60$  obs. per year in **bold** were used to test predictive models. SE: Standard Error, LCL / UCL: Lower and Upper 95% Confidence Limits.

## APPENDIX B

2007			Mixed Conifer			Ponderosa Pine		
Species common name	Severity	n	D	LCL	UCL	D	LCL	UCL
American Robin	High	0	0.12	0.04	0.32	0.05	0.01	0.15
	Low	3	0.06	0.02	0.21	0.02	0.01	0.10
	Unburned	27	0.09	0.04	0.22	0.04	0.01	0.10
Chipmunk spp.	High	1	0.09	0.02	0.35	0.31	0.09	1.00
	Low	13	0.09	0.03	0.34	0.31	0.10	0.98
	Unburned	38	0.11	0.04	0.30	0.36	0.15	0.86
Clark's Nutcracker	High	1	0.14	0.02	1.08	0.13	0.01	1.13
	Low	4	0.05	0.00	0.61	0.05	0.00	0.63
	Unburned	21	0.06	0.01	0.52	0.06	0.01	0.54
Downy Woodpecker	High	6	0.17	0.02	1.61	0.15	0.01	1.57
	Low	2	0.16	0.01	1.68	0.14	0.01	1.63
	Unburned	12	0.06	0.01	0.60	0.05	0.00	0.58
Golden-mantled Ground Squirrel	High	0	0.00	0.00	0.04	0.02	0.00	0.16
	Low	6	0.01	0.00	0.06	0.06	0.01	0.27
	Unburned	25	0.03	0.01	0.12	0.16	0.05	0.50
Hairy Woodpecker	High	37	0.42	0.16	1.07	0.45	0.16	1.26
	Low	19	0.35	0.12	1.02	0.38	0.11	1.20
	Unburned	41	0.11	0.04	0.30	0.12	0.04	0.35
Kaibab Squirrel	High	0	0.00	0.00	0.60	0.01	0.00	1.31
	Low	2	0.01	0.00	2.35	0.07	0.00	5.12
	Unburned	7	0.01	0.00	1.13	0.04	0.00	2.45
Mourning Dove	High	5	0.07	0.01	0.80	0.18	0.01	2.38
	Low	0	0.01	0.00	0.26	0.04	0.00	0.79
	Unburned	16	0.02	0.00	0.26	0.05	0.00	0.78
Northern Flicker	High	51	0.44	0.22	0.88	0.44	0.20	0.93
	Low	25	0.24	0.10	0.54	0.24	0.10	0.58
	Unburned	154	0.31	0.15	0.60	0.31	0.15	0.64
Red Squirrel	High	0	0.02	0.00	0.07	0.00	0.00	0.01
	Low	6	0.87	0.26	2.61	0.16	0.05	0.51
	Unburned	127	0.69	0.26	1.66	0.13	0.05	0.32
Steller's Jay	High	5	0.05	0.01	0.20	0.08	0.02	0.32
	Low	8	0.05	0.01	0.22	0.09	0.02	0.36
	Unburned	46	0.06	0.01	0.20	0.10	0.03	0.34
Three-toed Woodpecker	High	0	0.05	0.00	2.29	0.02	0.00	0.91
	Low	0	0.07	0.00	3.31	0.03	0.00	1.31
	Unburned	5	0.04	0.00	1.57	0.02	0.00	0.62
Williamson's Sapsucker	High	0	0.00	0.00	0.00	0.00	0.00	0.00
	Low	0	0.11	0.00	14.64	0.02	0.00	3.13
	Unburned	6	0.16	0.00	17.85	0.03	0.00	3.81

Appendix B. 2007 Density (D) estimates for birds and mammals per hectare with one sample period.

APPENDIX B

2008			Mixed Conifer			Ponderosa Pine		
Species common name	Severity	n	D	LCL	UCL	D	LCL	UCL
<b>American Robin</b>	High	29	0.35	0.14	0.87	0.14	0.05	0.43
	Low	0	0.18	0.06	0.58	0.07	0.02	0.28
	Unburned	60	0.27	0.12	0.58	0.11	0.04	0.29
<b>Chipmunk spp.</b>	High	1	0.17	0.05	0.63	0.57	0.18	1.79
	Low	11	0.17	0.05	0.62	0.58	0.19	1.75
	Unburned	78	0.21	0.08	0.54	0.68	0.29	1.55
Clark's Nutcracker	High	23	0.10	0.01	0.80	0.09	0.01	0.84
	Low	0	0.05	0.00	0.44	0.03	0.02	0.13
	Unburned	11	0.05	0.01	0.39	0.04	0.00	0.40
Downy Woodpecker	High	10	0.07	0.01	0.68	0.06	0.01	0.65
	Low	2	0.06	0.01	0.70	0.06	0.00	0.68
	Unburned	4	0.02	0.00	0.25	0.02	0.00	0.24
<b>Golden-mantled Ground Squirrel</b>	High	1	0.01	0.00	0.07	0.05	0.01	0.30
	Low	3	0.02	0.00	0.12	0.11	0.02	0.50
	Unburned	56	0.06	0.02	0.21	0.30	0.10	0.91
<b>Hairy Woodpecker</b>	High	51	0.49	0.19	1.26	0.53	0.19	1.50
	Low	22	0.41	0.14	1.21	0.45	0.14	1.44
	Unburned	38	0.13	0.05	0.35	0.14	0.05	0.41
Kaibab Squirrel	High	0	0.00	0.00	0.60	0.01	0.00	1.30
	Low	4	0.02	0.00	3.26	0.10	0.00	5.66
	Unburned	20	0.01	0.00	1.56	0.06	0.00	2.71
Mourning Dove	High	3	0.05	0.00	0.57	0.13	0.01	1.78
	Low	3	0.01	0.00	0.19	0.03	0.00	0.58
	Unburned	3	0.01	0.00	0.19	0.04	0.00	0.58
<b>Northern Flicker</b>	High	52	0.42	0.21	0.84	0.42	0.20	0.90
	Low	23	0.23	0.10	0.52	0.23	0.09	0.56
	Unburned	141	0.29	0.15	0.58	0.30	0.14	0.62
<b>Red Squirrel</b>	High	1	0.01	0.00	0.06	0.00	0.00	0.01
	Low	10	0.66	0.20	2.11	0.10	0.03	0.32
	Unburned	83	0.52	0.20	1.34	0.08	0.03	0.20
<b>Steller's Jay</b>	High	8	0.05	0.01	0.19	0.08	0.02	0.33
	Low	6	0.05	0.01	0.22	0.09	0.02	0.36
	Unburned	45	0.05	0.01	0.20	0.10	0.03	0.34
Three-toed Woodpecker	High	4	0.02	0.00	1.06	0.00	0.00	0.35
	Low	0	0.02	0.00	1.54	0.00	0.00	0.52
	Unburned	0	0.01	0.00	0.73	0.00	0.00	0.24
Williamson's Sapsucker	High	0	0.00	0.00	0.00	0.00	0.00	0.00
	Low	1	0.12	0.00	15.06	0.02	0.00	3.24
	Unburned	12	0.18	0.00	18.37	0.03	0.00	3.96

Appendix B (continue). 2008 Density (D) of birds and mammals per hectare by forest type and fire severity, with sample size (n) and 95% confidence limits (LCL / UCL). Species (**bold**) with  $n \geq 60$  detections per year were used to develop models to best predict species abundance.

## APPENDIX B

2009			Mixed Conifer			Ponderosa Pine		
Species common name	Severity	n	D	LCL	UCL	D	LCL	UCL
<b>American Robin</b>	High	79	0.56	0.23	1.36	0.24	0.08	0.69
	Low	12	0.29	0.09	0.90	0.12	0.03	0.45
	Unburned	53	0.43	0.20	0.91	0.18	0.07	0.46
<b>Chipmunk spp.</b>	High	11	0.20	0.05	0.72	0.65	0.21	2.03
	Low	16	0.20	0.06	0.70	0.66	0.22	1.98
	Unburned	76	0.24	0.09	0.62	0.77	0.34	1.75
Clark's Nutcracker	High	7	0.04	0.00	0.32	0.03	0.00	0.33
	Low	2	0.01	0.00	0.18	0.01	0.00	0.18
	Unburned	7	0.02	0.00	0.15	0.02	0.00	0.16
Downy Woodpecker	High	6	0.05	0.00	0.53	0.05	0.00	0.51
	Low	1	0.05	0.00	0.55	0.04	0.00	0.53
	Unburned	5	0.02	0.00	0.20	0.02	0.00	0.19
<b>Golden-mantled Ground Squirrel</b>	High	1	0.01	0.00	0.11	0.07	0.01	0.47
	Low	7	0.04	0.01	0.19	0.18	0.04	0.78
	Unburned	98	0.10	0.03	0.34	0.48	0.16	1.41
<b>Hairy Woodpecker</b>	High	39	0.49	0.19	1.25	0.53	0.18	1.48
	Low	33	0.41	0.14	1.20	0.44	0.14	1.42
	Unburned	39	0.13	0.05	0.34	0.14	0.05	0.41
Kaibab Squirrel	High	1	0.00	0.00	0.35	0.01	0.00	1.05
	Low	1	0.01	0.00	1.35	0.04	0.00	4.08
	Unburned	8	0.00	0.00	0.65	0.03	0.00	1.96
Mourning Dove	High	14	0.07	0.01	0.76	0.18	0.01	2.32
	Low	4	0.01	0.00	0.25	0.04	0.00	0.76
	Unburned	7	0.02	0.00	0.25	0.05	0.00	0.76
<b>Northern Flicker</b>	High	102	0.58	0.29	1.15	0.58	0.27	1.24
	Low	26	0.32	0.14	0.71	0.32	0.13	0.76
	Unburned	170	0.40	0.20	0.79	0.41	0.19	0.85
<b>Red Squirrel</b>	High	2	0.02	0.00	0.09	0.00	0.00	0.02
	Low	29	1.06	0.32	3.30	0.17	0.05	0.54
	Unburned	123	0.84	0.32	2.10	0.14	0.05	0.35
<b>Steller's Jay</b>	High	18	0.06	0.01	0.25	0.10	0.03	0.41
	Low	10	0.06	0.01	0.27	0.11	0.03	0.46
	Unburned	50	0.07	0.02	0.26	0.12	0.03	0.43
Three-toed Woodpecker	High	3	0.03	0.00	1.62	0.01	0.00	0.61
	Low	0	0.05	0.00	2.34	0.02	0.00	0.85
	Unburned	4	0.03	0.00	1.11	0.01	0.00	0.42
Williamson's Sapsucker	High	1	0.00	0.00	0.00	0.00	0.00	0.00
	Low	1	0.09	0.00	11.52	0.01	0.00	2.30
	Unburned	7	0.12	0.00	14.05	0.02	0.00	2.81

Appendix B (continue). 2009 Density (D) of birds and mammals per hectare by forest type and fire severity, with sample size (n) and 95% confidence limits (LCL / UCL). Species (**bold**) with  $n \geq 60$  detections per year were used to develop models to best predict species abundance.

## APPENDIX B

2010			Mixed Conifer			Ponderosa Pine		
Species common name	Severity	n	D	LCL	UCL	D	LCL	UCL
<b>American Robin</b>	High	29	0.51	0.20	1.23	0.21	0.07	0.62
	Low	10	0.26	0.08	0.81	0.11	0.03	0.41
	Unburned	92	0.39	0.18	0.82	0.16	0.06	0.41
<b>Chipmunk spp.</b>	High	20	0.25	0.07	0.91	0.82	0.26	2.51
	Low	19	0.25	0.07	0.89	0.83	0.27	2.45
	Unburned	92	0.30	0.11	0.78	0.97	0.43	2.17
Clark's Nutcracker	High	3	0.02	0.00	0.20	0.02	0.00	0.20
	Low	2	0.01	0.00	0.11	0.01	0.00	0.11
	Unburned	3	0.01	0.00	0.10	0.01	0.00	0.10
Downy Woodpecker	High	0	0.05	0.00	0.44	0.05	0.00	0.44
	Low	2	0.04	0.00	0.43	0.03	0.00	0.41
	Unburned	7	0.01	0.00	0.15	0.01	0.00	0.15
<b>Golden-mantled Ground Squirrel</b>	High	5	0.02	0.00	0.12	0.08	0.01	0.50
	Low	12	0.04	0.01	0.21	0.19	0.04	0.84
	Unburned	106	0.10	0.03	0.37	0.52	0.17	1.51
<b>Hairy Woodpecker</b>	High	21	0.23	0.08	0.62	0.25	0.08	0.73
	Low	13	0.19	0.06	0.59	0.21	0.06	0.70
	Unburned	20	0.06	0.02	0.17	0.07	0.02	0.20
Kaibab Squirrel	High	0	0.00	0.00	0.00	0.01	0.00	0.00
	Low	4	0.01	0.00	2.20	0.07	0.00	5.00
	Unburned	11	0.01	0.00	1.05	0.04	0.00	2.40
Mourning Dove	High	0	0.04	0.00	0.23	0.04	0.00	0.81
	Low	0	0.01	0.00	0.08	0.01	0.00	0.27
	Unburned	7	0.01	0.00	0.08	0.01	0.00	0.27
<b>Northern Flicker</b>	High	54	0.42	0.21	0.85	0.43	0.20	0.91
	Low	19	0.23	0.10	0.53	0.23	0.10	0.56
	Unburned	145	0.30	0.15	0.58	0.30	0.14	0.63
<b>Red Squirrel</b>	High	1	0.02	0.00	0.10	0.00	0.00	0.02
	Low	32	1.05	0.32	3.31	0.17	0.05	0.54
	Unburned	118	0.84	0.32	2.11	0.14	0.05	0.34
<b>Steller's Jay</b>	High	2	0.04	0.01	0.19	0.08	0.02	0.31
	Low	9	0.05	0.01	0.21	0.08	0.02	0.35
	Unburned	45	0.05	0.01	0.19	0.09	0.03	0.32
Three-toed Woodpecker	High	1	0.03	0.00	1.62	0.01	0.00	0.61
	Low	2	0.05	0.00	2.34	0.02	0.00	0.89
	Unburned	4	0.03	0.00	1.11	0.01	0.00	0.42
Williamson's Sapsucker	High	0	0.00	0.00	0.00	0.00	0.00	0.00
	Low	0	0.08	0.00	9.55	0.00	0.00	2.01
	Unburned	7	0.10	0.00	11.41	0.02	0.00	2.19

Appendix B (continue). 2010 Density (D) of birds and mammals per hectare by forest type and fire severity, with sample size (n) and 95% confidence limits (LCL / UCL). Species (bold) with  $n \geq 60$  detections per year were used to develop models to best predict species abundance.

APPENDIX C

	Forest Type	Fire Severity	Fire Severity Comparison		
			High - Low	High – No Burn	Low – No Burn
Tree Density	Mixed-Conifer	p <.0001	p <.0001	p <.0001	<b>p =.7615</b>
	Ponderosa Pine	p <.0001	p <.0001	p <.0001	<b>p =.7671</b>
Tree Density >20 cm dbh	Mixed-Conifer	p <.0001	p <.0001	p <.0001	<b>p =.9634</b>
	Ponderosa Pine	p <.0001	p <.0001	p <.0001	<b>p =.9526</b>
Snag Density	Mixed-Conifer	p =.0066	p =.0275	p =.0109	<b>p =.7916</b>
	Ponderosa Pine	p <.0001	<b>p =.3512</b>	p <.0002	p =.0077
Snag Density >20 cm dbh	Mixed-Conifer	p =.0054	<b>p =.1953</b>	p =.0039	<b>p =.7616</b>
	Ponderosa Pine	p <.0001	p =.0382	p <.0001	p <.0003
Canopy Cover	Mixed-Conifer	p <.0001	p <.0001	p <.0001	<b>p =.9970</b>
	Ponderosa Pine	p <.0169	p =.0103	p =.0162	<b>p =.8823</b>
Ground Cover	Mixed-Conifer	p <.0001	p <.0002	p <.0001	<b>p =.9969</b>
	Ponderosa Pine	<b>p =.6672</b>	<b>p =.6544</b>	<b>p =.8787</b>	<b>p =.7749</b>
Cone Productivity	Mixed-Conifer	p <.0001	p =.0220	p <.0001	<b>p =.2860</b>
	Ponderosa Pine	p <.0001	p <.0001	p <.0001	<b>p =.9554</b>
Sapling Counts	Mixed-Conifer	p <.0001	p <.0001	p <.0001	p =.0012
	Ponderosa Pine	p <.0001	p =.0016	<b>p =.8439</b>	p <.0001

Appendix C. Summary of p-values for a comparison of the means of forest metrics: by fire severity. Cover, productivity, and count comparisons are averaged across 2007-2010. P-values showing that difference between means were not significant at  $\alpha \leq 0.05$  are **bold**.

APPENDIX D

Mixed Conifer				Ponderosa Pine			
Forb	Unburned	Low	High	Forb	Unburned	Low	High
<i>Achillea millefolium</i>			X	<i>Ambrosia spp.</i>	X		
<i>Antennaria spp.</i>	X	X	X	<i>Antennaria spp.</i>	X	X	
<i>Chenopodium fremontii</i>		X	X	<i>Chenopodium fremontii</i>	X	X	X
<i>Cirsium spp.</i>			X	<i>Cirsium spp.</i>	X	X	X
<i>Colubrina Californica</i>			X	<i>Compositae</i>	X	X	
<i>Compositae</i>	X	X	X	<i>Erigeron formosissimus</i>			X
<i>Erigeron formosissimus</i>			X	<i>Erigeron macranthus</i>	X		
<i>Erigeron macranthus</i>	X	X		<i>Eriogonum racemosum</i>	X		X
<i>Fragaria vesca</i>	X			<i>Fragaria vesca</i>	X		
<i>Garryaceae</i>			X	<i>Gnaphalium arizonicum</i>	X		
<i>Goodyera oblongifolia</i>	X			<i>Helianthus</i>	X		
<i>Helianthus</i>			X	<i>Lactuca serriola</i>			X
<i>Leguminosea</i>	X			<i>Lotus wrightii</i>	X	X	X
<i>Lotus wrightii</i>	X	X		<i>Lupinus spp.</i>	X		
<i>Lupinus spp.</i>	X	X	X	<i>Oenothera spp.</i>	X		
<i>Portulacaceae</i>	X			<i>Rhamnaceae</i>			X
<i>Ranunculus</i>			X	<i>Solidago spp.</i>	X		X
<i>Rubus strigosus</i>	X			<i>Verbascum thapsus</i>	X		
<i>Senecio multilobatus</i>			X				
<i>Solidago spp.</i>	X	X					
Shrub				Shrub			
<i>Berberis repens</i>	X	X	X	<i>Berberis repens</i>	X	X	X
<i>Ceanothus fendleri</i>	X	X	X	<i>Ceanothus fendleri</i>	X	X	X
<i>Juniperus communis</i>	X			<i>Ericameria nauseosa</i>	X	X	X
<i>Ribes spp.</i>			X	<i>Juniperus communis</i>	X		
<i>Robinia neomexicana</i>			X	<i>Ribes spp.</i>	X	X	
<i>Sambucus spp.</i>			X	<i>Robinia neomexicana</i>	X	X	X
				<i>Rosacea arizonica</i>	X		
				<i>Sambucus spp.</i>	X		

Appendix D. Dominant Forbs and Shrubs found on transects from 2007 -2010.